

Global Warming of 1.5°C

An IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty.

Panmao Zhai
Co-Chair, IPCC Working Group I

Siem Reap, 27 May 2019

Global Warming of 1.5°C

An IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty.

The report in numbers

91 Authors from 40 Countries

133 Contributing authors

6000 Studies

1 113 Reviewers

42 001 Comments

Global warming of 1.5°C (SR1.5)

Chapter 1: Framing and context (integration WGI-WGII-III)

Chapter 2: Mitigation pathways compatible with 1.5°C in the context of sustainable development (integration WGI-WGIII, pathways)

Chapter 3: Impacts of 1.5°C global warming on natural and human systems (integration WGI-WGII, global – regional)

Chapter 4: Strengthening and implementing the global response to the threat of climate change (systems transitions, behaviour, dimensions of feasibility)

Chapter 5: Sustainable development, poverty eradication and reducing inequalities (ethics, equity, societal transformation, SDGs)

Understanding Global Warming of 1.5°C



Where are we?

Since pre-industrial times, human activities have caused approximately 1.0°C of global warming.

- Already seeing consequences for people, nature and livelihoods
- At current rate, would reach 1.5°C between 2030 and 2052
- Past emissions alone do not commit the world to 1.5°C


Ashley Cooper / Aurora Photos

ipcc

INTERGOVERNMENTAL PANEL ON climate change





Cumulative emissions of CO₂ and future non-CO₂ radiative forcing determine the probability of limiting warming to 1.5°C




Maximum temperature rise is determined by cumulative net CO₂ emissions and net non-CO₂ radiative forcing due to methane, nitrous oxide, aerosols and other anthropogenic forcing agents

Regional warming in the decade 2006-2015 relative to preindustrial


Annual average warming



December-January-February



June-July-August




Projected Climate Change, Potential Impacts and Associated Risks


Spatial patterns of changes in mean temperature

Global warming of 1.5°C

2°C




Temperature (°C)



Spatial patterns of changes in mean temperature and precipitation

Global warming of 1.5°C

2°C






26 CMIP5 models; hatching : 66% model agreement

Spatial patterns of changes in mean temperature and precipitation

Global warming of 1.5°C





2°C

Difference





Temperature (°C)

Temperature (°C)



Precipitation (%)

Precipitation (%)




Spatial patterns of changes in extreme temperature


Global warming of 1.5°C

2°C


*Number of hot days
(days)*



*Temperature of
hottest days (°C)*



*Temperature of
coldest nights (°C)*




Spatial patterns of changes in extreme temperature and precipitation


Global warming of 1.5°C

2°C


Number of hot days (days)




Temperature of hottest days (°C)



Temperature of coldest nights (°C)



Extreme precipitation (%)




Spatial patterns of changes in extreme temperature and precipitation

Global warming of 1.5°C

2°C

Difference


Number of hot days (days)




Temperature of hottest days (°C)



Temperature of coldest nights (°C)




Extreme precipitation (%)



At 1.5°C compared to 2°C

- Less extreme weather where people live, including extreme heat and rainfall
- By 2100, global mean sea level rise will be around 10 cm lower but may continue to rise for centuries
- Fewer people exposed to sea level rise



Jason Florio / Aurora Photos

ipcc

INTERGOVERNMENTAL PANEL ON climate change



Emergence and intensity of regional climate change hot spots

Arctic summer sea-ice

- *L* maintained; 50% or higher risk to be ice free; *VL* to be ice free
- Habitat (polar bear, whales, seals, sea birds) : losses; losses; critical losses
- Arctic fisheries : benefits; benefits; benefits

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

Emergence and intensity of regional climate change hot spots

Arctic land regions

- Cold extreme: warm up to 4.5° C (HC); warm up to 8° C (HC); VL drastic warming
- Tundra : L biome shifts; L more shifts; drastic biome shift possible (LC)
- Permafrost : L 17-44% reduction; L larger (28-53%); potential for collapse (LC)
- Boreal forest : increased mortality at S. boundary (MC); further (MC); potential dieback (LC)

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

Emergence and intensity of regional climate change hot spots

Alpine regions

- Biomes : *L* severe shift; *L* even more severe; *L* critical

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

Emergence and intensity of regional climate change hot spots

Mediterranean

- Extreme drought: increase probability(*MC*); robust increase(*MC*); robust and large increase(*MC*)
- Runoff decrease: about 9% (*MC*); about 17% (*MC*); substantial reductions (*MC*)
- Water deficit: risk (*MC*); higher risks (*MC*); very high risks (*MC*)

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

Emergence and intensity of regional climate change hot spots

Tropics

- # hot days and nights, heatwaves: **increases (HC)**; largest increase; **oppressive, VL health impact**
- Livestock heat stress : **increased**; onset of persistent (MC); **L persistent**
- Crop yields: **risks**; extensive risks (W. Africa, SE Asia, S. America); **VL substantial reductions**
- Rainforests : **reduced biomass**; larger reductions; **reduced extent (Central America), potential forest dieback (MC)**

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

Emergence and intensity of regional climate change hot spots

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

Southeast Asia

- ↗ flooding related to sea-level rise: risks; higher risks (*MC*); substantial increases in risk
- Asian monsoon : *LC*; *LC*; *L* increase in precipitation intensity
- Heavy precipitation: increase; stronger increase (*MC*); substantial increase
- Crop yield reductions: -; one third decline in per capita (*MC*); substantial reduction

Emergence and intensity of regional climate change hot spots

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

West African and the Sahel

- Monsoon : uncertain ; uncertain ; strengthening (*LC*)
- Hot nights, longer, more frequent heat waves: *L* ↗; *L* further ↗; *VL* substantial ↗
- ↘ in maize and sorghum production: *L*, about 40% ↘ suitable area; *L* larger; major regional food insecurities (*MC*)
- Undernutrition risks : increased; higher; high

Emergence and intensity of regional climate change hot spots

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

Southern Africa

- Water availability: reductions (*MC*); larger reductions (*MC*); large reductions (*MC*)
- # of hot nights and ↗ heat waves : increases (*HC*); further increase (*HC*); drastic increase (*HC*)
- Increased mortality from heat-waves: high risks; higher risks (*HC*);
substantial impact on health and mortality (*HC*)
- Undernutrition / dryland agriculture and livestock: high risk; higher risk (*HC*); very high risks

Emergence and intensity of regional climate change hot spots

Warming of 1.5° C or less

Warming of 1.5°C-2° C

Warming > 2° C

L, likely

VL, very likely

LC, low confidence

MC, medium confidence

HC, high confidence

Small islands:

- Inundation risk : land exposed; tens of thousands displaced ; substantial, widespread impacts
- Coastal flooding: risks; high risks ; substantial and widespread impacts
- Fresh water stress : increased; projected aridity; substantial and widespread impacts
- # of warm days : increase; further increase (70 warm days/year), persistent heat stress in cattle ; persistent heat stress
- Loss of coral reefs: 70-90%; most coral reefs ; loss of most coral reefs (VL)

THANK YOU FOR YOUR ATTENTION!

For more information:

Website: <http://ipcc.ch/>

IPCC Secretariat: ipcc-sec@wmo.int

IPCC Press Office: ipcc-media@wmo.int

Find us on:



ipcc

INTERGOVERNMENTAL PANEL ON climate change

