

1
2 **Annex III: Tables of historical and projected well-mixed greenhouse gas mixing ratios**
3 **and effective radiative forcing of all climate forcers**

10 **Coordinating Lead Authors:** Frank Dentener (EU, The Netherlands), Bradley Hall (USA), Chris Smith
11 (UK)

12
13 **Lead Authors:** Jinho Ahn (Republic of Korea), William Collins (UK), Christopher Jones (UK), Malte
14 Meinshausen (Australia/Germany),

15 **Contributing Authors:**

16 Ralph Keeling (USA), Paul Krummel (Australia), Jens Mühle (USA), Zebedee Nicholls (Australia), Isobel
17 Simpson (USA)

18 **Review Editors:**

19 Text in AR6 Body Text

20 **Date of Draft:**

21 3 May 2021

22 **Notes:**

23 TSU compiled version

1 AIII.1 Introduction

2 Annex III presents, in tabulated form, data related to historical and projected changes in greenhouse gas
 3 (GHG) mixing ratios and effective radiative forcing (ERF) of all climate forcers as assessed and used
 4 throughout Chapters 1-7. Where available 90 % confidence level uncertainties on observed GHG mixing
 5 ratios are given. For each species, the abundance is given as dry air mole fraction: ppm = micromoles per
 6 mole (10^{-6}); ppb = nanomoles per mole (10^{-9}); and ppt = picomoles per mole (10^{-12}). ERF is given in W m^{-2}
 7 for CO_2 , N_2O , and CH_4 and mW m^{-2} for other components.

8
 9 Pre-instrumental mixing ratios are estimated from ice core and firn air records that are described in Ahn et
 10 al., 2012; Bauska et al., 2015; Flückiger et al., 1999; Machida et al., 1995; Meinshausen et al., 2017;
 11 Mitchell et al., 2013; Rubino et al., 2020; Ryu et al., 2020; Siegenthaler et al., 2005; Sowers, 2001.
 12 Observed (instrumental) mixing ratios are described in Dlugokencky et al., 2011; Hall et al., 2011; Laube et
 13 al., 2016; Masarie and Tans, 2004; Montzka et al., 2009; Naus et al., 2019; Prinn et al., 2018; Rigby et al.,
 14 2014; Trudinger et al., 2004; Worton et al., 2006; Adcock et al., 2018; Droste et al., 2020; Leedham Elvidge
 15 et al., 2018; Mühlé et al., 2019; Simmonds et al., 2017.

16
 17 Projected concentrations for the 5 core scenarios discussed in the report (Section 1.6.1) are from (Gidden et
 18 al., 2019; Meinshausen et al., 2017, 2020; Velders et al., 2015). These scenarios span a wide range of
 19 plausible societal and climatic futures from potentially below 1.5°C best-estimate warming to over 4°C
 20 warming by 2100 (Section 4.3.4). Computational methods and assumptions to calculate historical and
 21 projected ERF are described in Chapter 7 and detailed information can be found in Chapter 7 Supplementary
 22 Material 7.SM.1.3 and 7.SM.1.4.

23
 24 Extended datasets and further auxiliary data are made available via <https://zenodo.org/xxxx>

25
 26 **Chemical Abbreviations and Symbols of components regulated under the Kyoto¹¹ and Montreal
 Protocols.**

| | | |
|---------------------------|---|--------------------|
| CO_2 | carbon dioxide | Kyoto |
| CH_4 | methane | Kyoto |
| N_2O | nitrous oxide | Kyoto |
| HFC | hydrofluorocarbon (a class of compounds: HFC-32, HFC-134a, ...) | Kyoto, Montreal |
| PFC | perfluorocarbon (a class of compounds: CF_4 , C_2F_6 , C_4F_{10} , ...) | Kyoto |
| SF_6 | sulphur hexafluoride | Kyoto |
| NF_3 | nitrogen trifluoride | Kyoto |
| CFC | chlorofluorocarbon (a class of compounds: CFCl_3 , CF_2Cl_2 , ...) | Montreal |
| HCFC | hydrochlorofluorocarbon (a class of compounds: HCFC-22, HCFC-141b, ...) | Montreal |
| CCl_4 | carbon tetrachloride | Montreal |
| CH_3CCl_3 | methyl chloroform | Montreal |
| Halons | Bromo(chloro)fluorocarbon (a class of compounds: CF_2ClBr “Halon-1211”; CBrF_3 “Halon-1301”; $\text{C}_2\text{Br}_2\text{F}_4$ “Halon-2402”) | Montreal |

27
 28 ¹¹ The Kyoto protocol (1 December 1997-31 December 2020) regulated a basket of 6 GHGs. The term Kyoto gases is
 widely used in the scientific literature

1 **List of Tables:**

| | |
|---------------|--|
| Table AIII.1 | Historical Abundances of greenhouse gases |
| Table AIII.1a | Historical abundances and ERF(Wm^{-2}) for $\text{CO}_2(\text{ppm})$, $\text{CH}_4(\text{ppb})$, and $\text{N}_2\text{O}(\text{ppb})$ |
| Table AIII.1b | Historical abundances (ppt) and ERF (mWm^{-2}) of NF_3 , SF_6 , and PFCs |
| Table AIII.1c | Historical abundances (ppt) and ERF [mWm^{-2}] of HFCs |
| Table AIII.1d | Historical abundances (ppt) and ERF [mWm^{-2}] of HCFCs |
| Table AIII.1e | Historical abundances (ppt) and ERF [mWm^{-2}] of CFCs |
| Table AIII.1f | Historical abundances (ppt) and ERF [mWm^{-2}] of CH_3CCl_3 , CCl_4 , CH_3Br , CHCl_3 and Halons. |
| Table AIII.2 | Future abundances of CO_2 , CH_4 , and N_2O for selected SSP scenarios [2020-2500] |
| Table AIII.3 | Effective Radiative Forcing (Wm^{-2}) time series of all climate forcers from 1750-2019 |
| Table AIII.4a | Effective Radiative Forcing (Wm^{-2}) time series of all climate forcers for SSP1-19 |
| Table AIII.4b | Effective Radiative Forcing (Wm^{-2}) time series of all climate forcers for SSP1-26 |
| Table AIII.4c | Effective Radiative Forcing (Wm^{-2}) time series of all climate forcers for SSP2-45 |
| Table AIII.4d | Effective Radiative Forcing (Wm^{-2}) time series of all climate forcers for SSP3-70 |
| Table AIII.4e | Effective Radiative Forcing (Wm^{-2}) time series of all climate forcers for SSP5-85 |
| Table AIII.4f | Effective Radiative Forcing (mWm^{-2}) time series of halogen components for selected scenarios and Kyoto and Montreal gases |
| Table AIII.5 | Total Anthropogenic and natural ERF relative to 1750 assessed in AR5 (RCP scenarios) and AR6 (SSP scenarios). |

2
3

1 **Table AIII.1a:** Historical abundances and ERF(Wm⁻²) for CO₂(ppm), CH₄(ppb), and N₂O(ppb)

| Year | CO ₂ | CH ₄ | N ₂ O | Year | CO ₂ | CH ₄ | N ₂ O | Year | CO ₂ | CH ₄ | N ₂ O |
|-------------------|-----------------|-----------------|------------------|------|--------------------|-------------------|--------------------|------------------------|-----------------|-----------------|------------------|
| 1750 ¹ | 278.3 | 729 | 270.1 | 1968 | 322.5 | 1372 | 295.2 | 1995 | 360.0 | 1748 | 311.4 |
| 1850 ¹ | 285.5 | 808 | 272.1 | 1969 | 323.4 | 1389 | 295.6 | 1996 | 361.8 | 1750 | 312.3 |
| 1860 ² | 286.8 | 822 | 273.2 | 1970 | 324.9 | 1411 | 296.0 | 1997 | 362.9 | 1754 | 313.1 |
| 1870 | 288.4 | 852 | 274.9 | 1971 | 325.5 | 1431 | 296.5 | 1998 | 365.5 | 1764 | 313.9 |
| 1880 | 290.4 | 868 | 276.6 | 1972 | 327.4 | 1449 | 296.9 | 1999 | 367.6 | 1772 | 314.9 |
| 1890 | 293.3 | 896 | 277.6 | 1973 | 330.0 | 1463 | 297.3 | 2000 | 368.8 | 1773 | 315.9 |
| 1900 | 296.4 | 925 | 278.9 | 1974 | 330.8 | 1476 | 297.8 | 2001 | 370.4 | 1772 | 316.6 |
| 1905 | 298.0 | 947 | 280.2 | 1975 | 330.9 | 1492 | 298.3 | 2002 | 372.4 | 1773 | 317.3 |
| 1910 | 300.0 | 974 | 281.8 | 1976 | 331.6 | 1509 | 298.8 | 2003 | 375.0 | 1777 | 318.0 |
| 1915 | 302.5 | 991 | 283.6 | 1977 | 333.4 | 1528 | 299.3 | 2004 | 376.8 | 1776 | 318.6 |
| 1920 | 304.8 | 1025 | 284.5 | 1978 | 335.0 | 1547 | 299.8 | 2005 | 378.8 | 1774 | 319.3 |
| 1925 | 306.3 | 1052 | 285.3 | 1979 | 336.6 | 1566 | 300.4 | 2006 | 381.0 | 1774 | 320.2 |
| 1930 | 307.1 | 1072 | 285.6 | 1980 | 338.8 ³ | 1585 | 301.1 ³ | 2007 | 382.7 | 1781 | 320.9 |
| 1935 | 308.6 | 1097 | 286.3 | 1981 | 340.0 | 1603 | 301.9 | 2008 | 384.8 | 1788 | 321.8 |
| 1940 | 311.7 | 1120 | 287.3 | 1982 | 340.8 | 1619 | 303.1 | 2009 | 386.3 | 1793 | 322.6 |
| 1945 | 312.7 | 1139 | 289.0 | 1983 | 342.4 | 1633 | 303.7 | 2010 | 388.6 | 1798 | 323.4 |
| 1950 | 313.1 | 1164 | 289.5 | 1984 | 344.0 | 1645 ³ | 304.3 | 2011 | 390.5 | 1803 | 324.4 |
| 1955 | 314.6 | 1207 | 290.7 | 1985 | 345.5 | 1657 | 304.9 | 2012 | 392.5 | 1808 | 325.3 |
| 1960 | 316.8 | 1264 | 292.1 | 1986 | 346.9 | 1670 | 305.8 | 2013 | 395.2 | 1814 | 326.2 |
| 1961 | 317.5 | 1269 | 292.5 | 1987 | 348.6 | 1680 | 306.0 | 2014 | 397.1 | 1823 | 327.4 |
| 1962 | 318.2 | 1282 | 292.8 | 1988 | 351.2 | 1693 | 306.7 | 2015 | 399.4 | 1834 | 328.3 |
| 1963 | 318.8 | 1301 | 293.2 | 1989 | 352.8 | 1707 | 307.8 | 2016 | 402.9 | 1842 | 329.1 |
| 1964 | 319.5 | 1317 | 293.6 | 1991 | 354.0 | 1714 | 308.7 | 2017 | 405.0 | 1849 | 330.0 |
| 1965 | 320.0 | 1331 | 293.9 | 1992 | 355.3 | 1728 | 309.4 | 2018 | 407.4 | 1858 | 331.2 |
| 1966 | 321.0 | 1342 | 294.4 | 1993 | 356.0 | 1735 | 309.9 | 2019 | 409.9 | 1866 | 332.1 |
| 1967 | 321.6 | 1354 | 294.8 | 1994 | 356.7 | 1737 | 310.3 | ERF⁴ | 2.16 | 0.54 | 0.21 |

3 **Notes:** ¹ 1750/1850 CO₂, CH₄, N₂O from multiple ice cores assessed in Chapter 2. Uncertainties (90 % CI) for 1750 are
4 2.9 ppm, 9.4 ppb and 6.0 ppb for CO₂, CH₄ and N₂O, respectively. Uncertainties for 1850 are 2.1 ppm, 13.8 ppb and 5.7
5 ppb, based on variations of ice cores. ² Mixing ratios from 1851- 1980/1984 are updated from the CMIP6 (Meinshausen
6 et al., 2017) dataset, using a linear time-dependent offset correction function. ³ CO₂ from NOAA network; CH₄, N₂O
7 from merged NOAA and AGAGE networks. Uncertainties (90 % CI) in 2019, derived from multiple global networks,
8 are 0.36 ppm, 3.3 ppb and 0.4 ppb for CO₂, CH₄ and N₂O, respectively, and do not include estimates of analytical
9 accuracy. Uncertainties for other years may differ. ⁴ ERF(2019-1750) from Chapter 7.

Table AIII.1b: Historical abundances (ppt) and ERF (mWm⁻²) of NF₃, SF₆, and perfluorocarbons

| Year | NF ₃ | SF ₆ | SO ₂ F ₂ | CF ₄ | C ₂ F ₆ | C ₃ F ₈ | c-C ₄ F ₈ | n-C ₄ F ₁₀ | n-C ₅ F ₁₂ | n-C ₆ F ₁₄ | i-C ₆ F ₁₄ | C ₇ F ₁₆ | C ₈ F ₁₈ |
|-------------|-----------------|-----------------|--------------------------------|-----------------|-------------------------------|-------------------------------|---------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|--------------------------------|--------------------------------|
| 1750 | 0.00 | 0.00 | 0.00 | 34.05 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1850 | 0.00 | 0.00 | 0.00 | 34.1 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1900 | 0.00 | 0.00 | 0.00 | 34.1 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1910 | 0.00 | 0.00 | 0.00 | 34.1 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1920 | 0.00 | 0.00 | 0.00 | 34.4 | 0.04 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1930 | 0.00 | 0.00 | 0.00 | 34.9 | 0.10 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1940 | 0.00 | 0.00 | 0.00 | 35.8 | 0.19 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1950 | 0.00 | 0.00 | 0.00 | 38.0 | 0.40 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1960 | 0.00 | 0.09 | 0.00 | 40.1 | 0.51 | 0.02 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1970 | 0.00 | 0.32 | 0.01 | 43.4 | 0.62 | 0.03 | 0.14 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1980 | 0.00 | 0.86 | 0.35 | 53.5 | 1.22 | 0.05 | 0.38 | 0.02 | 0.02 | 0.01 | 0.00 | 0.01 | 0.01 |
| 1990 | 0.01 | 2.35 | 0.68 | 63.8 | 2.07 | 0.12 | 0.76 | 0.07 | 0.05 | 0.03 | 0.015 | 0.03 | 0.02 |
| 2000 | 0.17 | 4.56 | 1.07 | 71.5 | 3.11 | 0.28 | 0.98 | 0.13 | 0.10 | 0.14 | 0.038 | 0.07 | 0.06 |
| 2010 | 0.73 | 7.01 | 1.63 | 78.3 | 4.09 | 0.54 | 1.26 | 0.17 | 0.12 | 0.21 | 0.055 | 0.10 | 0.09 |
| 2015 | 1.30 | 8.57 | 2.11 | 81.9 | 4.49 | 0.62 | 1.50 | 0.19 | 0.14 | 0.22 | 0.062 | 0.11 | n.a. |
| 2019 | 2.05 | 9.95 | 2.50 | 85.5 | 4.85 | 0.68 | 1.75 | n.a. | n.a. | n.a. | n.a. | n.a. | n.a. |
| <i>Unc.</i> | <i>0.03</i> | <i>0.03</i> | <i>0.05</i> | <i>0.2</i> | <i>0.05</i> | <i>0.01</i> | <i>0.06</i> | <i>n.a.</i> | <i>n.a.</i> | <i>n.a.</i> | <i>n.a.</i> | <i>n.a.</i> | <i>n.a.</i> |
| <i>ERF</i> | <i>0.4</i> | <i>5.6</i> | <i>0.5</i> | <i>5.1</i> | <i>1.3</i> | <i>0.2</i> | <i>0.5</i> | <i>0.1</i> | <i>0.1</i> | <i>0.1</i> | <i>0.0¹</i> | <i>0.1</i> | <i>0.1</i> |

Notes: Data merged from AR5 (1750;1850); CMIP6 compilation by Meinshausen et al. (2017) until ca. 1995, and data directly taken from merged AGAGE and NOAA networks, depending on date of availability. Perfluorocarbons from CMIP6 dataset (Meinshausen et al., 2017) or estimated from Droste et al. (2020), with CMIP6 n-C₆F₁₄ and C₇F₁₆ scaled to account for calibration changes in Droste et al. (2020). Uncertainties pertain to 2019, derived from observations made by global networks and literature, and do not include estimates of analytical accuracy. Uncertainties are not available for c-C₄F₈, n-C₄F₁₀, n-C₅F₁₂, n-C₆F₁₄, i-C₆F₁₄, C₇F₁₆, C₈F₁₈. ERF(2019-1750) from Chapter 7, except for c-C₄F₈, n-C₄F₁₀, n-C₅F₁₂, n-C₆F₁₄, i-C₆F₁₄, C₇F₁₆ use 2015 abundances and C₈F₁₈ which uses 2014 abundance. ¹below <0.5 mWm⁻²

Table AIII.1c: Historical abundances (ppt) and ERF [mWm⁻²] of HFCs

| Year | HFC-134a | HFC-23 | HFC-32 | HFC-125 | HFC-143a | HFC-152a | HFC-227ea | HFC-236fa | HFC-245fa | HFC-365mfc | HFC-43-10mee |
|-------------|-------------|------------|------------|------------|------------|------------|-------------|-------------|-------------|-------------|--------------|
| 1750 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1850 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1900 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1910 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1920 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1930 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1940 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1950 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1960 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1970 | 0.0 | 1.8 | 0.0 | 0. | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1980 | 0.2 | 3.9 | 0.0 | 0.1 | 0.1 | 0.0 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1990 | 0.5 | 8.3 | 0.1 | 0.1 | 0.5 | 0.2 | 0.01 | 0.00 | 0.00 | 0.01 | 0.00 |
| 2000 | 14.2 | 15.2 | 0.2 | 1.5 | 2.5 | 1.6 | 0.11 | 0.02 | 0.02 | 0.01 | 0.03 |
| 2010 | 57.5 | 23.3 | 3.8 | 8.8 | 10.8 | 6.2 | 0.66 | 0.09 | 1.34 | 0.55 | 0.20 |
| 2015 | 83.4 | 28.0 | 10.0 | 18.1 | 17.6 | 6.6 | 1.10 | 0.14 | 2.23 | 0.86 | 0.25 |
| 2019 | 107.6 | 32.4 | 20.0 | 29.4 | 24.0 | 7.1 | 1.59 | 0.19 | 3.06 | 1.09 | 0.29 |
| <i>Unc.</i> | <i>0.5</i> | <i>0.1</i> | <i>1.4</i> | <i>0.6</i> | <i>0.4</i> | <i>0.4</i> | <i>0.06</i> | <i>n.a.</i> | <i>0.06</i> | <i>0.14</i> | <i>n.a.</i> |
| <i>ERF</i> | 18.0 | 6.2 | 2.2 | 6.9 | 4.0 | 0.7 | 0.4 | 0.0 | 0.7 | 0.2 | 0.1 |

Notes: Data merged from AR5 (1750;1850); CMIP6 compilation by Meinshausen et al (2017)until ca. 1995, and data directly taken from merged AGAGE and NOAA networks, depending on date of availability for various components. Uncertainties pertain to 2019, derived from observations made by global networks, and do not include estimates of analytical accuracy. n.a.: not available. ERF(2019-1750) from Chapter 7.

Table AIII.1d: Historical abundances (ppt) and ERF [mWm⁻²] of HCFCs

| Year | HCFC-22 | HCFC-141b | HCFC-142b | HCFC-133a | HCFC-31 | HCFC-124 |
|-------------|---------|-----------|-----------|-----------|---------|----------|
| 1750 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1850 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1900 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1910 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1920 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1930 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1940 | 0.3 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1950 | 0.9 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1960 | 2.3 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1970 | 13.1 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 |
| 1980 | 44.6 | 0.0 | 0.4 | 0.01 | 0.00 | 0.00 |
| 1990 | 89.6 | 0.3 | 1.5 | 0.05 | 0.00 | 0.00 |
| 2000 | 141.8 | 12.7 | 11.4 | 0.11 | 0.027 | 0.00 |
| 2010 | 206.3 | 20.5 | 20.4 | 0.31 | 0.084 | 1.10 |
| 2015 | 233.3 | 24.2 | 22.2 | 0.40 | 0.084 | 1.02 |
| 2019 | 246.8 | 24.4 | 22.3 | n.a. | n.a. | n.a. |
| <i>Unc.</i> | 0.6 | 0.3 | 0.4 | n.a. | n.a. | n.a. |
| <i>ERF</i> | 52.8 | 3.9 | 4.3 | 0.1 | 0.0 | 0.2 |

Notes: 1750/1850 from AR5.; 1900-1970 from CMIP6 dataset in Meinshausen et al. (2017). 1980-1995 AGAGE, or data directly taken from merged AGAGE and NOAA networks, depending on date of availability for various components; HCFC-31 from Schoenenberger et al (2015), HCFC-124 from Simmonds et al. (2017) Uncertainties pertain to 2019, derived from observations made by global networks, and do not include estimates of analytical accuracy. For HCFC-133a, HCFC-31, and HCFC-124 abundances in 2019 and uncertainties are not available. ERF(2019-1750) from Chapter 7, except HCFC-133a, HCFC-31, and HCFC-124 which are for 2015.

Table AIII.1e: Historical abundances (ppt) and ERF [mWm⁻²] of CFCs

| Year | CFC-12 | CFC-11 | CFC-113 | CFC-114 | CFC-115 | CFC-13 | CFC-112 | CFC-112a | CFC-113a | CFC-114a |
|------|--------|--------|---------|---------|---------|--------|---------|----------|----------|----------|
| 1750 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1850 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1900 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1910 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1920 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1930 | 0.0 | 0.0 | 0.0 | 0.0 | 0.00 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1940 | 0.0 | 0.0 | 0.5 | 0.0 | 0.00 | 0.03 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1950 | 6.4 | 0.9 | 1.0 | 1.5 | 0.00 | 0.04 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1960 | 31.6 | 10.2 | 2.0 | 3.9 | 0.00 | 0.05 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1970 | 121.7 | 57.0 | 5.9 | 6.7 | 0.20 | 0.44 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1980 | 304.0 | 166.8 | 20.8 | 10.1 | 1.75 | 1.20 | 0.11 | 0.00 | 0.06 | 0.44 |
| 1990 | 483.1 | 258.1 | 70.6 | 15.6 | 5.46 | 2.42 | 0.31 | 0.00 | 0.16 | 0.91 |
| 2000 | 542.3 | 259.2 | 82.1 | 16.4 | 8.16 | 2.83 | 0.49 | 0.07 | 0.28 | 1.03 |
| 2010 | 530.9 | 239.4 | 75.2 | 16.3 | 8.38 | 3.04 | 0.45 | 0.07 | 0.41 | 1.06 |
| 2015 | 516.6 | 231.0 | 72.0 | 16.0 | 8.46 | 3.16 | 0.42 | 0.07 | 0.62 | 1.05 |
| 2019 | 503.1 | 226.2 | 69.8 | 16.0 | 8.67 | 3.28 | n.a | n.a | n.a | n.a |
| Unc. | 3.2 | 1.1 | 0.3 | 0.1 | 0.02 | 0.02 | n.a | n.a | n.a | n.a |
| ERF | 180.3 | 66.3 | 21.0 | 5.0 | 2.1 | 0.9 | 0.1 | 0.0 | 0.2 | 0.3 |

Notes: 1750/1850 from AR5.; 1900-1970 from CMIP6 dataset in Meinshausen et al. (2017). 1980-1995 AGAGE, or data directly taken from merged AGAGE and NOAA networks, depending on date of availability for various components; CFC-13 from Vollmer et al. (2018) until 2015 then AGAGE; CFC-114 and CFC-114a from AGAGE and Laube et al. (2016) . CFC-112 from Laube et al.(2014); CFC-112a from Adcock et al. (2018). Uncertainties pertain to 2019, derived from observations made by global networks, and do not include estimates of analytical accuracy. For CFC-112, CFC-112a, CFC-113a, and CFC-114a abundances and uncertainties for 2019 are not available. ERF(2019-1750) from Chapter 7, except for CFC-112, CFC-112a, CFC-113a, and CFC-114a for 2015

Table AIII.1f: Historical abundances (ppt) and ERF [mWm⁻²] of CH₃CCl₃, CCl₄, CH₃Br, CHCl₃ and Halons.

| Year | CH ₃ CCl ₃ | CCl ₄ | CH ₃ Cl | CH ₃ Br | CH ₂ Cl ₂ | CHCl ₃ | Halon-1211 | Halon-1301 | Halon-2402 |
|-------------|----------------------------------|------------------|--------------------|--------------------|---------------------------------|-------------------|-------------|-------------|-------------|
| 1750 | 0.00 | 0.03 | 457 | 5.30 | 7 | 4.8 | 0.00 | 0.00 | 0.00 |
| 1850 | 0.00 | 0.03 | 457 | 5.30 | 7 | 4.8 | 0.00 | 0.00 | 0.00 |
| 1900 | 0.00 | 0.03 | 457 | 5.30 | 7 | 4.8 | 0.00 | 0.00 | 0.00 |
| 1910 | 0.00 | 0.03 | 457 | 5.30 | 7 | 4.8 | 0.00 | 0.00 | 0.00 |
| 1920 | 0.00 | 1.2 | 457 | 5.30 | 7 | 4.8 | 0.00 | 0.00 | 0.00 |
| 1930 | 0.00 | 4.1 | 457 | 5.30 | 7 | 5.0 | 0.00 | 0.00 | 0.00 |
| 1940 | 0.00 | 14.1 | 457 | 5.66 | 7 | 5.3 | 0.00 | 0.00 | 0.00 |
| 1950 | 0.00 | 35.5 | 478 | 6.06 | 8 | 5.7 | 0.03 | 0.00 | 0.00 |
| 1960 | 1.70 | 53.2 | 512 | 6.50 | 11 | 6.4 | 0.02 | 0.00 | 0.00 |
| 1970 | 17.7 | 77.0 | 540 | 7.06 | 14 | 7.5 | 0.04 | 0.00 | 0.02 |
| 1980 | 85.9 | 93.8 | 549 | 7.77 | 18 | 8.8 | 0.71 | 0.38 | 0.15 |
| 1990 | 129.3 | 106.2 | 550 | 8.69 | 20 | 10.3 | 2.44 | 1.85 | 0.37 |
| 2000 | 45.4 | 98.1 | 547 | 9.09 | 20 | 7.5 | 4.12 | 2.82 | 0.48 |
| 2010 | 7.6 | 87.3 | 538 | 7.14 | 29 | 7.3 | 4.12 | 3.21 | 0.46 |
| 2015 | 3.1 | 81.6 | 547 | 6.68 | 35 | 8.6 | 3.66 | 3.31 | 0.42 |
| 2019 | 1.6 | 77.9 | 551 | 6.49 | 41 | 8.8 | 3.28 | 3.32 | 0.40 |
| <i>Unc.</i> | <i>0.1</i> | <i>0.7</i> | <i>5</i> | <i>0.07</i> | <i>6</i> | <i>0.3</i> | <i>0.05</i> | <i>0.07</i> | <i>0.03</i> |
| <i>ERF</i> | <i>0.1</i> | <i>12.9</i> | <i>0.4</i> | <i>0.0</i> | <i>1.0</i> | <i>0.3</i> | <i>1.0</i> | <i>1.0</i> | <i>0.1</i> |

Notes: 1750 from AR5.; 1850-1970 from CMIP6 dataset in Meinshausen (2017). 1980-2019 AGAGE or merged AGAGE and NOAA networks, depending on date of availability. ERF(2019-1750) from Chapter 7.

Table AIII.2: Future abundances of CO₂, CH₄, and N₂O for selected SSP scenarios [2020-2500]

| Year/ Scena rio | CO ₂ (ppm) | | | | | CH ₄ (ppb) | | | | | N ₂ O (ppb) | | | | |
|-----------------------|-----------------------|---------|---------|---------|---------|-----------------------|---------|---------|---------|---------|------------------------|---------|---------|---------|-----|
| | 2019* | 410 | | | | 1866 | | | | | 332 | | | | |
| SSP1-19 | SSP1-126 | SSP2-45 | SSP3-70 | SSP5-58 | SSP1-19 | SSP1-126 | SSP2-45 | SSP3-70 | SSP5-58 | SSP1-19 | SSP1-126 | SSP2-45 | SSP3-70 | SSP5-58 | |
| 2020 | 414 | 414 | 414 | 415 | 415 | 1894 | 1888 | 1911 | 1921 | 1907 | 332 | 332 | 332 | 332 | 332 |
| 2030 | 434 | 440 | 444 | 451 | 452 | 1796 | 1810 | 2002 | 2099 | 2018 | 337 | 337 | 340 | 341 | 341 |
| 2040 | 440 | 458 | 475 | 493 | 500 | 1593 | 1663 | 2045 | 2289 | 2209 | 341 | 341 | 348 | 351 | 350 |
| 2050 | 438 | 469 | 507 | 541 | 563 | 1428 | 1519 | 2020 | 2472 | 2446 | 344 | 344 | 356 | 362 | 358 |
| 2060 | 431 | 474 | 537 | 593 | 643 | 1305 | 1402 | 1942 | 2655 | 2613 | 346 | 346 | 363 | 373 | 366 |
| 2070 | 424 | 473 | 564 | 652 | 744 | 1220 | 1299 | 1854 | 2840 | 2670 | 348 | 348 | 369 | 385 | 374 |
| 2080 | 415 | 467 | 585 | 716 | 864 | 1150 | 1197 | 1779 | 3028 | 2652 | 350 | 349 | 373 | 397 | 380 |
| 2090 | 405 | 457 | 598 | 787 | 998 | 1088 | 1112 | 1719 | 3208 | 2549 | 352 | 352 | 376 | 409 | 387 |
| 2100 | 394 | 446 | 603 | 867 | 1135 | 1036 | 1056 | 1683 | 3372 | 2415 | 354 | 354 | 377 | 422 | 392 |
| 2200 | 343 | 403 | 643 | 1457 | 2108 | 929 | 928 | 1255 | 2572 | 1516 | 364 | 363 | 376 | 497 | 414 |
| 2300 | 342 | 396 | 621 | 1483 | 2162 | 872 | 864 | 1001 | 1988 | 1068 | 361 | 360 | 367 | 511 | 411 |
| 2400 | 339 | 389 | 598 | 1424 | 2080 | 871 | 864 | 999 | 1959 | 1038 | 358 | 358 | 362 | 514 | 408 |
| 2500 | 337 | 384 | 579 | 1371 | 2010 | 871 | 864 | 997 | 1938 | 1019 | 357 | 356 | 360 | 516 | 407 |

Note: *observed from Table AIII.1a. SSP GHG concentrations (Meinshausen et al., 2017, 2020) available at greenhousegases.science.unimelb.edu.au. Concentrations of halogens in electronic supplement. Major scenarios used in this report are selected.

Table AIII.3: Effective Radiative Forcing (W m^{-2}) time series of all climate forcers from 1750-2019.

| year | CO_2 | CH_4 | N_2O | Halogens | O_3 | $\text{H}_2\text{O STRAT}$ | Contrail-cirrus | Aerosol radiation interactions | Aerosol cloud interactions | BC on snow | Land use | Volcanic | Solar | Total anthropogenic | Total natural | Total |
|------|---------------|---------------|----------------------|----------|--------------|----------------------------|-----------------|--------------------------------|----------------------------|------------|----------|----------|-------|---------------------|---------------|-------|
| 1750 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.20 | 0.10 | 0.00 | 0.30 | 0.30 |
| 1850 | 0.14 | 0.05 | 0.01 | 0.00 | 0.03 | 0.00 | 0.00 | -0.01 | -0.07 | 0.01 | -0.03 | 0.19 | 0.01 | 0.13 | 0.20 | 0.33 |
| 1900 | 0.35 | 0.12 | 0.03 | 0.00 | 0.08 | 0.01 | 0.00 | -0.06 | -0.29 | 0.02 | -0.08 | 0.20 | -0.04 | 0.18 | 0.16 | 0.34 |
| 1910 | 0.41 | 0.15 | 0.04 | 0.00 | 0.09 | 0.01 | 0.00 | -0.10 | -0.42 | 0.03 | -0.10 | 0.20 | -0.02 | 0.12 | 0.18 | 0.30 |
| 1920 | 0.50 | 0.17 | 0.05 | 0.00 | 0.10 | 0.02 | 0.00 | -0.10 | -0.43 | 0.03 | -0.11 | 0.19 | 0.01 | 0.23 | 0.20 | 0.43 |
| 1930 | 0.54 | 0.20 | 0.06 | 0.00 | 0.12 | 0.02 | 0.00 | -0.11 | -0.46 | 0.03 | -0.13 | 0.19 | 0.02 | 0.26 | 0.21 | 0.48 |
| 1940 | 0.62 | 0.22 | 0.06 | 0.00 | 0.14 | 0.02 | 0.00 | -0.15 | -0.52 | 0.03 | -0.14 | 0.19 | 0.04 | 0.29 | 0.23 | 0.52 |
| 1950 | 0.65 | 0.24 | 0.07 | 0.01 | 0.17 | 0.02 | 0.00 | -0.15 | -0.55 | 0.03 | -0.14 | 0.18 | 0.06 | 0.35 | 0.24 | 0.59 |
| 1960 | 0.71 | 0.29 | 0.08 | 0.03 | 0.22 | 0.03 | 0.01 | -0.25 | -0.73 | 0.04 | -0.17 | 0.18 | 0.09 | 0.26 | 0.27 | 0.54 |
| 1970 | 0.85 | 0.36 | 0.09 | 0.08 | 0.28 | 0.03 | 0.02 | -0.38 | -0.92 | 0.05 | -0.18 | 0.05 | 0.08 | 0.29 | 0.13 | 0.42 |
| 1980 | 1.09 | 0.43 | 0.11 | 0.20 | 0.33 | 0.04 | 0.02 | -0.41 | -1.04 | 0.06 | -0.18 | 0.09 | 0.11 | 0.66 | 0.19 | 0.86 |
| 1990 | 1.33 | 0.49 | 0.13 | 0.33 | 0.36 | 0.04 | 0.03 | -0.38 | -1.05 | 0.07 | -0.19 | 0.14 | 0.11 | 1.17 | 0.24 | 1.42 |
| 2000 | 1.56 | 0.51 | 0.16 | 0.37 | 0.40 | 0.05 | 0.04 | -0.30 | -0.92 | 0.07 | -0.19 | 0.18 | 0.11 | 1.74 | 0.29 | 2.02 |
| 2010 | 1.85 | 0.52 | 0.18 | 0.39 | 0.44 | 0.05 | 0.04 | -0.27 | -0.99 | 0.08 | -0.20 | 0.14 | -0.01 | 2.10 | 0.13 | 2.23 |
| 2015 | 2.01 | 0.53 | 0.20 | 0.40 | 0.47 | 0.05 | 0.05 | -0.23 | -0.89 | 0.08 | -0.20 | 0.11 | 0.03 | 2.47 | 0.14 | 2.61 |
| 2019 | 2.16 | 0.54 | 0.21 | 0.41 | 0.47 | 0.05 | 0.06 | -0.22 | -0.84 | 0.08 | -0.20 | 0.14 | -0.02 | 2.72 | 0.12 | 2.84 |

Note: O_3 includes tropospheric and stratospheric O_3 , dominated by tropospheric O_3 . Stratospheric water vapour from methane oxidation is a linear function of the methane ERF (Section 7.3.2.6). Contrail forcing is a linear scaling of aviation NOx emissions, scaled to ERF in 2018 (Lee et al., 2021). Present-day aerosol forcing is assessed in Section 7.3.3 as -0.3 (-0.6 to 0.0) W m^{-2} for aerosol-radiation interactions and -1.0 (-1.7 to -0.3) W m^{-2} for aerosol-cloud interactions for the 2005-2014 mean relative to 1750. Land use change considers albedo and irrigation effects (Section 7.3.4.1). BC on snow forcing is linear with emissions of BC (Section 7.3.4.3). Volcanic forcing is positive in years without large volcanic eruptions such that the long-term pre-industrial (500 BCE to 1749 CE) mean volcanic forcing is zero. Solar forcing is derived from the ^{14}C reconstruction of total solar irradiance in the combined PMIP4/CMIP6 dataset (Jungclaus et al., 2017; Matthes et al., 2017). Present-day solar forcing is assessed in Section 7.3.4.4 as +0.01 (-0.06 to +0.08) W m^{-2} , based on the mean total solar irradiance from solar cycle 24 (2009-2019) compared a long pre-industrial baseline period (6754 BCE - 1744 CE); the 2019 ERF value differs from this as it represents a single year near the solar minimum. Natural is the sum of volcanic and solar forcing, while anthropogenic includes all others. Further details on methods for computing ERF are in Chapter 7 Supplementary Material 7.SM.1.3.

Table AIII.4a: Effective Radiative Forcing (Wm⁻²) time series of all climate forcers for SSP1-19

| year | CO ₂ | CH ₄ | N ₂ O | Halogens | O ₃ | H ₂ O STRAT | Contrail-cirrus | Aerosol radiation interactions | Aerosol cloud interactions | BC on snow | Land use | Volcanic | Solar | Total anthropogenic | Total natural | Total |
|------|-----------------|-----------------|------------------|----------|----------------|------------------------|-----------------|--------------------------------|----------------------------|------------|----------|----------|-------|---------------------|---------------|-------|
| 2020 | 2.22 | 0.55 | 0.21 | 0.40 | 0.42 | 0.05 | 0.05 | -0.20 | -0.81 | 0.08 | 0.20 | 0.06 | -0.02 | 2.77 | 0.04 | 2.81 |
| 2030 | 2.49 | 0.52 | 0.23 | 0.38 | 0.25 | 0.05 | 0.03 | -0.16 | -0.39 | 0.03 | 0.21 | 0.00 | -0.02 | 3.22 | -0.02 | 3.20 |
| 2040 | 2.56 | 0.44 | 0.24 | 0.32 | 0.19 | 0.04 | 0.02 | -0.16 | -0.28 | 0.02 | 0.21 | 0.00 | -0.01 | 3.18 | -0.01 | 3.18 |
| 2050 | 2.53 | 0.37 | 0.25 | 0.27 | 0.14 | 0.03 | 0.02 | -0.17 | -0.20 | 0.01 | 0.21 | 0.00 | 0.01 | 3.04 | 0.01 | 3.05 |
| 2060 | 2.45 | 0.31 | 0.26 | 0.23 | 0.12 | 0.03 | 0.01 | -0.17 | -0.16 | 0.01 | 0.21 | 0.00 | 0.01 | 2.87 | 0.01 | 2.88 |
| 2070 | 2.35 | 0.27 | 0.26 | 0.21 | 0.11 | 0.02 | 0.01 | -0.16 | -0.13 | 0.00 | 0.20 | 0.00 | 0.02 | 2.74 | 0.02 | 2.76 |
| 2080 | 2.23 | 0.24 | 0.27 | 0.19 | 0.09 | 0.02 | 0.01 | -0.15 | -0.09 | 0.00 | 0.19 | 0.00 | 0.02 | 2.62 | 0.02 | 2.64 |
| 2090 | 2.09 | 0.21 | 0.28 | 0.17 | 0.09 | 0.02 | 0.01 | -0.14 | -0.05 | 0.00 | 0.19 | 0.00 | 0.01 | 2.48 | 0.01 | 2.48 |
| 2100 | 1.92 | 0.18 | 0.28 | 0.15 | 0.08 | 0.02 | 0.01 | -0.13 | -0.01 | 0.00 | 0.18 | 0.00 | 0.00 | 2.33 | 0.00 | 2.33 |
| 2200 | 1.16 | 0.12 | 0.32 | 0.07 | 0.06 | 0.01 | 0.00 | -0.10 | 0.08 | -0.01 | 0.17 | 0.00 | 0.03 | 1.55 | 0.03 | 1.58 |
| 2300 | 1.14 | 0.09 | 0.31 | 0.04 | 0.03 | 0.01 | 0.00 | -0.09 | 0.14 | -0.01 | 0.17 | 0.00 | 0.00 | 1.49 | 0.00 | 1.49 |
| 2400 | 1.09 | 0.09 | 0.30 | 0.03 | 0.03 | 0.01 | 0.00 | -0.09 | 0.14 | -0.01 | 0.17 | 0.00 | 0.00 | 1.42 | 0.00 | 1.42 |
| 2500 | 1.05 | 0.09 | 0.30 | 0.02 | 0.03 | 0.01 | 0.00 | -0.09 | 0.14 | -0.01 | 0.17 | 0.00 | 0.00 | 1.38 | 0.00 | 1.38 |

Note: ERF based on future abundancies of WMGHGs listed in AIII.2 . See notes of AIII.3. Future ozone forcing uses projected emissions of carbon monoxide, volatile organic carbon, nitrogen oxides, and concentrations of methane, nitrous oxide and halogenated compounds with relationships to forcing derived from Thornhill et al. (2021b, 2021a). Future contrail forcing is a linear scaling of future NOx emissions (Smith et al., 2018) and scaled to year-2018 ERF (Lee et al., 2021). Future aerosol forcing is based on emissions of black carbon, organic carbon, sulphur dioxide and ammonia using method described in Smith et al. (2018). Land use forcing scales with cumulative emissions of future land-use CO₂ (Smith et al., 2018). Future volcanic forcing set to zero from a 10-year linear transition from the end of the historical period following Eyring et al. (2016). Solar forcing is set to zero from 2300 CE. Further details on methods for computing SSP-projection ERF are in Chapter 7 Supplementary Material 7.SM.1.4

Table AIII.4b: Effective Radiative Forcing (Wm⁻²) time series of all climate forcers for SSP1-26

| year | CO ₂ | CH ₄ | N ₂ O | Halogens | O ₃ | H ₂ O STRAT | Contrail-cirrus | Aerosol radiation interactions | Aerosol cloud interactions | BC on snow | Land use | Volcanic | Solar | Total anthropogenic | Total natural | Total | |
|------|-----------------|-----------------|------------------|----------|----------------|------------------------|-----------------|--------------------------------|----------------------------|------------|----------|----------|-------|---------------------|---------------|-------|------|
| 2020 | 2.22 | 0.55 | 0.21 | 0.40 | 0.42 | 0.05 | 0.05 | -0.20 | -0.81 | 0.08 | 0.20 | - | 0.06 | -0.02 | 2.76 | 0.04 | 2.80 |
| 2030 | 2.56 | 0.52 | 0.22 | 0.39 | 0.30 | 0.05 | 0.04 | -0.19 | -0.51 | 0.04 | 0.21 | 0.00 | -0.02 | 3.23 | -0.02 | 3.21 | |
| 2040 | 2.79 | 0.46 | 0.24 | 0.34 | 0.26 | 0.04 | 0.04 | -0.16 | -0.36 | 0.03 | 0.20 | 0.00 | -0.01 | 3.49 | -0.01 | 3.48 | |
| 2050 | 2.93 | 0.41 | 0.25 | 0.28 | 0.21 | 0.04 | 0.04 | -0.15 | -0.26 | 0.02 | 0.20 | 0.00 | 0.01 | 3.56 | 0.01 | 3.58 | |
| 2060 | 2.99 | 0.35 | 0.25 | 0.24 | 0.18 | 0.03 | 0.04 | -0.13 | -0.20 | 0.02 | 0.20 | 0.00 | 0.01 | 3.58 | 0.01 | 3.58 | |
| 2070 | 2.98 | 0.31 | 0.26 | 0.21 | 0.15 | 0.03 | 0.04 | -0.13 | -0.16 | 0.02 | 0.19 | 0.00 | 0.02 | 3.52 | 0.02 | 3.54 | |
| 2080 | 2.91 | 0.26 | 0.27 | 0.19 | 0.12 | 0.02 | 0.04 | -0.13 | -0.10 | 0.01 | 0.18 | 0.00 | 0.02 | 3.40 | 0.02 | 3.42 | |
| 2090 | 2.78 | 0.22 | 0.28 | 0.17 | 0.10 | 0.02 | 0.03 | -0.13 | -0.06 | 0.01 | 0.17 | 0.00 | 0.01 | 3.24 | 0.01 | 3.25 | |
| 2100 | 2.63 | 0.19 | 0.28 | 0.16 | 0.08 | 0.02 | 0.03 | -0.12 | -0.01 | 0.00 | 0.17 | 0.00 | 0.00 | 3.10 | 0.00 | 3.10 | |
| 2200 | 2.06 | 0.12 | 0.32 | 0.07 | 0.04 | 0.01 | 0.01 | -0.10 | 0.08 | -0.01 | 0.15 | 0.00 | 0.03 | 2.47 | 0.03 | 2.50 | |
| 2300 | 1.96 | 0.08 | 0.31 | 0.04 | 0.01 | 0.01 | 0.00 | -0.09 | 0.14 | -0.01 | 0.15 | 0.00 | 0.00 | 2.30 | 0.00 | 2.30 | |
| 2400 | 1.87 | 0.08 | 0.30 | 0.03 | 0.01 | 0.01 | 0.00 | -0.09 | 0.14 | -0.01 | 0.15 | 0.00 | 0.00 | 2.19 | 0.00 | 2.19 | |
| 2500 | 1.79 | 0.08 | 0.29 | 0.02 | 0.01 | 0.01 | 0.00 | -0.09 | 0.14 | -0.01 | 0.15 | 0.00 | 0.00 | 2.11 | 0.00 | 2.11 | |

3
4
5 Notes: see Table AIII.3 and Table AIII.4a.
6

Table AIII.4c: Effective Radiative Forcing (Wm⁻²) time series of all climate forcers for SSP2-45

| year | CO ₂ | CH ₄ | N ₂ O | Halogens | O ₃ | H ₂ O STRAT | Contrail-cirrus | Aerosol radiation interactions | Aerosol cloud interactions | BC on snow | Land use | Volcanic | Solar | Total anthropogenic | Total natural | Total | |
|------|-----------------|-----------------|------------------|----------|----------------|------------------------|-----------------|--------------------------------|----------------------------|------------|----------|----------|-------|---------------------|---------------|-------|------|
| 2020 | 2.22 | 0.56 | 0.21 | 0.40 | 0.45 | 0.05 | 0.05 | -0.18 | -0.88 | 0.10 | 0.20 | - | 0.06 | -0.02 | 2.79 | 0.04 | 2.83 |
| 2030 | 2.62 | 0.59 | 0.23 | 0.40 | 0.47 | 0.05 | 0.07 | -0.21 | -0.84 | 0.09 | 0.21 | 0.00 | -0.02 | 3.25 | -0.02 | 3.24 | |
| 2040 | 3.01 | 0.61 | 0.26 | 0.37 | 0.45 | 0.06 | 0.09 | -0.20 | -0.74 | 0.07 | 0.22 | 0.00 | -0.01 | 3.76 | -0.01 | 3.75 | |
| 2050 | 3.38 | 0.60 | 0.28 | 0.34 | 0.43 | 0.06 | 0.09 | -0.18 | -0.61 | 0.06 | 0.22 | 0.00 | 0.01 | 4.23 | 0.01 | 4.24 | |
| 2060 | 3.72 | 0.57 | 0.30 | 0.32 | 0.39 | 0.05 | 0.10 | -0.18 | -0.51 | 0.05 | 0.22 | 0.00 | 0.01 | 4.59 | 0.01 | 4.60 | |
| 2070 | 4.01 | 0.54 | 0.32 | 0.30 | 0.35 | 0.05 | 0.12 | -0.19 | -0.43 | 0.04 | 0.21 | 0.00 | 0.02 | 4.89 | 0.02 | 4.91 | |
| 2080 | 4.23 | 0.51 | 0.33 | 0.29 | 0.31 | 0.05 | 0.15 | -0.19 | -0.34 | 0.02 | 0.20 | 0.00 | 0.02 | 5.15 | 0.02 | 5.16 | |
| 2090 | 4.35 | 0.49 | 0.34 | 0.28 | 0.28 | 0.04 | 0.16 | -0.19 | -0.25 | 0.01 | 0.19 | 0.00 | 0.01 | 5.32 | 0.01 | 5.33 | |
| 2100 | 4.40 | 0.47 | 0.35 | 0.27 | 0.25 | 0.04 | 0.18 | -0.19 | -0.21 | 0.01 | 0.18 | 0.00 | 0.00 | 5.40 | 0.00 | 5.40 | |
| 2200 | 4.79 | 0.29 | 0.35 | 0.14 | 0.08 | 0.03 | 0.06 | -0.13 | -0.01 | -0.01 | 0.15 | 0.00 | 0.03 | 5.43 | 0.03 | 5.46 | |
| 2300 | 4.59 | 0.16 | 0.32 | 0.05 | 0.02 | 0.01 | 0.00 | -0.11 | 0.09 | -0.01 | 0.15 | 0.00 | 0.00 | 4.94 | 0.00 | 4.94 | |
| 2400 | 4.35 | 0.16 | 0.31 | 0.03 | 0.02 | 0.01 | 0.00 | -0.11 | 0.09 | -0.01 | 0.15 | 0.00 | 0.00 | 4.67 | 0.00 | 4.67 | |
| 2500 | 4.17 | 0.16 | 0.30 | 0.03 | - | 0.01 | 0.00 | -0.11 | 0.09 | -0.01 | 0.15 | 0.00 | 0.00 | 4.47 | 0.00 | 4.47 | |

3
4 Notes: see Table AIII.3 and Table AIII.4a.
5
6

1 **Table AIII.4d:** Effective Radiative Forcing (Wm^{-2}) time series of all climate forcers for SSP3-70
2

| year | CO_2 | CH_4 | N_2O | Halogens | O_3 | $\text{H}_2\text{O STRAT}$ | Contrail-cirrus | Aerosol radiation interactions | Aerosol cloud interactions | BC on snow | Land use | Volcanic | Solar | Total anthropogenic | Total natural | Total |
|------|---------------|---------------|----------------------|----------|--------------|----------------------------|-----------------|--------------------------------|----------------------------|------------|----------|----------|-------|---------------------|---------------|-------|
| 2020 | 2.23 | 0.56 | 0.21 | 0.40 | 0.51 | 0.05 | 0.05 | -0.19 | -1.02 | 0.12 | 0.20 | 0.06 | -0.02 | 2.72 | 0.04 | 2.76 |
| 2030 | 2.71 | 0.63 | 0.23 | 0.39 | 0.57 | 0.06 | 0.07 | -0.19 | -1.04 | 0.12 | 0.21 | 0.00 | -0.02 | 3.33 | -0.02 | 3.31 |
| 2040 | 3.22 | 0.69 | 0.26 | 0.37 | 0.62 | 0.06 | 0.08 | -0.19 | -1.06 | 0.13 | 0.22 | 0.00 | -0.01 | 3.97 | -0.01 | 3.96 |
| 2050 | 3.76 | 0.75 | 0.29 | 0.36 | 0.66 | 0.07 | 0.09 | -0.20 | -1.06 | 0.13 | 0.23 | 0.00 | 0.01 | 4.61 | 0.01 | 4.63 |
| 2060 | 4.31 | 0.80 | 0.33 | 0.35 | 0.69 | 0.07 | 0.09 | -0.20 | -1.03 | 0.12 | 0.24 | 0.00 | 0.01 | 5.30 | 0.01 | 5.30 |
| 2070 | 4.87 | 0.86 | 0.36 | 0.35 | 0.71 | 0.08 | 0.10 | -0.20 | -0.99 | 0.12 | 0.25 | 0.00 | 0.02 | 6.00 | 0.02 | 6.02 |
| 2080 | 5.45 | 0.91 | 0.39 | 0.35 | 0.73 | 0.08 | 0.10 | -0.20 | -0.95 | 0.11 | 0.26 | 0.00 | 0.02 | 6.71 | 0.02 | 6.72 |
| 2090 | 6.04 | 0.96 | 0.42 | 0.35 | 0.75 | 0.09 | 0.10 | -0.21 | -0.92 | 0.11 | 0.26 | 0.00 | 0.01 | 7.43 | 0.01 | 7.44 |
| 2100 | 6.64 | 1.00 | 0.45 | 0.36 | 0.77 | 0.09 | 0.11 | -0.20 | -0.87 | 0.10 | 0.27 | 0.00 | 0.00 | 8.18 | 0.00 | 8.18 |
| 2200 | 10.00 | 0.77 | 0.64 | 0.22 | 0.38 | 0.07 | 0.04 | -0.16 | -0.32 | 0.03 | 0.29 | 0.00 | 0.03 | 11.37 | 0.03 | 11.40 |
| 2300 | 10.11 | 0.58 | 0.68 | 0.08 | 0.15 | 0.05 | 0.00 | -0.14 | 0.02 | -0.01 | 0.29 | 0.00 | 0.00 | 11.24 | 0.00 | 11.24 |
| 2400 | 9.84 | 0.57 | 0.69 | 0.06 | 0.14 | 0.05 | 0.00 | -0.14 | 0.02 | -0.01 | 0.29 | 0.00 | 0.00 | 10.94 | 0.00 | 10.94 |
| 2500 | 9.60 | 0.56 | 0.70 | 0.06 | 0.13 | 0.05 | 0.00 | -0.14 | 0.02 | -0.01 | 0.29 | 0.00 | 0.00 | 10.68 | 0.00 | 10.68 |

3 Notes: see Table AIII.3 and Table AIII.4a.
4
5
6

1 **Table AIII.4e:** Effective Radiative Forcing (Wm^{-2}) time series of all climate forcers for SSP5-85
2

| year | CO_2 | CH_4 | N_2O | Halogens | O_3 | $\text{H}_2\text{O STRAT}$ | Contrail-cirrus | Aerosol radiation interactions | Aerosol cloud interactions | BC on snow | Land use | Volcanic | Solar | Total anthropogenic | Total natural | Total | |
|------|---------------|---------------|----------------------|----------|--------------|----------------------------|-----------------|--------------------------------|----------------------------|------------|----------|----------|-------|---------------------|---------------|-------|-------|
| 2020 | 2.23 | 0.56 | 0.21 | 0.40 | 0.48 | 0.05 | 0.06 | -0.14 | -0.86 | 0.11 | 0.20 | 0.06 | -0.02 | 2.90 | 0.04 | 2.94 | |
| 2030 | 2.71 | 0.60 | 0.23 | 0.41 | 0.49 | 0.06 | 0.08 | -0.13 | -0.71 | 0.09 | 0.21 | 0.00 | -0.02 | 3.61 | 0.02 | 3.59 | |
| 2040 | 3.30 | 0.66 | 0.26 | 0.41 | 0.54 | 0.06 | 0.09 | -0.14 | -0.67 | 0.09 | 0.23 | 0.00 | -0.01 | 4.37 | 0.01 | 4.37 | |
| 2050 | 4.00 | 0.74 | 0.28 | 0.42 | 0.56 | 0.07 | 0.10 | -0.15 | -0.57 | 0.07 | 0.23 | 0.00 | 0.01 | 5.29 | 0.01 | 5.30 | |
| 2060 | 4.79 | 0.79 | 0.30 | 0.45 | 0.60 | 0.07 | 0.12 | -0.18 | -0.58 | 0.06 | 0.24 | 0.00 | 0.01 | 6.21 | 0.01 | 6.22 | |
| 2070 | 5.68 | 0.81 | 0.32 | 0.50 | 0.60 | 0.07 | 0.14 | -0.20 | -0.55 | 0.06 | 0.24 | 0.00 | 0.02 | 7.19 | 0.02 | 7.21 | |
| 2080 | 6.62 | 0.80 | 0.34 | 0.55 | 0.56 | 0.07 | 0.15 | -0.20 | -0.48 | 0.05 | 0.23 | 0.00 | 0.02 | 8.24 | 0.02 | 8.25 | |
| 2090 | 7.54 | 0.77 | 0.36 | 0.58 | 0.49 | 0.07 | 0.15 | -0.21 | -0.38 | 0.03 | 0.23 | 0.00 | 0.01 | 9.18 | 0.01 | 9.19 | |
| 2100 | 8.38 | 0.73 | 0.37 | 0.60 | 0.40 | 0.07 | 0.15 | -0.21 | -0.27 | 0.02 | 0.23 | 0.00 | 0.00 | 10.01 | 0.00 | 10.00 | |
| 2200 | 12.30 | 0.40 | 0.42 | 0.32 | 0.06 | 0.04 | 0.05 | -0.16 | -0.07 | 0.00 | 0.22 | 0.00 | 0.03 | 13.14 | 0.03 | 13.17 | |
| 2300 | 12.46 | 0.19 | 0.42 | 0.08 | - | 0.11 | 0.02 | 0.00 | -0.14 | 0.03 | -0.01 | 0.22 | 0.00 | 0.00 | 12.72 | 0.00 | 12.72 |
| 2400 | 12.22 | 0.18 | 0.42 | 0.06 | 0.13 | 0.02 | 0.00 | -0.14 | 0.03 | -0.01 | 0.22 | 0.00 | 0.00 | 12.43 | 0.00 | 12.43 | |
| 2500 | 12.01 | 0.17 | 0.41 | 0.06 | 0.14 | 0.02 | 0.00 | -0.14 | 0.03 | -0.01 | 0.22 | 0.00 | 0.00 | 12.19 | 0.00 | 12.19 | |

3

4 Notes: see Table AIII.3 and Table AIII.4a.

5

6

Table AIII.4f: Effective Radiative Forcing (mW m^{-2}) time series of halogen components for selected scenarios and Kyoto and Montreal gases.

| | SSP1-19 | | | | | | SSP3-70 | | | | | | SSP5-85 | | | | | |
|---------------|---------|--|-------------|---|-------|----------|---------|--|-------------|---|-------|----------|---------|--|-------------|---|-------|----------|
| 2019 (obs) | 40 | 14 | 338 | 17 | 54 | 394 | 40 | 14 | 311 | 17 | 54 | 367 | 40 | 14 | 311 | 17 | 54 | 367 |
| year | HFCs | NF ₃ , SF ₆ , PFCs | CFCS, HCFCs | CH ₃ CCl ₃ , CCL ₄ , CH ₃ Cl, CH ₃ Br | Kyoto | Montreal | HFCs | NF ₃ , SF ₆ , PFCs | CFCS, HCFCs | CH ₃ CCl ₃ , CCL ₄ , CH ₃ Cl, CH ₃ Br | Kyoto | Montreal | HFCs | NF ₃ , SF ₆ , PFCs | CFCS, HCFCs | CH ₃ CCl ₃ , CCL ₄ , CH ₃ Cl, CH ₃ Br | Kyoto | Montreal |
| 2020 | 43 | 14 | 330 | 16 | 57 | 390 | 42 | 14 | 330 | 16 | 56 | 389 | 43 | 14 | 330 | 17 | 57 | 390 |
| 2030 | 53 | 16 | 296 | 12 | 70 | 308 | 68 | 18 | 295 | 13 | 86 | 308 | 83 | 18 | 294 | 14 | 101 | 308 |
| 2040 | 41 | 17 | 249 | 9 | 59 | 258 | 92 | 21 | 248 | 10 | 113 | 258 | 130 | 21 | 246 | 11 | 151 | 257 |
| 2050 | 33 | 18 | 209 | 7 | 51 | 216 | 116 | 24 | 208 | 8 | 140 | 216 | 180 | 25 | 207 | 8 | 205 | 215 |
| 2060 | 28 | 19 | 180 | 5 | 47 | 185 | 138 | 27 | 178 | 6 | 164 | 184 | 238 | 28 | 177 | 7 | 266 | 184 |
| 2070 | 25 | 19 | 157 | 4 | 45 | 161 | 157 | 29 | 155 | 5 | 187 | 160 | 304 | 31 | 154 | 6 | 336 | 160 |
| 2080 | 24 | 20 | 139 | 3 | 44 | 141 | 176 | 32 | 136 | 4 | 208 | 140 | 370 | 35 | 135 | 5 | 405 | 140 |
| 2090 | 23 | 20 | 123 | 2 | 43 | 125 | 195 | 35 | 119 | 3 | 229 | 123 | 420 | 38 | 118 | 4 | 459 | 122 |
| 2100 | 22 | 20 | 110 | 1 | 43 | 111 | 212 | 38 | 105 | 3 | 250 | 108 | 447 | 41 | 104 | 4 | 489 | 107 |
| 2200 | 12 | 22 | 38 | 0 | 34 | 38 | 132 | 56 | 31 | 0 | 187 | 31 | 225 | 61 | 29 | 1 | 286 | 30 |

Notes: ERF for 2019 was calculated using concentrations list in Table AIII.1, 2020 and onward are scenario projections. ERF was calculated using concentrations of individual halogen components. Updated from (Meinshausen et al., 2017, 2020) available at greenhousegases.science.unimelb.edu.au.

Table AIII.5:Total Anthropogenic and natural ERF relative to 1750 assessed in AR5 (RCP scenarios) and AR6 (SSP scenarios).

| | | 2030 | | | 2050 | | | 2090 | | |
|-----|--|----------------|---------|------------------|----------------|---------|------------------|----------------|---------|-------------------|
| | | Anthro-pogenic | Natural | Total | Anthro-pogenic | Natural | Total | Anthro-pogenic | Natural | Total |
| AR5 | RCP2.6 ¹ | 2.52 | | 2.50 ± 0.51 | 2.64 | | 2.65 ± 0.47 | 2.35 | | 2.44 ± 0.49 |
| | RCP4.5 ¹ | 2.67 | | 2.61 ± 0.54 | 3.42 | | 3.25 ± 0.56 | 3.91 | | 3.78 ± 0.58 |
| | RCP6.0 ¹ | 2.52 | | 2.41 ± 0.60 | 3.20 | | 3.07 ± 0.61 | 4.93 | | 4.64 ± 0.71 |
| | RCP8.5 ¹ | 2.91 | | 2.92 ± 0.57 | 4.37 | | 4.21 ± 0.63 | 7.32 | | 7.13 ± 0.89 |
| AR6 | RCP2.6 ² | 2.85 | -0.02 | 2.83 (1.93-3.61) | 3.11 | 0.01 | 3.12 (2.33-3.76) | 2.70 | 0.01 | 2.71 (2.04-3.22) |
| | RCP4.5 ² | 3.01 | -0.02 | 2.99 (2.12-3.84) | 3.90 | 0.01 | 3.91 (3.07-4.68) | 4.48 | 0.01 | 4.49 (3.75-5.15) |
| | RCP6.0 ² | 2.84 | -0.02 | 2.82 (1.88-3.69) | 3.56 | 0.01 | 3.58 (2.59-4.47) | 5.51 | 0.01 | 5.52 (4.50-6.38) |
| | RCP8.5 ² | 3.36 | -0.02 | 3.34 (2.40-4.23) | 5.01 | 0.01 | 5.03 (4.06-5.92) | 8.46 | 0.01 | 8.46 (7.23-9.61) |
| | SSP1-1.9 ³ | 3.22 | -0.02 | 3.20 (2.62-3.75) | 3.04 | 0.01 | 3.05 (2.58-3.50) | 2.48 | 0.01 | 2.48 (2.12-2.83) |
| | SSP1-2.6 ³ | 3.23 | -0.02 | 3.21 (2.57-3.84) | 3.56 | 0.01 | 3.58 (3.02-4.11) | 3.24 | 0.01 | 3.25 (2.81-3.67) |
| | SSP2-4.5 ³ | 3.25 | -0.02 | 3.24 (2.40-4.08) | 4.23 | 0.01 | 4.24 (3.46-5.02) | 5.32 | 0.01 | 5.33 (4.56-6.05) |
| | SSP3-7.0 ³ | 3.33 | -0.02 | 3.31 (2.30-4.31) | 4.61 | 0.01 | 4.63 (3.51-5.73) | 7.43 | 0.01 | 7.44 (6.17-8.70) |
| | SSP3-7.0_lowNTCF ⁴ | 3.40 | -0.02 | 3.48 (2.53-4.24) | 4.91 | 0.01 | 4.92 (4.13-5.70) | 7.73 | 0.01 | 7.74 (6.70-8.76) |
| | SSP3-7.0_lowNTCFCH ₄ ⁵ | 3.26 | -0.02 | 3.24 (2.40-4.09) | 4.28 | 0.01 | 4.29 (3.56-5.01) | 6.67 | 0.01 | 6.67 (5.75-7.57) |
| | SSP4-3.4 ³ | 3.26 | -0.02 | 3.24 (2.33-4.11) | 3.90 | 0.01 | 3.91 (3.06-4.68) | 4.18 | 0.01 | 4.19 (3.49-4.83) |
| | SSP4-6.0 ³ | 3.31 | -0.02 | 3.30 (2.35-4.23) | 4.44 | 0.01 | 4.46 (3.49-5.42) | 6.12 | 0.01 | 6.13 (5.21-7.02) |
| | SSP5-3.4_over ³ | 3.62 | -0.02 | 3.60 (2.82-4.40) | 5.04 | 0.01 | 5.05 (4.29-5.79) | 3.93 | 0.01 | 3.93 (3.40-4.45) |
| | SSP5-8.5 ³ | 3.61 | -0.02 | 3.59 (2.80-4.38) | 5.29 | 0.01 | 5.30 (4.44-6.17) | 9.18 | 0.01 | 9.19 (7.96-10.40) |

Notes ¹Table AII.6.8 and 6.10 in Annex II of the IPCC AR5 WG1 report (Prather et al., 2013), for which total ERF is derived from CMIP5 models and anthropogenic ERF from a simple climate model emulator. ²RCPs calculated for ERF as described in Chapter 7 Supplementary Material 7.SM.1.4. Similar to AR5, an emulator was used in AR6 to derive anthropogenic ERF (Cross Chapter Box 7.1 and Chapter 7 Supplementary Material 7.SM.1.4). Further discussion on the difference between AR5 and AR6 radiative forcing in section 4.6.2. ³ ScenarioMIP (O’Neill et al., 2016).

⁴AerChemMIP (Collins et al., 2017) ⁵ Allen et al. (2021)

References

- Adcock, K. E., Reeves, C. E., Gooch, L. J., Leedham Elvidge, E. C., Ashfold, M. J., Brenninkmeijer, C. A. M., et al. (2018). Continued increase of CFC-113a (CCl₃CF₃) mixing ratios in the global atmosphere: emissions, occurrence and potential sources. *Atmos. Chem. Phys.* 18, 4737–4751. doi:10.5194/acp-18-4737-2018.
- Allen, R. J., Horowitz, L. W., Naik, V., Oshima, N., O'Connor, F. M., Turnock, S., et al. (2021). Significant climate benefits from near-term climate forcer mitigation in spite of aerosol reductions. *Environ. Res. Lett.* doi:10.1088/1748-9326/abe06b.
- Collins, W. J., Lamarque, J.-F., Schulz, M., Boucher, O., Eyring, V., Hegglin, M. I., et al. (2017). AerChemMIP: quantifying the effects of chemistry and aerosols in CMIP6. *Geosci. Model Dev.* 10, 585–607. doi:10.5194/gmd-10-585-2017.
- Slugokenky, E. J., Nisbet, E. G., Fisher, R., and Lowry, D. (2011). Global atmospheric methane: Budget, changes and dangers. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* doi:10.1098/rsta.2010.0341.
- Droste, E. S., Adcock, K. E., Ashfold, M. J., Chou, C., Fleming, Z., Fraser, P. J., et al. (2020). Trends and emissions of six perfluorocarbons in the Northern Hemisphere and Southern Hemisphere. *Atmos. Chem. Phys.* 20, 4787–4807. doi:10.5194/acp-20-4787-2020.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., et al. (2016). Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.* 9, 1937–1958. doi:10.5194/gmd-9-1937-2016.
- Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., et al. (2019). Global emissions pathways under different socioeconomic scenarios for use in CMIP6: a dataset of harmonized emissions trajectories through the end of the century. *Geosci. Model Dev.* 12, 1443–1475. doi:10.5194/gmd-12-1443-2019.
- Hall, B. D., Dutton, G. S., Mondeel, D. J., Nance, J. D., Rigby, M., Butler, J. H., et al. (2011). Improving measurements of SF₆ for the study of atmospheric transport and emissions. *Atmos. Meas. Tech.* doi:10.5194/amt-4-2441-2011.
- Jungclaus, J. H., Bard, E., Baroni, M., Braconnot, P., Cao, J., Chini, L. P., et al. (2017). The PMIP4 contribution to CMIP6 - Part 3: The last millennium, scientific objective, and experimental design for the PMIP4 past1000 simulations. *Geosci. Model Dev.* 10, 4005–4033. doi:10.5194/gmd-10-4005-2017.
- Laube, J. C., Mohd Hanif, N., Martinerie, P., Gallacher, E., Fraser, P. J., Langenfelds, R., et al. (2016). Tropospheric observations of CFC-114 and CFC-114a with a focus on long-term trends and emissions. *Atmos. Chem. Phys.* doi:10.5194/acp-16-15347-2016.
- Laube, J. C., Newland, M. J., Hogan, C., Brenninkmeijer, C. A. M., Fraser, P. J., Martinerie, P., et al. (2014). Newly detected ozone-depleting substances in the atmosphere. *Nat. Geosci.* 7, 266–269. doi:10.1038/ngeo2109.
- Lee, D. S., Fahey, D. W., Skowron, A., Allen, M. R., Burkhardt, U., Chen, Q., et al. (2021). The contribution of global aviation to anthropogenic climate forcing for 2000 to 2018. *Atmos. Environ.* 244, 117834. doi:<https://doi.org/10.1016/j.atmosenv.2020.117834>.
- Leedham Elvidge, E., Bönisch, H., Brenninkmeijer, C. A. M., Engel, A., Fraser, P. J., Gallacher, E., et al. (2018). Evaluation of stratospheric age of air from CF4, C2F6, C3F8, CHF3, HFC-125, HFC-227ea and SF6; implications for the calculations of halocarbon lifetimes, fractional release factors and ozone depletion potentials. *Atmos. Chem. Phys.* 18, 3369–3385. doi:10.5194/acp-18-3369-2018.
- Masarie, K. A., and Tans, P. P. (2004). Extension and integration of atmospheric carbon dioxide data into a globally consistent measurement record. *J. Geophys. Res.* doi:10.1029/95jd00859.
- Matthes, K., Funke, B., Andersson, M. E., Barnard, L., Beer, J., Charbonneau, P., et al. (2017). Solar forcing for CMIP6 (v3.2). *Geosci. Model Dev.* 10, 2247–2302. doi:10.5194/gmd-10-2247-2017.
- Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., et al. (2020). The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500. *Geosci. Model Dev.* 13, 3571–3605. doi:10.5194/gmd-13-3571-2020.
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., et al. (2017). Historical greenhouse gas concentrations for climate modelling (CMIP6). *Geosci. Model Dev.* 10, 2057–2116. doi:10.5194/gmd-10-2057-2017.
- Montzka, S. A., Hall, B. D., and Elkins, J. W. (2009). Accelerated increases observed for hydrochlorofluorocarbons since 2004 in the global atmosphere. *Geophys. Res. Lett.* doi:10.1029/2008GL036475.
- Mühle, J., Trudinger, C. M., Western, L. M., Rigby, M., Vollmer, M. K., Park, S., et al. (2019). Perfluorocyclobutane (PFC-318, C4F₈) in the global atmosphere. *Atmos. Chem. Phys.* 19, 10335–10359. doi:10.5194/acp-19-10335-2019.
- Naus, S., Montzka, S. A., Pandey, S., Basu, S., Slugokenky, E. J., and Krol, M. (2019). Constraints and biases in a tropospheric two-box model of OH. *Atmos. Chem. Phys.* doi:10.5194/acp-19-407-2019.
- O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., et al. (2016). The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geosci. Model Dev.* 9, 3461–3482. doi:10.5194/gmd-9-3461-2016.
- Prather, M., Flato, G., Friedlingstein, P., Jones, C., Lamarque, J. F., Liao, H., et al. eds. (2013). “IPCC, 2013: Annex II: Do Not Cite, Quote or Distribute

- 1 Climate System Scenario Tables," in *Climate Change 2013: The Physical Science Basis. Contribution of Working*
2 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D.
3 Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia (Cambridge, United Kingdom and
4 New York, NY, USA: Cambridge University Press), 52. Available at: https://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_AnnexII_FINAL.pdf.
- 5 Prinn, R. G., Weiss, R. F., Arduini, J., Arnold, T., Langley Dewitt, H., Fraser, P. J., et al. (2018). History of chemically
6 and radiatively important atmospheric gases from the Advanced Global Atmospheric Gases Experiment
7 (AGAGE). *Earth Syst. Sci. Data.* doi:10.5194/essd-10-985-2018.
- 8 Rigby, M., Prinn, R. G., O'Doherty, S., Miller, B. R., Ivy, D., Mühle, J., et al. (2014). Recent and future trends in
9 synthetic greenhouse gas radiative forcing. *Geophys. Res. Lett.* 41, 2623–2630. doi:10.1002/2013GL059099.
- 10 Schoenenberger, F., Vollmer, M. K., Rigby, M., Hill, M., Fraser, P. J., Krumbel, P. B., Langenfelds, R. L., R., S., T., et
11 al. (2015). First observations, trends, and emissions of HCFC-31 (CH₂ClF) in the global atmosphere. *Geophys.*
12 *Res. Lett.*, 42, 7817–. doi:10.1002/2015GL064709.
- 13 Simmonds, P. G., Rigby, M., McCulloch, A., O'Doherty, S., Young, D., Mühle, J., et al. (2017). Changing trends and
14 emissions of hydrochlorofluorocarbons (HCFCs) and their hydrofluorocarbon (HFCs) replacements. *Atmos.*
15 *Chem. Phys.* 17, 4641–4655. doi:10.5194/acp-17-4641-2017.
- 16 Smith, C. J., Forster, P. M., Allen, M., Leach, N., Millar, R. J., Passerello, G. A., et al. (2018). FAIR v1.3: a simple
17 emissions-based impulse response and carbon cycle model. *Geosci. Model Dev.* 11, 2273–2297.
18 doi:10.5194/gmd-11-2273-2018.
- 19 Thornhill, G., Collins, W., Olivié, D., Skeie, R. B., Archibald, A., Bauer, S., et al. (2021a). Climate-driven chemistry
20 and aerosol feedbacks in CMIP6 Earth system models. *Atmos. Chem. Phys.* 21, 1105–1126. doi:10.5194/acp-21-
21 1105-2021.
- 22 Thornhill, G. D., Collins, W. J., Kramer, R. J., Olivié, D., Skeie, R. B., O'Connor, F. M., et al. (2021b). Effective
23 radiative forcing from emissions of reactive gases and aerosols -- a multi-model comparison. *Atmos. Chem. Phys.*
24 21, 853–874. doi:10.5194/acp-21-853-2021.
- 25 Trudinger, C. M., Etheridge, D. M., Sturrock, G. A., Fraser, P. J., Krumbel, P. B., and McCulloch, A. (2004).
26 Atmospheric histories of halocarbons from analysis of Antarctic firn air: Methyl bromide, methyl chloride,
27 chloroform, and dichloromethane. *J. Geophys. Res. D Atmos.* doi:10.1029/2004JD004932.
- 28 Velders, G. J. M., Fahey, D. W., Daniel, J. S., Andersen, S. O., and McFarland, M. (2015). Future atmospheric
29 abundances and climate forcings from scenarios of global and regional hydrofluorocarbon (HFC) emissions.
30 *Atmos. Environ.* 123, 200–209. doi:<https://doi.org/10.1016/j.atmosenv.2015.10.071>.
- 31 Vollmer, M. K., Mühle, J., Trudinger, C. M., Rigby, M., Montzka, S. A., Harth, C. M., et al. (2016). Atmospheric
32 histories and global emissions of halons H-1211 (CBrClF₂), H-1301 (CBrF₃), and H-2402 (CBrF₂CBrF₂). *J.*
33 *Geophys. Res.* doi:10.1002/2015JD024488.
- 34 Worton, D. R., Sturges, W. T., Schwander, J., Mulvaney, R., Barnola, J. M., and Chappellaz, J. (2006). 20th century
35 trends and budget implications of chloroform and related tri-and dihalomethanes inferred from firn air. *Atmos.*
36 *Chem. Phys.* doi:10.5194/acp-6-2847-2006.
- 37
- 38
- 39