

# Oceans and Coastal Ecosystems and Their Services Supplementary Material

**Coordinating Lead Authors:** Sarah R. Cooley (USA) and David S. Schoeman (Australia)

**Lead Authors:** Laurent Bopp (France), Philip Boyd (Australia/UK), Simon Donner (Canada), Shin-Ichi Ito (Japan), Wolfgang Kiessling (Germany), Paulina Martinetto (Argentina), Elena Ojea (Spain), Marie-Fanny Racault (UK/France), Björn Rost (Germany), Mette Skern-Mauritzen (Norway), Dawit Yemane Ghebrehiwet (South Africa/Eritrea)

**Contributing Authors:** Ibidun Adelekan (Nigeria), Johann D. Bell (Australia), Julia Blanchard (Australia), Jessica Bolin (Australia), William W. L. Cheung (Canada), Andrés Cisneros-Montemayor (Canada/Mexico), Sam Dupont (Sweden/Belgium), Stephanie Dutkiewicz (USA), Thomas Frölicher (Switzerland), Juan Diego Gaitán-Espitia (Hong Kong, Special Administrative Region, China/Colombia), Jorge García Molinos (Japan/Spain), Bruce Glavovic (New Zealand/South Africa), Helen Gurney-Smith (Canada), Marjolijn Haasnoot (The Netherlands), Rebecca Harris (Australia), Stephanie Henson (UK), Manuel Hidalgo (Spain), Elisabeth Holland (Fiji), Robert Kopp (USA), Rebecca Kordas (UK/USA), Lester Kwiatkowski (France/UK), Judy Lawrence (New Zealand), Nadine Le Bris (France), Gonéri Le Cozannet (France), Salvador E. Lluch-Cota (Mexico), Cheryl Logan (USA), Felix Mark (Germany), Yunus Mgaya (Tanzania), Coleen Moloney (South Africa), Aditi Mukherji (Nepal), Norma Patricia Muñoz Sevilla (Mexico), Patrick Nunn (Australia), Dieter Piepenburg (Germany), Gregoire Randin (Fiji/France/Switzerland), Nussaibah B. Raja (Germany/Mauritius), Anusha Rajkaran (South Africa), Anthony J. Richardson (Australia), Stephanie Roe (Philippines/USA), Raquel Ruiz Diaz (Spain), Diana Salili (Vanuatu), Jean-Baptiste Sallée (France), Kylie Scales (Australia/UK), Daniela Schmidt (UK/Germany), Michelle Scobie (Trinidad and Tobago), Craig T. Simmons (Australia), Chandni Singh (India), Aimée Slangen (The Netherlands), Seree Supratid (Thailand), Olivier Torres (France), Andrew Yool (UK)

**Review Editors:** Karim Hilmi (Morocco) and Lisa Levin (USA)

**Chapter Scientist:** Jessica Bolin (Australia)

**This Supplementary Material should be cited as:**

Cooley, S., D. Schoeman, L. Bopp, P. Boyd, S. Donner, D.Y. Ghebrehiwet, S.-I. Ito, W. Kiessling, P. Martinetto, E. Ojea, M.-F. Racault, B. Rost, and M. Skern-Mauritzen, 2022: Oceans and Coastal Ecosystems and Their Services Supplementary Material. In: *Climate Change 2022: Impacts, Adaptation and Vulnerability*. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegria, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Available from <https://www.ipcc.ch/report/ar6/wg2/>.

## Table of Contents

|                   |  |    |
|-------------------|--|----|
| <b>SM3.1</b>      | <b>Section 3.1</b>   | 3  |
| SM3.1.1           | SROCC Burning Embers   | 3  |
| <b>SM3.2</b>      | <b>Section 3.3</b>   | 4  |
| SM3.2.1           | Combined Climate Stressors   | 4  |
| SM3.2.2           | Understanding Sources' Uncertainty in Climate Projections from Marine Ecological Models        | 4  |
| <b>SM3.3</b>      | <b>Section 3.4</b>   | 5  |
| SM3.3.1           | Assessment of the Impact of Storms and Tropical Cyclones on Estuaries and Soft-Sediment Coasts | 5  |
| SM3.3.2           | Additional Background for the Assessment of Semi-Enclosed Seas                                 | 5  |
| SM3.3.3           | Calculating Changes in Phenology Shifts  | 5  |
| <b>SM3.4</b>      | <b>Section 3.5</b>   | 23 |
| <b>SM3.5</b>      | <b>Section 3.6</b>   | 26 |
| SM3.5.1           | Assessment of the Potential of Marine and Coastal Adaptation Solutions                         | 26 |
| SM3.5.2           | Fisheries Adaptation to Climate Change   | 30 |
| SM3.5.3           | Multilateral Environmental Agreements and Climate Change                                       | 40 |
| SM3.5.4           | Data Supporting Figure 3.25  | 42 |
| SM3.5.5           | Data Supporting Figure 3.26  | 45 |
| <b>References</b> |  | 55 |



**SM3.1 Section 3.1****SM3.1.1 SROCC Burning Embers****Impacts and risks to ocean ecosystems from climate change**

Figure SM3.1 | Projected impacts and risks are shown for ocean regions and ecosystems, as assessed by SROCC (IPCC, 2019).

## SM3.2 Section 3.3

### SM3.2.1 Combined Climate Stressors

#### The effects of single and combined stressors on the performance of marine ectothermic animals



**Figure SM3.2 | The effects of single and combined stressors on the performance of marine ectothermic animals.** Performance relates to survival, development, condition, growth, calcification and metabolism, which ultimately modulate population extinction risk. Solid symbols and vertical bars indicate effect size  $\pm 95\%$  confidence (extremely likely) intervals. The interaction of rising temperature and oxygen limitation has the most detrimental effect on organismal performance. (Modified from Reddin et al., 2020).

### SM3.2.2 Understanding Sources' Uncertainty in Climate Projections from Marine Ecological Models

The use of global and regional marine ecosystem and fisheries models (e.g., in ensembles such as Fish-MIP; Tittensor et al., 2018) provides opportunities to investigate the influence of different sources of uncertainty on model projections at different temporal and spatial scales. Fish-MIP global modelling experiments have shown that different model assumptions regarding the effects of temperature and changes in primary and secondary production, combined with the degree of food-web complexity and feedbacks, can lead to substantial differences in marine animal biomass under future projections under separate and combined physical and biogeochemical drivers (Heneghan et al., 2021). These drivers include directional differences in projections, particularly at smaller spatial scales (Tittensor et al., 2018), and in comparison with regional marine ecosystem models (Pethybridge et al., 2020). It is clear that much more work is needed to advance the nature of coupling physical, biogeochemical and ecological models as well as appropriate combinations of models and data across different scales.

Global Fish-MIP model ensemble experiments have been restricted to the use of only a few ESMs which does not provide enough information to assess whether there is a decrease in the uncertainty of Fish-MIP models compared with CMIP ensembles (Lotze et al., 2019). Single-model ensembles have been used, however, to assess the relative influence of different sources of uncertainty on impacts of regional projections, spanning a wider range of ESM outputs (Payne et al., 2016). For example, using a regional model of the Bering Sea, Reum et al. (2020) showed that in the short- to medium term, ecological processes contributed substantially to model uncertainty, but that ESM and scenario uncertainty became the overriding sources of variation in the long term. Similar results have been found with species distribution models at the global scale, but when examined for single species or regions, internal variability of ecological models also can be very large (Cheung et al., 2016a; Cheung et al., 2016b), highlighting the need to track multiple sources of uncertainty for confidence assessment in models.

## SM3.3 Section 3.4

### SM3.3.1 Assessment of the Impact of Storms and Tropical Cyclones on Estuaries and Soft-Sediment Coasts

Estuarine ecosystems and their phytoplankton (Carrasco Navas-Parejo et al., 2020), vegetation (Mo et al., 2020), fish (Matich et al., 2020) and birds (Field et al., 2019; Wilkinson et al., 2019) can be resilient to the impacts of storms and tropical cyclones (*medium confidence*). Nevertheless, the passage of storms can also manifest subtly in, for example, reduced body condition of juvenile fish (Matich et al., 2020), and can also have counterintuitive effects by, for example, reducing erosion rates by moving sediments onshore (Wang et al., 2020). In some instances, heavy precipitation can flush estuaries, resulting in net erosion; in others, runoff from adjacent land can increase nutrient loads, causing or exacerbating eutrophication, stimulating HABs (Philips et al., 2020) and sometimes causing large-scale marine mammal, bird and fish kills (Adams et al., 2019).

Effects of tropical cyclones on mangroves are variable (*high confidence*). They range from beneficial (Castañeda-Moya et al., 2020; Feher et al., 2020), through minor damage (Armitage et al., 2019) or recovery over a period of less than a decade (Branoff, 2020), culminating in regime shifts involving peat collapse and transition to mudflats (Chambers et al., 2019; Osland et al., 2020). Salt-marsh vegetation is more resilient to intense storms (*medium confidence*) (Armitage et al., 2019; Mo et al., 2020), with sediments showing limited amounts of long-term erosion, and sometimes even accretion (Xie et al., 2017), especially when suspended sediments are not advected too far away from the site of erosion (Leonardi et al., 2018).

On sandy beaches (Section 3.4.2.6), although extreme storms can both remove and create habitat for shorebirds, losses of existing habitat are generally small, even when considering only the impacted portion of species' distributions (Maslo et al., 2019). When considering the full species' distributions, impacts are often negligible (Field et al., 2019). Projected loss of beach habitat to SLR and urbanisation remains a substantial risk, however.

### SM3.3.2 Additional Background for the Assessment of Semi-Enclosed Seas

Recent warming and changes in other environmental conditions in semi-enclosed seas (SES), with ecological and biogeochemical ramifications, have been attributed to climate change (*high confidence*) (e.g., Adloff et al., 2015; Shirvani et al., 2015; The BACC II Author Team, 2015). The Persian Gulf, characterised by extreme seasonal fluctuations in temperature and salinity, has shown an increase in frequency of extreme events, increases in salinity and declines in oxygen content. The latter changes, combined with substantial reduction in freshwater discharge and an increase in coastal eutrophication, have triggered changes in biogeochemical cycles (*high confidence*) (Al-Said et al., 2018; Ben-Hasan et al., 2018; Al-Yamani and Naqvi, 2019; Ben-Hasan and Christensen, 2019). In the Red Sea, ocean acidification and MHWs are the main climate-

induced drivers, along with warming (*high confidence*) (Steiner et al., 2018; Genevier et al., 2019). Increasing MHW frequency, decreases in precipitation, accelerating SLR and extreme winter weather events have been reported for the Mediterranean Sea (*high confidence*) (Adloff et al., 2015; Cramer et al., 2018; Darmaraki et al., 2019). Important changes in thermohaline circulation have been reported in the Black Sea, including erosion of the cold intermediate layer that controls the important exchange of oxygen between surface and deep water masses (*high confidence*) (Cannaby et al., 2015; Miladinova et al., 2017). In the Baltic Sea, changes in rainfall and river runoff have been described, with a decreased influx of seawater (from the North Sea) and prolonged hypoxia in deeper parts of the basin representing major stressors for different ecosystem components (*high confidence*) (The BACC II Author Team, 2015; Räisänen, 2017).

The size and number of OMZs are increasing worldwide and in most SES (*high confidence*) (Global Ocean Oxygen Network, 2018), with growing impacts on fish species and ecosystem function. In the Persian Gulf and Red Sea, increasing nutrient loads associated with coastal activities and warming have increased the size of OMZs (*high confidence*) (Al-Said et al., 2018; Lachkar et al., 2019). These OMZs represent an even greater problem in the Black and Baltic Seas, with broad implications for ecosystem function and services (e.g., Levin et al., 2009), especially where actions to reduce nutrient loading from land have been unable to reduce the OMZ coverage (*high confidence*) (Carstensen et al., 2014; Miladinova et al., 2017; Global Ocean Oxygen Network, 2018). In the Baltic Sea, OMZs are affecting the spawning areas of cod, *Gadus morhua* (*high confidence*) (Hinrichsen et al., 2016), while in the Black Sea, the combined effect of OMZs and warming is influencing the distribution and physiology of fish species, and their migration and schooling behaviour in the overwintering grounds (*medium confidence*) (Güraslan et al., 2017). Cascading effects on food webs have been reported in the Baltic, where detrimental effects of changing oxygen levels on zooplankton production as well as pelagic and piscivorous fish are influencing seasonal succession and species composition of phytoplankton (*high confidence*) (Viitasalo et al., 2015).

### SM3.3.3 Calculating Changes in Phenology Shifts

#### SM3.3.3.1 Database

This analysis updates the database of Poloczanska et al. (2013), which had 109 time series from 31 studies and was used in AR5 WGI Chapter 30 (Hoegh-Guldberg et al., 2014). The new studies were found by searching ISI Web of Science and Google Scholar using keywords 'phenology' and 'marine'. As in the original database, all observations had to meet three criteria:

- Criterion 1. Time series had to be at least 19 years in duration to minimise bias associated with short-term responses to natural climate variability (median time-series duration was 45 years).
- Criterion 2. The end date of the time series had to be 1990 or later.
- Criterion 3. Studies had to directly test, or at a minimum discuss, their results in relation to expected impacts of climate change (Poloczanska et al., 2013).

This process resulted in the addition of 306 time series from 79 studies.

Each observation in the database was a time series of a species or a group of species (e.g., total zooplankton). Observations included cases where phenological responses were consistent with climate change (earlier occurrence with warming), inconsistent with climate change or there was no change. Data were included from time series that were continuous ( $n = 388$ ), intermittent ( $n = 14$ ) and from two periods in time ( $n = 13$ ). Quantitative estimates of shifts in phenology were taken directly from the published papers, calculated from information in tables or in the supplementary materials, or digitised from figures. We used data reported as days per decade to examine the mean rates of shifts.


To avoid duplication and to minimise spatial autocorrelation, the latitude and longitude of each observation was used to assign each observation to a Longhurst Province, a commonly used global bioregionalisation (Longhurst et al., 1995). We considered time series of the same species in different Longhurst Provinces as unique observations, but where we found multiple time series for the same species in a Longhurst Province, we used the mean phenological shift and randomly selected an observation for consistency (see 'Analysis' below). The data used in the analysis below are available in Table SM3.1.

3SM

### SM3.3.3.2 Analysis

To estimate rates of consistency with climate change among taxonomic groups (taxa), we analysed the observations using a generalised linear model with a binomial error structure and a binary response variable (0 = Not consistent; 1 = Consistent) ( $n = 413$ ). To analyse the magnitude and direction of observed phenological shifts (days per decade) by different taxonomic groups, we used a linear model with a normal error structure ( $n = 312$ ). In both models, the same four predictors were used. First was the taxonomic group (with levels for Phytoplankton, Zooplankton, Meroplankton, Benthic invertebrates, Plants, Fish, Reptiles, Seabirds, Mammals), so we could test for differences across groups in Consistency and the magnitude and direction of phenological shift. We included the remaining predictors to account for potential issues associated with publication bias (Parmesan and Yohe, 2003). The second predictor in the models, number of species in each study, was included because previous analyses have found that studies that included few species (e.g., single-species studies) tend to have a larger effect size (Parmesan and Yohe, 2003). The third predictor was the duration of the study, because shorter studies might be more likely to report more consistent impacts of climate change and greater phenology shifts. The final predictor included was the mid-year of the time series, to test whether more-

### Observed responses to climate change based on a systematic review of the Web of Science for marine phenology studies longer than 19 years in duration



**Figure SM3.3 | Observed responses to climate change based on a systematic review of the Web of Science for marine phenology studies longer than 19 years in duration.** Error envelopes indicate 95% (*extremely likely*) confidence interval.

(a) Proportion of phenology observations (showing means and *extremely likely* range) that are attributed to climate change (i.e., generally showing earlier timing) by duration of study in years (adjusted for taxon). The dashed line at 0.5 indicates random chance.

(b) Shifts in timing that are attributed to climate change by duration of study (adjusted for taxon). The dashed line at 0 indicates no shift. Negative shifts are earlier, positive shifts are later. The observations summarised in panels (a) and (b) have been collected over the global oceans (see locations in Figure 3.16a) and include  $n = 297,277$  observations of phenology shifts that are attributed (at least partly) to climate change, and  $n = 1168$  observations of phenology shifts that are inconsistent with climate change (see Section 3.4.3.2).

recent time series might be more likely to observe more consistent impacts of climate change and greater phenology shifts as climate change accelerates. For modelling consistency, we removed from the model any taxonomic group that had no variance (i.e., all time

series were consistent with climate change) because their inclusion destabilised error estimation. These points were plotted in the final output without error (since it is impossible to determine error without variance).

**Table SM3.1** | Data used to create Figure 3.16 in Section 3.4.3, with Longhurst codes included. Shift is in days per decade

| Reference                     | Database | Long-hurst | Latitude | Lon-gitude | Group           | Species  | Last year | Median year | Duration | Number of replicates | Con-sistent | Shift |
|-------------------------------|----------|------------|----------|------------|-----------------|--|-----------|-------------|----------|----------------------|-------------|-------|
| Atkinson et al. (2015)        | Current  | NECS       | 50.25    | -4.21      | Holozooplankton | <i>Acartia clausi</i>                            | 2012      | 2000        | 25       | 24                   | 1           | -6.4  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00       | Holozooplankton | <i>Acartia</i> spp.                              | 1998      | 1982        | 45       | 49                   | 1           | -0.81 |
| Costello et al. (2006)        | AR5      | NWCS       | 41.50    | -71.35     | Holozooplankton | <i>Acartia tonsa</i>                             | 2003      | 1977        | 53       | 2                    | 1           | 0     |
| Greve et al. (1996)           | Current  | NECS       | 54.19    | 7.90       | Holozooplankton | <i>Actinotrocha</i> spp.                         | 1994      | 1984        | 21       | 25                   | 1           | NA    |
| Descamps et al. (2019)        | Current  | PSAW       | 52.35    | -176.93    | Seabirds        | <i>Aethia cristatella</i>                        | 2015      | 2001.5      | 28       | 16                   | 0           | 0     |
| Descamps et al. (2019)        | Current  | PSAW       | 52.35    | -176.93    | Seabirds        | <i>Aethia psittacula</i>                         | 2015      | 2003        | 25       | 16                   | 1           | -0.08 |
| Descamps et al. (2019)        | Current  | PSAW       | 52.35    | -176.93    | Seabirds        | <i>Aethia pusilla</i>                            | 2015      | 1998.5      | 34       | 16                   | 0           | 0.09  |
| Descamps et al. (2019)        | Current  | PSAW       | 52.35    | -176.93    | Seabirds        | <i>Aethia pygmaea</i>                            | 2015      | 2001.5      | 28       | 16                   | 1           | -0.18 |
| Greve et al. (1996)           | Current  | NECS       | 54.19    | 7.90       | Holozooplankton | <i>Aglantha digitale</i>                         | 1994      | 1984        | 21       | 25                   | 1           | NA    |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94       | Fish            | <i>Agonus cataphractus</i>                       | 2013      | 1986.5      | 54       | 30                   | 1           | NA    |
| Greve et al. (1996)           | Current  | NECS       | 54.19    | 7.90       | Holozooplankton | <i>Alaurina</i> spp.                             | 1994      | 1984        | 21       | 25                   | 1           | NA    |
| Wanless et al. (2009)         | AR5      | NECS       | 56.18    | -2.56      | Seabirds        | <i>Alca torda</i>                                | 2006      | 1990.5      | 36       | 7                    | 0           | -0.04 |
| Descamps et al. (2019)        | Current  | SARC       | 70.37    | 31.13      | Seabirds        | <i>Alca torda</i>                                | 2013      | 1997.5      | 32       | 16                   | 1           | -0.03 |
| Bucci et al. (2020)           | Current  | NWCS       | 44.90    | -66.70     | Phytoplankton   | <i>Alexandrium catenella</i>                     | 2014      | 2001        | 27       | 1                    | 1           | -7.5  |
| Atkinson et al. (2015)        | Current  | NECS       | 50.25    | -4.21      | Phytoplankton   | <i>All other colourless dinoflagellates</i>      | 2012      | 2000        | 25       | 24                   | 0           | 6.4   |
| Moe et al. (2009)             | AR5      | ARCT       | 77.00    | 15.55      | Seabirds        | <i>Alle alle</i>                                 | 2012      | 1992.5      | 46       | 2                    | 1           | -0.66 |
| Lombardo et al. (2019)        | Current  | NWCS       | 36.00    | -76.50     | Fish            | <i>Alosa aestivalis</i>                          | 2016      | 1994.5      | 44       | 1                    | 1           | -4.33 |
| Chevillot et al. (2017)       | Current  | NECS       | 45.42    | -0.86      | Fish            | <i>Alosa alosa</i>                               | 2010      | 1997.5      | 26       | 7                    | 1           | NA    |
| Chevillot et al. (2017)       | Current  | NECS       | 45.42    | -0.86      | Holozooplankton | <i>Alosa bifilosa</i>                            | 2010      | 1997.5      | 26       | 7                    | 1           | NA    |
| Chevillot et al. (2017)       | Current  | NECS       | 45.42    | -0.86      | Fish            | <i>Alosa fallax</i>                              | 2010      | 1997.5      | 26       | 7                    | 1           | NA    |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00     | Fish            | <i>Alosa pseudoharengus</i>                      | 2015      | 1992.5      | 48       | 35                   | 1           | -3    |
| Cobb (2020)                   | Current  | NWCS       | 43.27    | -70.98     | Fish            | <i>Alosa pseudoharengus and Alosa aestivalis</i> | 2016      | 1997.5      | 38       | 1                    | 1           | -3.5  |
| Chevillot et al. (2017)       | Current  | NECS       | 45.42    | -0.86      | Fish            | <i>Alosa regius</i>                              | 2010      | 1997.5      | 26       | 7                    | 0           | NA    |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                            | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|------------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Quinn and Adams, 1996)           | Current  | CCAL       | 45.64    | -121.94   | Fish            | <i>Alosa sapidissima</i>           | 1992      | 1965        | 55       | 1                    | 1          | -6.89  |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Alosa sapidissima</i>           | 2012.5    | 1993.25     | 48       | 35                   | 1          | -4     |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Amblyraja radiate</i>           | 2014      | 1990.5      | 48       | 35                   | 0          | NA     |
| Burthe et al. (2012)             | Current  | NECS       | 56.50    | -1.50     | Fish            | <i>Ammodytes marinus</i>           | 2006      | 1994.5      | 24       | 4                    | 0          | 2.58   |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Ammodytes tobianus</i>          | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Holozooplankton | <i>Amphidinium crassum</i>         | 2016      | 2004.5      | 24       | 29                   | 1          | -7.24  |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Anarhichas lupus</i>            | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Meroplankton    | <i>Anemone larvae</i>              | 2012      | 2000        | 25       | 24                   | 1          | -1.6   |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Anguilla anguilla</i>           | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Thaxton et al. (2020)            | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Anguilla rostrata</i>           | 2013      | 1999.5      | 28       | 9                    | 1          | -0.1   |
| Chambers et al. (2014)           | Current  | AUSW       | -28.77   | 113.85    | Seabirds        | <i>Anous stolidus</i>              | 2010      | 2000.5      | 20       | 4                    | 0          | NA     |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Aphanizomenon spp.</i>          | 2016      | 2004.5      | 24       | 29                   | 1          | -3.03  |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Holozooplankton | Appendicularians                   | 2012      | 2000        | 25       | 24                   | 0          | 2      |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Aptenodytes forsteri</i>        | 2004      | 1977        | 55       | 9                    | 0          | 0.31   |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | Arachnactes                        | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Argentina sialis</i>            | 2008      | 1979.5      | 58       | 43                   | 0          | 2      |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Argyropelecus sladeni</i>       | 2008      | 1979.5      | 58       | 43                   | 1          | -5.85  |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Arnoglossus laterna</i>         | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | <i>Asterias rubens</i>             | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Asterionellopsis glacialis</i>  | 2011      | 1985.166    | 45       | 49                   | 1          | -9.5   |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Atherina presbyter</i>          | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Van Walraven et al. (2015)       | Current  | NECS       | 53.57    | 6.94      | Holozooplankton | <i>Aurelia aurita</i>              | 2010      | 1985        | 51       | 2                    | 1          | -7.66  |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | Autotrophic dinoflagellate biomass | 2012      | 2000        | 25       | 24                   | 0          | 1.6    |
| Ramp et al. (2015)               | Current  | NWCS       | 49.90    | -64.50    | Mammals         | <i>Balaenoptera physalus</i>       | 2010      | 1997        | 27       | 2                    | 1          | -10.37 |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Bathylagus pacificus</i>        | 2008      | 1979.5      | 58       | 43                   | 0          | 4.6    |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Bathylagus wesethi</i>          | 2008      | 1979.5      | 58       | 43                   | 0          | 4.4    |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Belone belone</i>               | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                       | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|-------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Schlüter et al. (2010)           | AR5      | NECS       | 54.18    | 7.90      | Holozooplankton | <i>Beroe gracilis</i>         | 2004      | 1989.5      | 30       | 2                    | 1          | -9.33  |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | Bodonids                      | 2012      | 2000        | 25       | 24                   | 1          | -5.2   |
| Thaxton et al. (2020)            | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Brevoortia tyrannus</i>    | 2013      | 1999.5      | 28       | 9                    | 1          | -10.9  |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Brosme brosme</i>          | 2014      | 1990.5      | 48       | 35                   | 0          | NA     |
| McGinty et al. (2011)            | Current  | NADR       | 49.50    | -9.00     | Holozooplankton | <i>Calanus finmarchicus</i>   | 2008      | 1983        | 51       | 7                    | 1          | -1.56  |
| McGinty et al. (2011)            | Current  | NASE       |          | -15.00    | Holozooplankton | <i>Calanus finmarchicus</i>   | 2008      | 1983        | 51       | 7                    | 1          | -0.59  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Calanus finmarchicus</i>   | 2006.4    | 1984.7      | 45       | 49                   | 1          | -1.84  |
| McGinty et al. (2011)            | Current  | SARC       | 60.00    | -10.00    | Holozooplankton | <i>Calanus finmarchicus</i>   | 2008      | 1983        | 51       | 7                    | 1          | -2.97  |
| Persson et al. (2012)            | Current  | SARC       | 66.33    | 33.66     | Holozooplankton | <i>Calanus glacialis</i>      | 2010      | 1987        | 47       | 2                    | 1          | NA     |
| McGinty et al. (2011)            | Current  | NADR       | 49.50    | -9.00     | Holozooplankton | <i>Calanus helgolandicus</i>  | 2008      | 1983        | 51       | 7                    | 1          | -2.3   |
| McGinty et al. (2011)            | Current  | NASE       | 41.00    | -15.00    | Holozooplankton | <i>Calanus helgolandicus</i>  | 2008      | 1983        | 51       | 7                    | 1          | -1.48  |
| McGinty et al. (2011)            | Current  | NECS       | 53.00    | -6.00     | Holozooplankton | <i>Calanus helgolandicus</i>  | 2008.4    | 1989.9      | 39       | 7                    | 1          | 0.94   |
| McGinty et al. (2011)            | Current  | SARC       | 60.00    | -10.00    | Holozooplankton | <i>Calanus helgolandicus</i>  | 2008      | 1983        | 51       | 7                    | 1          | -2.08  |
| Burthe et al. (2012)             | Current  | NECS       | 56.50    | -1.50     | Holozooplankton | <i>Calanus I-IV</i>           | 2006      | 1994.5      | 24       | 4                    | 1          | -6.33  |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Callionymus lyra</i>       | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Cancer borealis</i>        | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Cancer irroratus</i>       | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Candacia armata</i>        | 2007      | 1990        | 45       | 49                   | 1          | 0.73   |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Meroplankton    | <i>Carcinus maenas</i>        | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Monsinjon et al. (2019)          | Current  | BRAZ       | -12.57   | -38.00    | Reptiles        | <i>Caretta caretta</i>        | 2014      | 2001.5      | 26       | 1                    | 1          | -2.2   |
| Mazaris et al. (2008)            | AR5      | MEDI       | 37.73    | 20.89     | Reptiles        | <i>Caretta caretta</i>        | 2002      | 1993        | 19       | 1                    | 1          | -11.58 |
| Hawkes et al. (2007)             | AR5      | NWCS       | 33.83    | -77.95    | Reptiles        | <i>Caretta caretta</i>        | 2005      | 1994        | 23       | 1                    | 1          | 0      |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Catharacta maccormicki</i> | 2004      | 1977        | 55       | 9                    | 0          | 0.11   |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Centropages hamatus</i>    | 2002      | 1980        | 45       | 49                   | 1          | -3.13  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Centropages typicus</i>    | 2008.666  | 1990.333    | 45       | 49                   | 1          | -4.47  |

| Reference                     | Database | Long-hurst | Latitude | Longitude | Group           | Species                          | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|-------------------------------|----------|------------|----------|-----------|-----------------|----------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Centropristes striata</i>     | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Sauve et al. (2019)           | Current  | BPLR       | 71.33    | -155.68   | Seabirds        | <i>Cephus grylle mandtii</i>     | 2017      | 1996.5      | 42       | 1                    | 1          | -1.86  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Ceratium furca</i>            | 2011      | 1985.166    | 45       | 49                   | 1          | -2.49  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Ceratium fusus</i>            | 2011      | 1985.166    | 45       | 49                   | 1          | -8.22  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Ceratium horridum</i>         | 2009      | 1983.5      | 45       | 49                   | 1          | -4.84  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Ceratium lineatum</i>         | 2009      | 1983.5      | 45       | 49                   | 1          | -4.08  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Ceratium longipes</i>         | 2009      | 1983.5      | 45       | 49                   | 0          | -0.21  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Ceratium macroceros</i>       | 2009      | 1983.5      | 45       | 49                   | 1          | -8.79  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Ceratium tripos</i>           | 2009      | 1983.5      | 45       | 49                   | 1          | -7.27  |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Ceratoscopelus townsendi</i>  | 2008      | 1979.5      | 58       | 43                   | 1          | -0.7   |
| Scharfe and Wiltshire (2019)  | Current  | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Ceratulina pelagica</i>       | 2015      | 1988.5      | 54       | 8                    | 1          | 0.39   |
| Bertram et al. (2001)         | AR5      | ALSK       | 50.87    | -129.08   | Seabirds        | <i>Cerorhinca monocerata</i>     | 2007      | 1996        | 25       | 4                    | 1          | -4.44  |
| Chivers et al. (2020)         | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Chaetoceros (Hyalochaete)</i> | 2016      | 1987        | 59       | 12                   | 1          | -1.75  |
| Chivers et al. (2020)         | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Chaetoceros (Phaeoceros)</i>  | 2016      | 1987        | 59       | 12                   | 0          | -0.34  |
| Scharfe and Wiltshire (2019)  | Current  | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Chaetoceros spp.</i>          | 2015.333  | 1993.833    | 54       | 8                    | 1          | -10.43 |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | Chaetognaths                     | 2007      | 1990        | 45       | 49                   | 1          | -2.13  |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Chauliodus macouni</i>        | 2008      | 1979.5      | 58       | 43                   | 1          | -0.6   |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Chelidonichthys lucerna</i>   | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |
| Émond et al. (2020)           | Current  | NWCS       | 47.50    | -63.00    | Meroplankton    | <i>Chionoecetes opilio</i>       | 2012      | 1997        | 31       | 2                    | 1          | NA     |
| Guinder et al. (2010)         | Current  | FKLD       | -38.72   | -62.27    | Phytoplankton   | <i>Chl-a</i>                     | 2007      | 1992.5      | 30       | 1                    | 1          | -12.33 |
| Philippart et al. (2003)      | AR5      | NECS       | 53.00    | 4.80      | Phytoplankton   | <i>Chl-a</i>                     | 2011.5    | 1998.625    | 29       | 2                    | 1          | -0.42  |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Chromis punctipinnis</i>      | 2008      | 1979.5      | 58       | 43                   | 0          | 3      |
| Van Walraven et al. (2015)    | Current  | NECS       | 53.57    | 6.94      | Holozooplankton | <i>Chrysaora hysoscella</i>      | 2010      | 1985        | 51       | 2                    | 1          | -10.01 |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Chrysochromulina spp.</i>     | 2016      | 2004.5      | 24       | 29                   | 0          | 1.94   |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                         | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|---------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Meroplankton    | <i>Cirripede ciprid</i>         | 2007      | 1990        | 45       | 49                   | 1          | 1.52   |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Meroplankton    | <i>Cirripede nauplii</i>        | 2003      | 1992        | 21       | 25                   | 1          | 1.6    |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Citharichthys sordidus</i>   | 2008      | 1979.5      | 58       | 43                   | 0          | 2.95   |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Citharichthys stigmatus</i>  | 2008      | 1979.5      | 58       | 43                   | 0          | 16     |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Clione limacina</i>          | 2002      | 1980        | 45       | 49                   | 0          | 3.21   |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Clupea harengus</i>          | 2004.5    | 1985.75     | 54       | 30                   | 1          | -7.69  |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Clupea harengus</i>          | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | Coccolithophore                 | 2012      | 2000        | 25       | 24                   | 0          | 9.6    |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | Colourless flagellates          | 2012      | 2000        | 25       | 24                   | 1          | -7.2   |
| Burthe et al. (2012)             | Current  | NECS       | 56.50    | -1.50     | Holozooplankton | <i>Copepod nauplii</i>          | 2009      | 1997.25     | 24       | 4                    | 1          | -2.11  |
| Edwards et al. (2006)            | AR5      | NECS       | 56.00    | 3.00      | Holozooplankton | Copepods                        | 2005      | 1981.5      | 48       | 1                    | 1          | NA     |
| Edwards et al. (2006)            | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Corycaeus</i> spp.           | 2002      | 1980        | 45       | 49                   | 0          | 5.57   |
| Philippart et al. (2003)         | AR5      | NECS       | 53.00    | 4.80      | Invertebrates   | <i>Crangon crangon</i>          | 2001      | 1992        | 19       | 2                    | 1          | -30.39 |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Meroplankton    | Crangonidae                     | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Cherkiss et al. (2020)           | Current  | CARB       | 25.16    | -80.83    | Reptiles        | <i>Crocodylus acutus</i>        | 2016      | 1998.25     | 39       | 1                    | 1          | -5.8   |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Holozooplankton | Ctenophores                     | 2012      | 2000        | 25       | 24                   | 1          | -26.4  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | Cumacea                         | 2002      | 1980        | 45       | 49                   | 1          | -2.77  |
| Hosia et al. (2014)              | Current  | NECS       | 58.42    | 8.75      | Holozooplankton | <i>Cyanea</i> spp.              | 2010.5    | 1993.5      | 19       | 1                    | 0          | 2.59   |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Cyclopterus lumpus</i>       | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Cyclothona signata</i>       | 2008      | 1979.5      | 58       | 43                   | 1          | -4.35  |
| Chivers et al. (2020)            | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Cylindrotheca closterium</i> | 2016      | 1987        | 59       | 12                   | 0          | -0.68  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Meroplankton    | <i>Cyphonautes larvae</i>       | 2002      | 1980        | 45       | 49                   | 1          | -2.72  |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Danaphos oculatus</i>        | 2008      | 1979.5      | 58       | 43                   | 1          | -9.3   |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Daption capense</i>          | 2004      | 1977        | 55       | 9                    | 0          | 1.28   |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Meroplankton    | <i>Decapod larvae</i>           | 2007      | 1990        | 45       | 49                   | 1          | -0.99  |

| Reference                     | Database | Long-hurst | Latitude | Longitude | Group           | Species                           | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|-------------------------------|----------|------------|----------|-----------|-----------------|-----------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Hauser et al. (2017)          | Current  | BPLR       | 69.47    | -171.69   | Mammals         | <i>Delphinapterus leucas</i>      | 2012      | 2002.5      | 20       | 1                    | 1          | 11.33  |
| Scharfe and Wiltshire (2019)  | Current  | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Detonula pumila</i>            | 2015      | 1988.5      | 54       | 8                    | 1          | -4.33  |
| Wiltshire and Manly (2004)    | AR5      | NECS       | 54.18    | 7.90      | Phytoplankton   | Diatoms                           | 2009      | 1994.125    | 41       | 1                    | 1          | -4.55  |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Dicentrarchus labrax</i>       | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Dinobryon faculiferum</i>      | 2016      | 2004.5      | 24       | 29                   | 1          | -16.3  |
| Tunin-Ley et al. (2009)       | AR5      | NASE       | 41.00    | -14.00    | Phytoplankton   | Dinoflagellates                   | 2005      | 1956.5      | 98       | 1                    | 1          | NA     |
| Hjerne et al. (2019)          | Current  | NECS       | 58.80    | 17.63     | Phytoplankton   | Dinoflagellates                   | 2011      | 1997.25     | 35       | 1                    | 1          | -3.76  |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Dinophysis acuminata</i>       | 2016      | 2004.5      | 24       | 29                   | 1          | -16.37 |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Dinophysis</i> spp.            | 2009      | 1983.5      | 45       | 49                   | 1          | -4.73  |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Diogenichthys attanticus</i>   | 2008      | 1979.5      | 58       | 43                   | 1          | -2.3   |
| Atkinson et al. (2015)        | Current  | NECS       | 50.25    | -4.21     | Holozooplankton | <i>Ditrichocorycaeus anglicus</i> | 2012      | 2000        | 25       | 24                   | 0          | 0.8    |
| Chivers et al. (2020)         | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Ditylum brightwelli</i>        | 2015.5    | 1987.75     | 59       | 12                   | 1          | -2.94  |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Dolichospermum</i> spp.        | 2016      | 2004.5      | 24       | 29                   | 1          | -27.45 |
| Langan et al. (2021)          | Current  | NWCS       | 41.50    | -71.40    | Fish            | <i>Doryteuthis pealeii</i>        | 2016      | 1987.5      | 58       | 5                    | 1          | -2.88  |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Holozooplankton | <i>Ebria tripartita</i>           | 2016      | 2004.5      | 24       | 29                   | 1          | -24.99 |
| Greve et al. (1996)           | Current  | NECS       | 54.19    | 7.90      | Meroplankton    | <i>Echinocardium cordatum</i>     | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Meroplankton    | <i>Echinoderm larvae</i>          | 2007      | 1990        | 45       | 49                   | 1          | -7.16  |
| Chevillot et al. (2017)       | Current  | NECS       | 45.42    | -0.86     | Fish            | <i>Engraulis encrasicolus</i>     | 2011.5    | 1992        | 26       | 7                    | 1          | NA     |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Engraulis mordax</i>           | 2008      | 1979.5      | 58       | 43                   | 1          | -3     |
| Chivers et al. (2020)         | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Eucampia zodiacus</i>          | 2015.666  | 1987.5      | 59       | 12                   | 0          | 6.23   |
| Hindell et al. (2012)         | Current  | SANT       | -54.62   | 158.85    | Seabirds        | <i>Eudyptes schlegeli</i>         | 1999      | 1981.5      | 36       | 1                    | 1          | -1.08  |
| Cullen et al. (2009)          | AR5      | AUSW       | -38.50   | 145.16    | Seabirds        | <i>Eudyptula minor</i>            | 2002.5    | 1985.25     | 40       | 1                    | 0          | 3.45   |
| Atkinson et al. (2015)        | Current  | NECS       | 50.25    | -4.21     | Holozooplankton | <i>Euterpina acutifrons</i>       | 2012      | 2000        | 25       | 24                   | 0          | 5.2    |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Eutriptiella</i> spp.          | 2016      | 2004.5      | 24       | 29                   | 1          | -0.28  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Evadne</i> spp.                | 2003      | 1988.25     | 45       | 49                   | 1          | -6.41  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Meroplankton    | Fish eggs                         | 2002.666  | 1988        | 45       | 49                   | 1          | -5.84  |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                          | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|----------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Greve et al. (1996)              | AR5      | NECS       | 54.19    | 7.90      | Meroplankton    | Fish larvae                      | 1998.8    | 1986.4      | 19       | 25                   | 1          | -17.68 |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Fragilaria</i> spp.           | 2009      | 1983.5      | 45       | 49                   | 1          | -4.09  |
| Wanless et al. (2009)            | AR5      | NECS       | 56.18    | -2.56     | Seabirds        | <i>Fratercula arctica</i>        | 2006      | 1990.5      | 36       | 7                    | 0          | 1.11   |
| Descamps et al. (2019)           | Current  | SARC       | 69.07    | 15.17     | Seabirds        | <i>Fratercula arctica</i>        | 2016      | 1999        | 35       | 16                   | 1          | -0.32  |
| Bertram et al. (2001)            | AR5      | ALSK       | 50.87    | -129.08   | Seabirds        | <i>Fratercula cirrhata</i>       | 2005.333  | 1992.333    | 25       | 4                    | 1          | -7.24  |
| Descamps et al. (2019)           | Current  | ALSK       | 54.18    | -164.83   | Seabirds        | <i>Fratercula corniculata</i>    | 2015      | 2001.5      | 28       | 16                   | 0          | 0.26   |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | <i>Fritillaria borealis</i>      | 1994      | 1984        | 21       | 25                   | 0          | NA     |
| Wanless et al. (2009)            | AR5      | NECS       | 56.18    | -2.56     | Seabirds        | <i>Fulmarus glacialis</i>        | 2006      | 1988.5      | 36       | 7                    | 1          | 0.16   |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Fulmarus glacialisoides</i>   | 2004      | 1977        | 55       | 9                    | 0          | 3.87   |
| Morgan et al. (2013)             | Current  | GFST       | 43.00    | -51.00    | Fish            | <i>Gadus morhua</i>              | 2009      | 1992        | 35       | 2                    | 0          | 6.35   |
| McQueen and Marshall (2017)      | Current  | NECS       | 53.50    | -5.00     | Fish            | <i>Gadus morhua</i>              | 2013.666  | 1994.833    | 32       | 2                    | 1          | -10.57 |
| Morgan et al. (2013)             | Current  | NWCS       | 48.00    | -51.00    | Fish            | <i>Gadus morhua</i>              | 2012.333  | 1990.5      | 38       | 2                    | 0          | 10.66  |
| McQueen and Marshall (2017)      | Current  | SARC       | 60.00    | 1.00      | Fish            | <i>Gadus morhua</i>              | 2014      | 1999.5      | 30       | 2                    | 1          | -6.58  |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Meroplankton    | <i>Galathea</i> spp.             | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | Gammarid amphipods               | 2007      | 1990        | 45       | 49                   | 1          | -0.97  |
| Chevillot et al. (2017)          | Current  | NECS       | 45.42    | -0.86     | Fish            | <i>Gasterosteus aculeatus</i>    | 2011.5    | 1992        | 26       | 7                    | 1          | NA     |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Meroplankton    | Gastropod larvae                 | 2003      | 1992        | 21       | 25                   | 1          | 0.8    |
| Scharfe and Wiltshire (2019)     | Current  | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Guindardia delicatula</i>     | 2015      | 1988.5      | 54       | 8                    | 1          | -11.8  |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | <i>Gyrodinium</i> spp.           | 2012      | 2000        | 25       | 24                   | 1          | -15.6  |
| Chivers et al. (2020)            | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Gyrosigma</i> spp.            | 2016      | 1987        | 59       | 12                   | 1          | 0.4    |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Harpacticoida</i>             | 2002      | 1980        | 45       | 49                   | 1          | -4.98  |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Helicolenus dactylopterus</i> | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Hemiselmis</i> spp.           | 2016      | 2004.5      | 24       | 29                   | 0          | 7.57   |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Hemitripterus americanus</i>  | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Heterocapsa rotunda</i>       | 2016      | 2004.5      | 24       | 29                   | 1          | -15.74 |

| Reference                     | Database | Long-hurst | Latitude | Longitude | Group           | Species                             | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|-------------------------------|----------|------------|----------|-----------|-----------------|-------------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Heterocapsa triquetra</i>        | 2016      | 2004.5      | 24       | 29                   | 0          | 10.65  |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Hippoglossina oblonga</i>        | 2015      | 1989        | 48       | 35                   | 1          | -3.14  |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Hippoglossoides platessoides</i> | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Hippoglossus hippoglossus</i>    | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Homarus americanus</i>           | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Émond et al. (2020)           | Current  | NWCS       | 47.50    | -63.00    | Meroplankton    | <i>Hyas</i> spp.                    | 2012      | 1997        | 31       | 2                    | 1          | NA     |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Hyperoplus lanceolatus</i>       | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Icichthys lockingtoni</i>        | 2008      | 1979.5      | 58       | 43                   | 1          | -5.6   |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Idiacanthus antrostomus</i>      | 2008      | 1979.5      | 58       | 43                   | 1          | -3     |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Illex illecebrosus</i>           | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Holozooplankton | <i>Katablepharis</i> spp.           | 2016      | 2004.5      | 24       | 29                   | 1          | -7.28  |
| Atkinson et al. (2015)        | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | <i>Katodinium</i> spp.              | 2012      | 2000        | 25       | 24                   | 1          | -3.2   |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Labidocera wollastoni</i>        | 2002      | 1980        | 45       | 49                   | 1          | -2.52  |
| Thaxton et al. (2020)         | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Lagodon rhomboides</i>           | 2013      | 1999.5      | 28       | 9                    | 1          | -1.5   |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Meroplankton    | Lamellibranch larvae                | 2002.666  | 1988        | 45       | 49                   | 1          | 4.22   |
| Greve et al. (1996)           | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | <i>Lanice conchilega</i>            | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Descamps et al. (2019)        | Current  | ALSK       | 54.18    | -164.83   | Seabirds        | <i>Larus glaucescens</i>            | 2015      | 2004.5      | 22       | 16                   | 1          | -0.08  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | Larvacea                            | 2002      | 1980        | 45       | 49                   | 1          | -6.68  |
| Thaxton et al. (2020)         | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Leiostomus xanthurus</i>         | 2013      | 1999.5      | 28       | 9                    | 1          | -2.1   |
| Scharfe and Wiltshire (2019)  | Current  | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Leptocylindricus minimus</i>     | 2015      | 1988.5      | 54       | 8                    | 1          | -28.31 |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Lestidiops ringens</i>           | 2008      | 1979.5      | 58       | 43                   | 1          | -7.3   |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Leucoraja erinacea</i>           | 2014      | 1990.5      | 48       | 35                   | 0          | NA     |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Leucoraja ocellata</i>           | 2014      | 1990.5      | 48       | 35                   | 0          | NA     |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Leuroglossus stibius</i>         | 2008      | 1979.5      | 58       | 43                   | 0          | 1.2    |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Limacina retroversa</i>          | 2002      | 1980        | 45       | 49                   | 0          | 8.37   |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Limanda ferruginea</i>           | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                                | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|--|-----------|-------------|----------|----------------------|------------|--------|
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Limanda limanda</i>                 | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Meroplankton    | <i>Liocarcinus</i> spp.                | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Liparis liparis</i>                 | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Lipagus ochotensis</i>              | 2008      | 1979.5      | 58       | 43                   | 1          | -3.2   |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Liza aurata</i>                     | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Loligo pealeii</i>                  | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Lophius gastrophysus</i>            | 2014      | 1990.5      | 48       | 35                   | 0          | NA     |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Lyopsetta exilis</i>                | 2008      | 1979.5      | 58       | 43                   | 1          | -0.7   |
| Beukema et al. (2009)            | AR5      | NECS       | 53.00    | 6.00      | Invertebrates   | <i>Macoma balthica rubra</i>           | 2007      | 1988        | 39       | 1                    | 1          | -5.38  |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Macronectes giganteus</i>           | 2004      | 1982        | 45       | 9                    | 0          | 1.91   |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | <i>Magelona</i> spp.                   | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Holozooplankton | <i>Medusae</i>                         | 2012      | 2000        | 25       | 24                   | 0          | 2.4    |
| Ramp et al. (2015)               | Current  | NWCS       | 49.90    | -64.50    | Mammals         | <i>Megaptera novaeangliae</i>          | 2010      | 1998.5      | 24       | 2                    | 1          | -11.67 |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Melamphaes lugubris</i>             | 2008      | 1979.5      | 58       | 43                   | 0          | 3      |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Melanogrammus aeglefinus</i>        | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Merluccius bilinearis</i>           | 2014      | 1990.5      | 48       | 35                   | 0          | NA     |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Merluccius productus</i>            | 2008      | 1979.5      | 58       | 43                   | 1          | -3     |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | <i>Mesodinium rubrum</i>               | 2014      | 2002.25     | 25       | 24                   | 0          | -14.28 |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Holozooplankton | <i>Metridia lucens</i>                 | 2012      | 2000        | 25       | 24                   | 1          | -8.4   |
| Thaxton et al. (2020)            | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Micropogonias undulatus</i>         | 2013      | 1999.5      | 28       | 9                    | 1          | -14.4  |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Holozooplankton | <i>Microsetella</i> spp.               | 2012      | 2000        | 25       | 24                   | 0          | 11.2   |
| Costello et al. (2006)           | AR5      | NWCS       | 41.50    | -71.35    | Holozooplankton | <i>Mnemiopsis leidyi</i>               | 2003      | 1977        | 53       | 2                    | 1          | -11.13 |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Monoraphidium contortum</i>         | 2016      | 2004.5      | 24       | 29                   | 1          | -13.92 |
| Peer and Miller (2014)           | Current  | NWCS       | 37.52    | -76.10    | Fish            | <i>Morone saxatilis</i>                | 2012      | 1995.5      | 20       | 1                    | 1          | -3     |
| Wanless et al. (2009)            | AR5      | NECS       | 55.00    | -5.00     | Seabirds        | <i>Morus bassanus</i>                  | 2007      | 1993.5      | 28       | 1                    | 0          | 2.1    |
| Thaxton et al. (2020)            | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Mugil cephalus</i>                  | 2013      | 1999.5      | 28       | 9                    | 1          | -9.1   |
| Langan et al. (2021)             | Current  | NWCS       | 41.50    | -71.40    | Fish            | <i>Myoxocephalus octodecemspinosus</i> | 2016      | 1987.5      | 58       | 5                    | 0          | 5.03   |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                               | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|---------------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Myoxocephalus scorpius</i>         | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |
| Thaxton et al. (2020)            | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Myrophis punctatus</i>             | 2013      | 1999.5      | 28       | 9                    | 1          | -14.5  |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Myxocephalus octodecemspinosus</i> | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Nannobrachium regale</i>           | 2008      | 1979.5      | 58       | 43                   | 0          | 0.7    |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Nannobrachium ritteri</i>          | 2008      | 1979.5      | 58       | 43                   | 1          | -4.8   |
| Chivers et al. (2020)            | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Navicula</i> spp.                  | 2016      | 1987        | 59       | 12                   | 0          | -1.57  |
| Bertram et al. (2001)            | AR5      | BERS       | 50.00    | 145.00    | Holozooplankton | <i>Neocalanus plumchrus</i>           | 1996      | 1985.5      | 22       | 4                    | 1          | -22    |
| Bertram et al. (2001)            | AR5      | CCAL       | 48.65    | -126.67   | Holozooplankton | <i>Neocalanus plumchrus</i>           | 1998      | 1986.5      | 24       | 4                    | 1          | -13    |
| Mackas et al. (1998)             | AR5      | PSAE       | 50.00    | -145.00   | Holozooplankton | <i>Neocalanus plumchrus</i>           | 1996      | 1982        | 29       | 1                    | 1          | -10.34 |
| Chevillot et al. (2017)          | Current  | NECS       | 45.42    | -0.86     | Holozooplankton | <i>Neomysis integer</i>               | 2010      | 1997.5      | 26       | 7                    | 1          | NA     |
| McGeady et al. (2021)            | Current  | NECS       | 53.75    | -4.75     | Meroplankton    | <i>Nephrops norvegicus</i>            | 2010      | 1996        | 29       | 1                    | 1          | -11.94 |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Phytoplankton   | <i>Noctiluca scintillans</i>          | 2003.75   | 1989.25     | 21       | 25                   | 1          | 4.03   |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Nodularia</i> spp.                 | 2016      | 2004.5      | 24       | 29                   | 0          | 21.25  |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | <i>Obelia</i> spp.                    | 1994      | 1984        | 21       | 25                   | 0          | NA     |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Oceanites oceanicus</i>            | 2004      | 1981.5      | 46       | 9                    | 0          | 1.01   |
| Descamps et al. (2019)           | Current  | ALSK       | 54.18    | -164.83   | Seabirds        | <i>Oceanodroma furcata</i>            | 2015      | 1998.5      | 34       | 16                   | 1          | -0.74  |
| Descamps et al. (2019)           | Current  | ALSK       | 54.18    | -164.83   | Seabirds        | <i>Oceanodroma leucorhoa</i>          | 2015      | 2005        | 21       | 16                   | 1          | -0.53  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Odontella aurita</i>               | 2008      | 1983.666    | 45       | 49                   | 1          | -4.69  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Odontella sinensis</i>             | 2011      | 1985.166    | 45       | 49                   | 1          | 3.41   |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | <i>Oikopleura dioica</i>              | 1994      | 1984.25     | 21       | 25                   | 1          | -2.83  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Oithona</i> spp.                   | 2007      | 1990        | 45       | 49                   | 1          | -1.65  |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Holozooplankton | <i>Oncaeae</i> spp.                   | 2012      | 2000        | 25       | 24                   | 0          | 0.4    |
| Kovach et al. (2013)             | Current  | ALSK       | 58.38    | -134.65   | Fish            | <i>Oncorhynchus clarkii</i>           | 2010      | 1995        | 31       | 2                    | 1          | -1.15  |
| Taylor (2008)                    | Current  | ALSK       | 58.38    | -134.65   | Fish            | <i>Oncorhynchus gorbuscha</i>         | 2007.5    | 1992        | 34       | 1                    | 1          | -3.21  |
| Kovach et al. (2015)             | Current  | ALSK       | 58.38    | -134.64   | Fish            | <i>Oncorhynchus keta</i>              | 2010      | 1995.5      | 30       | 4                    | 1          | -3.93  |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                          | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|----------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Rubenstein et al. (2019)         | Current  | CCAL       | 49.65    | -125.44   | Fish            | <i>Oncorhynchus keta</i>         | 2016.333  | 2003.333    | 34       | 2                    | 1          | -4.34  |
| Kovach et al. (2015)             | Current  | ALSK       | 58.38    | -134.64   | Fish            | <i>Oncorhynchus kisutch</i>      | 2010      | 1995.5      | 30       | 4                    | 1          | -4.64  |
| Rubenstein et al. (2019)         | Current  | CCAL       | 49.65    | -125.44   | Fish            | <i>Oncorhynchus kisutch</i>      | 2015      | 1998.5      | 34       | 2                    | 1          | -8     |
| Robards and Quinn (2002)         | Current  | CCAL       | 47.44    | -120.84   | Fish            | <i>Oncorhynchus mykiss</i>       | 1998      | 1977.125    | 49       | 1                    | 0          | -0.19  |
| Kovach et al. (2015)             | Current  | ALSK       | 58.38    | -134.64   | Fish            | <i>Oncorhynchus nerka</i>        | 2010      | 1995.5      | 30       | 4                    | 0          | 1.61   |
| Crozier et al. (2011)            | Current  | CCAL       | 46.24    | -124.00   | Fish            | <i>Oncorhynchus nerka</i>        | 2000.333  | 1976        | 75       | 1                    | 1          | -4.22  |
| Kovach et al. (2015)             | Current  | ALSK       | 58.38    | -134.64   | Fish            | <i>Oncorhynchus tshawytscha</i>  | 2010      | 1995.5      | 30       | 4                    | 0          | 0.81   |
| Kovach et al. (2012)             | Current  | ALSK       | 58.38    | -134.65   | Fish            | <i>Oncorhyncus gorbuscha</i>     | 2011      | 1997        | 29       | 1                    | 1          | NA     |
| Morita (2018)                    | Current  | KURO       | 43.27    | 141.37    | Fish            | <i>Oncorhyncus gorbuscha</i>     | 2017      | 2005        | 25       | 1                    | 1          | -2.92  |
| Chambers et al. (2014)           | Current  | AUSW       | -32.30   | 115.69    | Seabirds        | <i>Onychoprion anaethetus</i>    | 2011      | 1998.5      | 26       | 4                    | 0          | NA     |
| Chambers et al. (2014)           | Current  | AUSW       | -28.77   | 113.85    | Seabirds        | <i>Onychoprion fuscata</i>       | 2010      | 2000.5      | 20       | 4                    | 0          | NA     |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Oocystis</i> spp.             | 2016      | 2004.5      | 24       | 29                   | 1          | -54.08 |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | <i>Ophiura</i> spp.              | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Ahas and Aasa (2006)             | AR5      | NECS       | 58.20    | 24.30     | Fish            | <i>Osmerus eperlanus</i>         | 2003      | 1978.5      | 48       | 1                    | 1          | -0.08  |
| Atkinson et al. (2015)           | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | <i>Other ciliates</i>            | 2012      | 2000        | 25       | 24                   | 1          | -28.8  |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Oxyjulis californica</i>      | 2008      | 1979.5      | 58       | 43                   | 1          | -6.9   |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Pagodroma nivea</i>           | 2004      | 1987        | 35       | 9                    | 0          | -0.23  |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Meroplankton    | <i>Pagurus</i> spp.              | 1994      | 1984        | 21       | 25                   | 0          | NA     |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Para-pseudocalanus</i> spp.   | 1998      | 1982        | 45       | 49                   | 1          | -2.71  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Paralia sulcata</i>           | 2009      | 1983.5      | 45       | 49                   | 1          | -1.83  |
| Thaxton et al. (2020)            | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Paralichthys alboguttata</i>  | 2013      | 1999.5      | 28       | 9                    | 1          | -13.9  |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Paralichthys californicus</i> | 2008      | 1979.5      | 58       | 43                   | 0          | 5.3    |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Paralichthys dentatus</i>     | 2014.333  | 1992.5      | 48       | 35                   | 1          | -5.17  |
| Thaxton et al. (2020)            | Current  | NWCS       | 34.60    | -77.20    | Meroplankton    | <i>Paralichthys lethostigma</i>  | 2013      | 1999.5      | 28       | 9                    | 1          | -2.8   |
| Greve et al. (1996)              | Current  | NECS       | 54.19    | 7.90      | Holozooplankton | <i>Paramysis</i> spp.            | 1994      | 1984        | 21       | 25                   | 0          | NA     |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Parophrys vetulus</i>         | 2008      | 1979.5      | 58       | 43                   | 1          | -1.8   |

| Reference                     | Database | Long-hurst | Latitude | Longitude | Group           | Species                          | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|-------------------------------|----------|------------|----------|-----------|-----------------|----------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Moore et al. (2011)           | AR5      | NECS       | 50.58    | -4.32     | Invertebrates   | <i>Patella depressa</i>          | 2007      | 1976.5      | 62       | 2                    | 1          | -10.2  |
| Moore et al. (2011)           | AR5      | NECS       | 50.58    | -4.32     | Invertebrates   | <i>Patella vulgata</i>           | 2007      | 1976.5      | 62       | 2                    | 0          | 3.3    |
| Langan et al. (2021)          | Current  | NWCS       | 41.50    | -71.40    | Fish            | <i>Peprius triacanthus</i>       | 2016      | 1987.5      | 58       | 5                    | 1          | -6.36  |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Peridiniella catenata</i>     | 2016      | 2004.5      | 24       | 29                   | 1          | -38.61 |
| Atkinson et al. (2015)        | Current  | NECS       | 50.25    | -4.21     | Phytoplankton   | <i>Phaeocystis</i> spp.          | 2012      | 2000        | 25       | 24                   | 1          | -6     |
| Frederiksen et al. (2004)     | AR5      | NECS       | 56.18    | -2.55     | Seabirds        | <i>Phalacrocorax aristotelis</i> | 2005      | 1989.25     | 34       | 3                    | 1          | -3.72  |
| Descamps et al. (2019)        | Current  | BERS       | 57.13    | -170.28   | Seabirds        | <i>Phalacrocorax urile</i>       | 2015      | 2000        | 31       | 16                   | 1          | -0.16  |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Pholis gunnellus</i>          | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |
| Burthe et al. (2012)          | Current  | NECS       | 56.50    | -1.50     | Phytoplankton   | Phytoplankton Colour Index       | 2006      | 1994.5      | 24       | 4                    | 1          | -5.76  |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Plagioselmis prolonga</i>     | 2016      | 2004.5      | 24       | 29                   | 0          | 5.1    |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Planctonema lauterbornii</i>  | 2016      | 2004.5      | 24       | 29                   | 0          | 3.14   |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Planktolyngba</i> spp.        | 2016      | 2004.5      | 24       | 29                   | 1          | -26.63 |
| Schlüter et al. (2010)        | AR5      | NECS       | 54.18    | 7.90      | Holozooplankton | <i>Pleurobrachia pileus</i>      | 2000.666  | 1987.833    | 30       | 2                    | 1          | -10.47 |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Pleuronectes platessa</i>     | 2008.5    | 1986.75     | 54       | 30                   | 0          | -10    |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Pleuronichthys verticalis</i> | 2008      | 1979.5      | 58       | 43                   | 0          | 5.6    |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Podon</i> spp.                | 2002.666  | 1988        | 45       | 49                   | 1          | -2.52  |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Pollachius pollachius</i>     | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Pollachius virens</i>         | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Pollachius virens</i>         | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Meroplankton    | Polychaete larvae                | 2007      | 1990        | 45       | 49                   | 1          | 1.52   |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Pomatoschistus minutus</i>    | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Diaz-Almela et al. (2007)     | AR5      | MEDI       | 41.00    | 7.00      | Plants          | <i>Posidonia oceanica</i>        | 2004      | 1988.5      | 32       | 1                    | 1          | NA     |
| Langan et al. (2021)          | Current  | NWCS       | 41.50    | -71.40    | Fish            | <i>Prionotus evolans</i>         | 2016      | 1987.5      | 58       | 5                    | 1          | -7.74  |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Proboscia alata</i>           | 2009      | 1983.5      | 45       | 49                   | 1          | -6.78  |
| Chivers et al. (2020)         | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Proboscia inermis</i>         | 2016      | 1987        | 59       | 12                   | 0          | 1.68   |
| Scharfe and Wiltshire (2019)  | Current  | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Prorocentrum micans</i>       | 2015      | 1988.5      | 54       | 8                    | 1          | -8.65  |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                               | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|---------------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Prorocentrum</i> spp.              | 2009      | 1983.5      | 45       | 49                   | 1          | -5.72  |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Protomyctophum crockeri</i>        | 2008      | 1979.5      | 58       | 43                   | 1          | -1.4   |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Holozooplankton | <i>Protoperidinium bipes</i>          | 2016      | 2004.5      | 24       | 29                   | 1          | -21.35 |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Holozooplankton | <i>Protoperidinium brevipes</i>       | 2016      | 2004.5      | 24       | 29                   | 1          | -2.76  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Protoperidinium</i> spp.           | 2010      | 1989        | 45       | 49                   | 1          | -3.85  |
| Chivers et al. (2020)            | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Pseudo-nitzschia delicatissima</i> | 2016      | 1987        | 59       | 12                   | 1          | -3.35  |
| Chivers et al. (2020)            | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Pseudo-nitzschia seriata</i>       | 2016      | 1987        | 59       | 12                   | 1          | -1.46  |
| Scharfe and Wiltshire (2019)     | Current  | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Pseudo-nitzschia</i> spp.          | 2015      | 1988.5      | 54       | 8                    | 1          | -28.31 |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Pseudocalanus elongatus</i>        | 2007      | 1990        | 45       | 49                   | 1          | -4.09  |
| Persson et al. (2012)            | Current  | SARC       | 66.33    | 33.66     | Holozooplankton | <i>Pseudocalanus minutus</i>          | 2010      | 1987        | 47       | 2                    | 1          | NA     |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Pseudopedinella</i> spp.           | 2016      | 2004.5      | 24       | 29                   | 1          | -26.66 |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Pseudopleuronectes americanus</i>  | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Schroeder et al. (2009)          | AR5      | CCAL       | 37.70    | -123.00   | Seabirds        | <i>Ptychoramphus aleuticus</i>        | 2006      | 1991.666    | 19       | 2                    | 0          | 4      |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Pygocelis adeliae</i>              | 2004      | 1977        | 55       | 9                    | 0          | 0.51   |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Pyramimonas</i> spp.               | 2016      | 2004.5      | 24       | 29                   | 1          | -15.81 |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Rhizosolenia hebetata</i>          | 2009      | 1983.5      | 45       | 49                   | 1          | -0.89  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Rhizosolenia imbricata</i>         | 2011      | 1985.166    | 45       | 49                   | 1          | -8.59  |
| Scharfe and Wiltshire (2019)     | Current  | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Rhizosolenia setigera</i>          | 2015      | 1988.5      | 54       | 8                    | 1          | -10.22 |
| Chivers et al. (2020)            | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Rhizosolenia styliformis</i>       | 2016      | 1987        | 59       | 12                   | 1          | -2.31  |
| Volkov and Pozdnyakov (2021)     | Current  | BPLR       | 73.00    | 125.00    | Seabirds        | <i>Rhodostethia rosea</i>             | 2013      | 1997.5      | 32       | 1                    | 1          | NA     |
| Byrd et al. (2008)               | AR5      | BERS       | 57.00    | -169.00   | Seabirds        | <i>Rissa brevirostris</i>             | 2005      | 1990        | 31       | 4                    | 1          | -8.35  |
| Descamps et al. (2019)           | Current  | PSAW       | 52.35    | -176.93   | Seabirds        | <i>Rissa brevirostris</i>             | 2015      | 2000        | 31       | 16                   | 1          | -0.44  |
| Descamps et al. (2019)           | Current  | ALSK       | 58.92    | -152.17   | Seabirds        | <i>Rissa tridactyla</i>               | 2016      | 1999        | 35       | 16                   | 0          | 0.04   |
| Byrd et al. (2008)               | AR5      | BERS       | 57.00    | -169.00   | Seabirds        | <i>Rissa tridactyla</i>               | 2005      | 1990        | 31       | 4                    | 1          | -6.1   |

| Reference                     | Database | Long-hurst | Latitude | Longitude | Group          | Species                     | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|-------------------------------|----------|------------|----------|-----------|----------------|-----------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Moe et al. (2009)             | AR5      | BPLR       | 78.90    | 12.22     | Seabirds       | <i>Rissa tridactyla</i>     | 2008      | 1989        | 39       | 2                    | 0          | 1.3    |
| Frederiksen et al. (2004)     | AR5      | NECS       | 56.18    | -2.55     | Seabirds       | <i>Rissa tridactyla</i>     | 2005      | 1990.75     | 22       | 3                    | 0          | 4.81   |
| Greve et al. (1996)           | Current  | NECS       | 54.19    | 7.90      | Holzooplankton | <i>Sagitta spp.</i>         | 1994      | 1984        | 21       | 25                   | 1          | NA     |
| Otero et al. (2014)           | Current  | ARCT       | 65.75    | -14.90    | Fish           | <i>Salmo salar</i>          | 2008      | 1998.5      | 20       | 3                    | 1          | NA     |
| Otero et al. (2014)           | Current  | BPLR       | 53.56    | -56.35    | Fish           | <i>Salmo salar</i>          | 2008      | 1989        | 39       | 3                    | 1          | NA     |
| Kennedy and Crozier (2010)    | AR5      | NECS       | 55.20    | -6.53     | Fish           | <i>Salmo salar</i>          | 2007.611  | 1992.416    | 31       | 1                    | 1          | 3.31   |
| Juanes et al. (2004)          | AR5      | NWCS       | 41.25    | -72.35    | Fish           | <i>Salmo salar</i>          | 2000.162  | 1986.135    | 23       | 1                    | 1          | -7.9   |
| Otero et al. (2014)           | Current  | SARC       | 70.03    | 22.96     | Fish           | <i>Salmo salar</i>          | 2009      | 1998.5      | 23       | 3                    | 1          | NA     |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish           | <i>Salmo trutta</i>         | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Kovach et al. (2013)          | Current  | ALSK       | 58.38    | -134.65   | Fish           | <i>Salvelinus malma</i>     | 2010      | 1995        | 31       | 2                    | 1          | -0.68  |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish           | <i>Sardina pilchardus</i>   | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton   | <i>Sardinops sagax</i>      | 2008      | 1979.5      | 58       | 43                   | 1          | -11.1  |
| Fitchett et al. (2019)        | Current  | EAFR       | -29.85   | 31.02     | Fish           | <i>Sardinops sagax</i>      | 2012      | 1979.5      | 66       | 1                    | 0          | 1.3    |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton   | <i>Scomber japonicus</i>    | 2008      | 1979.5      | 58       | 43                   | 1          | -1     |
| Kanamori et al. (2019)        | Current  | KURO       | 35.00    | 141.00    | Fish           | <i>Scomber japonicus</i>    | 2017      | 1997.5      | 40       | 1                    | 0          | 4.24   |
| Jansen and Gislason (2011)    | Current  | NECS       | 56.95    | 11.30     | Fish           | <i>Scomber scombrus</i>     | 2010.5    | 1992        | 22       | 1                    | 1          | -23.4  |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish           | <i>Scomber scombrus</i>     | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish           | <i>Scophthalmus maximus</i> | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| van Walraven et al. (2017)    | Current  | NECS       | 53.57    | 6.94      | Fish           | <i>Scophthalmus rhombus</i> | 2013      | 1986.5      | 54       | 30                   | 0          | NA     |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish           | <i>Scophthalmus aquosus</i> | 2014      | 1990.5      | 48       | 35                   | 0          | NA     |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton   | <i>Sebastes aurora</i>      | 2008      | 1979.5      | 58       | 43                   | 1          | -5.4   |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton   | <i>Sebastes diploproa</i>   | 2008      | 1979.5      | 58       | 43                   | 1          | -12.3  |
| Henderson et al. (2017)       | Current  | NWCS       | 40.00    | -71.00    | Fish           | <i>Sebastes fasciatus</i>   | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton   | <i>Sebastes goodei</i>      | 2008      | 1979.5      | 58       | 43                   | 0          | 6.7    |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton   | <i>Sebastes jordani</i>     | 2008      | 1979.5      | 58       | 43                   | 1          | -2.8   |
| Asch (2015)                   | Current  | CCAL       | 34.00    | -122.00   | Meroplankton   | <i>Sebastes paucispinis</i> | 2008      | 1979.5      | 58       | 43                   | 0          | 1.6    |
| Atkinson et al. (2015)        | Current  | NECS       | 50.25    | -4.21     | Holzooplankton | Siphonophores               | 2012      | 2000        | 25       | 24                   | 1          | -2.4   |
| Edwards and Richardson (2004) | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton  | <i>Skeletonema costatum</i> | 2012      | 1986        | 45       | 49                   | 0          | -2.19  |
| Forsblom et al. (2019)        | Current  | NECS       | 59.70    | 24.50     | Phytoplankton  | <i>Skeletonema marinoi</i>  | 2016      | 2004.5      | 24       | 29                   | 1          | -52.51 |

| Reference                       | Data-base | Long-hurst | Latitude | Lon-gitude | Group           | Species                             | Last year | Median year | Dura-tion | Number of repli-cates | Con-sistent | Shift  |
|---------------------------------|-----------|------------|----------|------------|-----------------|-------------------------------------|-----------|-------------|-----------|-----------------------|-------------|--------|
| Atkinson et al. (2015)          | Current   | NECS       | 50.25    | -4.21      | Phytoplankton   | Small <i>Peridinium</i> spp.        | 2012      | 2000        | 25        | 24                    | 1           | -12.8  |
| Forsblom et al. (2019)          | Current   | NECS       | 59.70    | 24.50      | Phytoplankton   | <i>Snowella</i> spp.                | 2016      | 2004.5      | 24        | 29                    | 0           | 6.32   |
| Fincham et al. (2013)           | Current   | NECS       | 51.00    | -4.75      | Fish            | <i>Solea solea</i>                  | 2008.875  | 1992.125    | 27        | 1                     | 1           | -7.45  |
| Wanless et al. (2009)           | AR5       | NECS       | 56.18    | -2.56      | Seabirds        | <i>Somateria mollissima</i>         | 2006      | 1988.5      | 36        | 7                     | 1           | 0.19   |
| D'Alba et al. (2010)            | AR5       | SARC       | 64.02    | -22.71     | Seabirds        | <i>Somateria mollissima</i>         | 2006      | 1991.5      | 30        | 1                     | 1           | -2.8   |
| Chambers et al. (2014)          | Current   | SANT       | -44.04   | 65.22      | Seabirds        | <i>Spheniscus magellanicus</i>      | 2006      | 1994.5      | 24        | 4                     | 0           | NA     |
| Greve et al. (1996)             | Current   | NECS       | 54.19    | 7.90       | Holozooplankton | <i>Spioniden metatrocha</i>         | 1994      | 1984        | 21        | 25                    | 0           | NA     |
| van Walraven et al. (2017)      | Current   | NECS       | 53.57    | 6.94       | Fish            | <i>Sprattus sprattus</i>            | 2013      | 1986.5      | 54        | 30                    | 0           | NA     |
| Henderson et al. (2017)         | Current   | NWCS       | 40.00    | -71.00     | Fish            | <i>Squalus acanthias</i>            | 2014      | 1990.5      | 48        | 35                    | 0           | NA     |
| Asch (2015)                     | Current   | CCAL       | 34.00    | -122.00    | Meroplankton    | <i>Stenobrachius leucopsarus</i>    | 2008      | 1979.5      | 58        | 43                    | 1           | -1.7   |
| Langan et al. (2021)            | Current   | NWCS       | 41.50    | -71.40     | Fish            | <i>Stenotomus chrysops</i>          | 2016      | 1987.5      | 58        | 5                     | 1           | -2.1   |
| Wanless et al. (2009)           | AR5       | NECS       | 56.18    | -2.56      | Seabirds        | <i>Sterna hirundo</i>               | 2006      | 1988.5      | 36        | 7                     | 1           | -0.45  |
| Wanless et al. (2009)           | AR5       | NECS       | 56.18    | -2.56      | Seabirds        | <i>Sterna paradisaea</i>            | 2002      | 1976        | 36        | 7                     | 1           | -2.26  |
| Wanless et al. (2009)           | AR5       | NECS       | 56.18    | -2.56      | Seabirds        | <i>Sterna sandvicensis</i>          | 2006      | 1988.5      | 36        | 7                     | 1           | -0.18  |
| Asch (2015)                     | Current   | CCAL       | 34.00    | -122.00    | Meroplankton    | <i>Stomias atriventer</i>           | 2008      | 1979.5      | 58        | 43                    | 0           | 1.2    |
| Atkinson et al. (2015)          | Current   | NECS       | 50.25    | -4.21      | Phytoplankton   | <i>Strombidium</i> spp.             | 2012      | 2000        | 25        | 24                    | 1           | -3.2   |
| Atkinson et al. (2015)          | Current   | NECS       | 50.25    | -4.21      | Holozooplankton | <i>Subeucalanus</i> spp.            | 2012      | 2000        | 25        | 24                    | 0           | 2      |
| Asch (2015)                     | Current   | CCAL       | 34.00    | -122.00    | Meroplankton    | <i>Symbolophorus californiensis</i> | 2008      | 1979.5      | 58        | 43                    | 1           | -6.15  |
| van Walraven et al. (2017)      | Current   | NECS       | 53.57    | 6.94       | Fish            | <i>Syngnathus acus</i>              | 2013      | 1986.5      | 54        | 30                    | 1           | NA     |
| Descamps et al. (2019)          | Current   | ALSK       | 54.18    | -164.83    | Seabirds        | <i>Synthliboramphus antiquus</i>    | 2015      | 2006        | 19        | 16                    | 1           | -0.21  |
| Asch (2015)                     | Current   | CCAL       | 34.00    | -122.00    | Meroplankton    | <i>Tarletonbeania crenularis</i>    | 2008      | 1979.5      | 58        | 43                    | 1           | 1.65   |
| Forsblom et al. (2019)          | Current   | NECS       | 59.70    | 24.50      | Phytoplankton   | <i>Teleaulax</i> spp.               | 2016      | 2004.5      | 24        | 29                    | 1           | -45.2  |
| Forsblom et al. (2019)          | Current   | NECS       | 59.70    | 24.50      | Holozooplankton | <i>Telonema subtile</i>             | 2016      | 2004.5      | 24        | 29                    | 1           | -35.06 |
| Greve et al. (1996)             | Current   | NECS       | 54.19    | 7.90       | Holozooplankton | <i>Temora longicaudata</i>          | 1994      | 1984        | 21        | 25                    | 1           | NA     |
| Edwards and Richardson (2004)   | AR5       | NECS       | 55.00    | 5.00       | Holozooplankton | <i>Temora longicornis</i>           | 2006.666  | 1991.5      | 45        | 49                    | 0           | -3.17  |
| Appelqvist and Havenhand (2016) | Current   | NECS       | 58.25    | 11.35      | Invertebrates   | <i>Teredo navalis</i>               | 2006      | 1988.5      | 36        | 1                     | 1           | 3.92   |
| Asch (2015)                     | Current   | CCAL       | 34.00    | -122.00    | Meroplankton    | <i>Tetragonurus cuvieri</i>         | 2008      | 1979.5      | 58        | 43                    | 0           | 3.5    |

| Reference                        | Database | Long-hurst | Latitude | Longitude | Group           | Species                           | Last year | Median year | Duration | Number of replicates | Consistent | Shift  |
|----------------------------------|----------|------------|----------|-----------|-----------------|-----------------------------------|-----------|-------------|----------|----------------------|------------|--------|
| Wiltshire et al. (2010)          | AR5      | NECS       | 54.18    | 7.90      | Phytoplankton   | <i>Thalassionema nitzschoides</i> | 2006      | 1984        | 45       | 1                    | 1          | NA     |
| Chivers et al. (2020)            | Current  | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Thalassionema nitzscioides</i> | 2015.5    | 1987.75     | 59       | 12                   | 1          | 1.33   |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Thalassiosira baltica</i>      | 2016      | 2004.5      | 24       | 29                   | 1          | -21.41 |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Thalassiosira</i> spp.         | 2012      | 1986        | 45       | 49                   | 0          | -3.36  |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Phytoplankton   | <i>Thalassiothrix longissima</i>  | 2009      | 1983.5      | 45       | 49                   | 1          | -1.94  |
| Barbraud and Weimerskirch (2006) | AR5      | APLR       | -66.66   | 140.00    | Seabirds        | <i>Thalassica antarctica</i>      | 2004      | 1992        | 25       | 9                    | 0          | 12.64  |
| Dufour et al. (2010)             | AR5      | NADR       | 46.00    | -13.00    | Fish            | <i>Thunnus alalunga</i>           | 2009      | 1990.333    | 39       | 2                    | 1          | -2.77  |
| Dufour et al. (2010)             | AR5      | NADR       | 46.00    | -13.00    | Fish            | <i>Thunnus thynnus</i>            | 2005      | 1993.833    | 25       | 2                    | 1          | -10    |
| Edwards and Richardson (2004)    | AR5      | NECS       | 55.00    | 5.00      | Holozooplankton | <i>Tomopteris</i> spp.            | 2002      | 1980        | 45       | 49                   | 1          | -4.19  |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Trachurus symmetricus</i>      | 2008      | 1979.5      | 58       | 43                   | 1          | -6.7   |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Triphoturus mexicanus</i>      | 2008      | 1979.5      | 58       | 43                   | 1          | -6.2   |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Trisopterus luscus</i>         | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |
| van Walraven et al. (2017)       | Current  | NECS       | 53.57    | 6.94      | Fish            | <i>Trisopterus minutus</i>        | 2013      | 1986.5      | 54       | 30                   | 1          | NA     |
| Descamps et al. (2019)           | Current  | ALSK       | 54.18    | -164.83   | Seabirds        | <i>Uria aalge</i>                 | 2015      | 1998.5      | 34       | 16                   | 0          | 0.07   |
| Byrd et al. (2008)               | AR5      | BERS       | 57.00    | -169.00   | Seabirds        | <i>Uria aalge</i>                 | 2005      | 1990        | 31       | 4                    | 0          | 1.3    |
| Schroeder et al. (2009)          | AR5      | CCAL       | 37.70    | -123.00   | Seabirds        | <i>Uria aalge</i>                 | 2006.666  | 1992.333    | 19       | 2                    | 1          | -2.4   |
| Frederiksen et al. (2004)        | AR5      | NECS       | 56.18    | -2.55     | Seabirds        | <i>Uria aalge</i>                 | 2005.6    | 1990.8      | 21       | 3                    | 0          | 0.82   |
| Descamps et al. (2019)           | Current  | ALSK       | 54.18    | -164.83   | Seabirds        | <i>Uria lomvia</i>                | 2016      | 1999        | 35       | 16                   | 1          | -0.31  |
| Byrd et al. (2008)               | AR5      | BERS       | 57.00    | -169.00   | Seabirds        | <i>Uria lomvia</i>                | 2005      | 1990        | 31       | 4                    | 0          | 2.36   |
| Gaston et al. (2005)             | AR5      | BPLR       | 74.04    | -90.03    | Seabirds        | <i>Uria lomvia</i>                | 2003.5    | 1992.5      | 26       | 1                    | 1          | -2.5   |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Urophycis chuss</i>            | 2014.666  | 1989.5      | 48       | 35                   | 0          | 8.86   |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Urophycis regia</i>            | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Henderson et al. (2017)          | Current  | NWCS       | 40.00    | -71.00    | Fish            | <i>Urophycis tenuis</i>           | 2014      | 1990.5      | 48       | 35                   | 1          | NA     |
| Cherry et al. (2013)             | Current  | BPLR       | 59.71    | -85.25    | Mammals         | <i>Ursus maritimus</i>            | 2013.714  | 2001.285    | 19       | 1                    | 1          | -1.46  |
| Asch (2015)                      | Current  | CCAL       | 34.00    | -122.00   | Meroplankton    | <i>Vinciguerria lucetia</i>       | 2008      | 1979.5      | 58       | 43                   | 1          | -5.6   |
| Forsblom et al. (2019)           | Current  | NECS       | 59.70    | 24.50     | Phytoplankton   | <i>Woronichinia</i> spp.          | 2016      | 2004.5      | 24       | 29                   | 1          | -13.92 |

| Reference                  | Data-base | Long-hurst | Latitude | Lon-gitude | Group | Species                   | Last year | Median year | Dura-tion | Number of repli-cates | Con-sistent | Shift |
|----------------------------|-----------|------------|----------|------------|-------|---------------------------|-----------|-------------|-----------|-----------------------|-------------|-------|
| Henderson et al. (2017)    | Current   | NWCS       | 40.00    | -71.00     | Fish  | <i>Zoarces americanus</i> | 2015      | 1989        | 48        | 35                    | 1           | 9.78  |
| van Walraven et al. (2017) | Current   | NECS       | 53.57    | 6.94       | Fish  | <i>Zoarces viviparus</i>  | 2013      | 1986.5      | 54        | 30                    | 1           | NA    |

Notes:

NECS: North-east Atlantic shelves; PSAW: Western Pacific sub-Arctic gyres; SARC: Atlantic sub-Arctic; NWCS: North-west Atlantic shelves; ARCT: Atlantic Arctic; CCAL: California Current; APLR: Austral Polar; NADR: North Atlantic Drift; NASE: North-east Atlantic subtropical gyre; BRAZ: Brazilian current coast; MED: Mediterranean Sea; BPLR: Boreal polar; ALSK: Alaska coastal downwelling; FKLD: South-west Atlantic shelves; CARB: Caribbean; SANT: Sub-Antarctic water ring; AUSW: Western Australian and Indonesian coast; GFST: Gulf Stream; BERS: North Pacific Epicontinental Sea; KURO: Kuroshio current; PSAW: Western Pacific sub-arctic gyres; SANT: Sub-Arctic water ring; EAFR: East African coast

## SM3.4 Section 3.5

**Table SM3.2 |** Climate-sensitive human pathogens associated with an aquatic-system infection route<sup>a</sup>

| Pathogen                  | How climate might affect disease occurrence   | Strength of ev-idence for water-related infections                            | Type of infection route | Type of study linking climate and infection |
|---------------------------|---|---|-------------------------|---|
| <i>Acanthamoeba</i>       | <i>A. polyphaga</i> linked to contact-lens washing and hygiene is important. Infections linked to flooding (presumed contamination of potable water). Water contamination links.  | Freshwater/seawater – Strong  | Waterborne              | PMCC; LR                                    |
| <i>Adenovirus</i>         | Subgroups A–E cause upper respiratory infections, conjunctivitis, febrile illness, sore throat and swollen glands. The enteric subgroup F adenoviruses Ad40 and Ad41 cause gastroenteritis in children. Contamination of groundwater used as a drinking-water source and from faecal or respiratory contamination of untreated recreational waters. Swimming pool outbreaks.  | Freshwater/seawater – Moderate  | Waterborne              | POWE; OI                                    |
| <i>Astrovirus</i>         | Astroviruses cause diarrhoea in children under 5 years old. Viruses are excreted in faeces, and they will be present in sewage. Contact with contaminated recreational waters may be a risk factor. Outbreaks are often mixed. Outbreak linked to flood-water contamination of shellfish with several viruses.  | Freshwater/seawater – Weak  | Waterborne              | POWE  |
| <i>Campylobacter</i> spp. | The commonest bacterial cause of diarrhoea. Most infections are sporadic, but waterborne outbreaks are linked to camp sites, travelling abroad, hospitals and large communities. Infection is commonly derived from contaminated poultry, and water for the chicken flocks may be one source of contamination. <i>Campylobacter</i> spp. are spiral/curved organisms when isolated from patients and change to a more resistant coccoid stage when present in water. Most human infections are caused by <i>C. jejuni</i> , <i>C. coli</i> and <i>C. lari</i> . <i>Campylobacter fetus</i> subsp. <i>fetus</i> can cause human infections, with septicaemia and gall bladder infection being more common than with the other species. <i>C. upsaliensis</i> , <i>C. hyoilectinalis</i> subsp. <i>lawsonii</i> and <i>C. hyoilectinalis</i> subsp. <i>hyoilectinalis</i> are occasionally isolated from diarrhoeal patients. Infection through contaminated drinking water and contaminated water runoffs associated with heavy rainfall.  | Freshwater/seawater – Strong, outbreaks represent a small percentage of cases | Waterborne              | PORA; EACI                                  |
| <i>Cryptosporidium</i>    | Cause diarrhoea in young mammals and in humans but cannot grow in the environment. Large waterborne outbreaks have been reported throughout the world. Oocysts are excreted in faeces and sewage. Many species ( <i>C. hominis</i> , <i>C. parvum</i> , <i>C. meleagridis</i> , <i>C. cuniculus</i> , <i>C. ubiquitum</i> , <i>C. viatorum</i> , <i>C. canis</i> , <i>C. felis</i> , <i>C. suis</i> , <i>C. scrofarum</i> , <i>C. bovis</i> and <i>C. muris</i> , and genotypes of <i>C. parvum</i> ) can cause human disease. Rainfall can contribute to drinking-water contamination from both human and animal faeces.   | Freshwater/seawater – Strong  | Waterborne              | SA; SFA; OI; POWE; FTA; RSA                 |
| <i>Cyanobacteria</i> spp. | Grow as blooms or mats, mostly within freshwater bodies. There are a large variety of species, many producing potent toxins that can cause acute and chronic disease in mammals, including humans. The toxins include microcystins, nodularins, anatoxins, saxitoxins, apllysiatoxins, cylindrospermopsins, beta-methyl-amino-L-alanine (BMAA) and lipopolysaccharides. Algal blooms are more commonly found in eutrophic (eutrophic waters have a high concentration of nutrients) inland waters. Human health risks arise if the water is consumed untreated, if people bathe or participate in water-contact sports in waters with a scum or heavy bloom and if contaminated water is used in renal dialysis. There have been some notable outbreaks associated with cyanobacterial toxins with a high mortality rate in dialysis patients. There are also associations between exposure to cyanobacterial toxins and long-term health risks including cancer. The risks from BMAA linked to neurological disease are unclear. Climate influence on algal blooms. Human recreational and drinking-water exposures. | Freshwater/seawater – Strong for outbreaks linked to peritoneal dialysis      | Water toxicosis         | CEO   |

| Pathogen   | How climate might affect disease occurrence   | Strength of evidence for water-related infections | Type of infection route             | Type of study linking climate and infection |
|--|---|---|-------------------------------------|---|
| <i>Cyanobacteria—<i>Microcystis</i> spp.</i>       | <i>M. aeruginosa</i> is a common cyanobacteria found in eutrophic waters. It can cause hepatic failure and diarrhoea in humans and other animals. An association was found between drinking water from a reservoir contaminated with <i>M. aeruginosa</i> and raised liver enzymes in a population in New South Wales, Australia. Other toxic species include <i>M. viridis</i> and <i>M. botrys</i> .  | Freshwater – Strong                               | Water toxicosis                     | GDSE  |
| <i>Dinoflagellates and diatoms</i>                 | These are protozoan organisms that can produce a range of potent toxins. They occur predominantly in saltwater and, under the right conditions, can produce blooms that cause 'red tides' that can cause toxic effects in fish and other sea life. The toxins can accumulate within shellfish, causing paralytic shellfish poisoning (PSP), diarrhoeic shellfish poisoning (DSP), neurotoxic shellfish poisoning (NSP) and amnesic shellfish poisoning (ASP). Some of the toxins can also accumulate through passing up the food chain to produce carnivorous fish that are toxic (ciguatera toxin). Coastal blooms causing respiratory symptoms, ciguatera and shellfish poisoning. Blooms of dinoflagellates are linked to weather and nutrients. | Freshwater/seawater – Strong                      | Toxin contamination of marine foods | CEO   |
| <i>Dracunculus medinensis</i>                      | <i>Dracunculus medinensis</i> life cycle involves the water flea Cyclops. It is the cause of dracuniasis. Human infection results from the consumption of water contaminated with infected water fleas. The adult worm emerges on the foot or leg, and rhabditoid larvae are released into the water where they re-infect water fleas. There is a WHO-led worldwide programme to eradicate Guinea worm disease. Rainfall contamination of source waters. Infection is associated with water scarcity and the start of the rainy season.   | Freshwater – Strong                               | Waterborne                          | RSA   |
| <i>Enteroviruses</i>                               | Gross contamination of drinking water leading to enterovirus outbreaks.   | Freshwater/seawater – Moderate                    | Waterborne                          | OI; QMRA                                    |
| <i>Escherichia coli—<i>Shiga cytotoxigenic</i></i> | Infection through contaminated drinking water and contaminated water runoffs associated with heavy rains.   | Freshwater/seawater – Strong                      | Waterborne                          | CA; OI                                      |
| <i>Fasciola hepatica</i>                           | A liver fluke (helminth) that is common in herbivores that graze in wet pasture. The parasite requires a snail as an intermediate host, and man is occasionally infected through the consumption of aquatic plants, particularly watercress, contaminated with the metacercaria.  | Freshwater – Strong                               | Water based                         | RAI   |
| <i>Giardia</i> spp.                                | Grows attached to the small intestinal lining and causes malabsorption in people. The parasite can be isolated from the faeces of wild and domestic animals, and waterborne outbreaks are usually associated with recreational water use. The parasite cyst, which is found in faeces, is moderately resistant to chlorine. The modes of transmission remain unclear. Giardia can be transmitted through recreational and drinking water, although hygiene is also important.   | Freshwater/seawater – Strong                      | Waterborne                          | CSS   |
| <i>Hepatitis A</i>                                 | Hepatitis A virus causes hepatitis and can be acquired person to person, through contaminated water, shellfish and foods eaten raw or washed in contaminated water, and waterborne routes. Infection resulting from sewage contamination of source waters and shellfish. Some rainfall associations.  | Freshwater/seawater – Strong                      | Waterborne                          | PORA; SFA                                   |
| <i>Hepatitis E</i>                                 | Hepatitis E virus has a genome of single-stranded RNA. Epidemiological evidence suggests that the disease can be transmitted by drinking water contaminated with faeces or contact with an environment contaminated with faeces. Pigs may be an important reservoir of infection. Infections in the UK are associated with overseas travel. Large waterborne outbreaks.   | Freshwater/seawater – Strong                      | Waterborne                          | OI  |
| <i>Leptospira</i> spp.                             | Tightly coiled spiral bacteria that cause Weil's disease (jaundice) in people. Infection is from rodents and agricultural and domestic animals, usually through exposure to contaminated water or urine. Drinking or exposing wounds or mucous membranes to contaminated water can result in infection. Infection through natural water contaminated by rodent urine and occasionally through non-chlorinated drinking water. Outbreaks follow heavy rainfall and flooding, and occasionally abnormally low rainfall.   | Freshwater – Strong                               | Waterborne                          | OI; RILO; SA; RCS; NBM; CSS                 |
| <i>Microsporidia</i>                               | <i>Enterocytozoon bieneusi</i> infection is linked to transmission through food and water. Encephalitozoon hellem keratoconjunctivitis possibly related to water or mud. Link to rainy season in Singapore.   | Freshwater/seawater – Weak                        | Waterborne                          | RSA   |
| <i>Naegleria fowleri</i>                           | Colonises thermally polluted waters. Infections in the Southern USA are seasonal, with more in the summer. Infections in cattle are also seasonal. Infections may increase in some countries with warmer temperatures. Runoff from heavy rains introduces this organism into lakes, ponds and surface waters.   | Freshwater – Strong, links to water contamination | Waterborne                          | SCS   |
| <i>Norovirus</i>                                   | Is mostly transmitted person to person. Transmission has also been indicated via contaminated ice, stored water on cruise ships, borehole water and contaminated recreational bathing waters. Municipal drinking-water supplies have been implicated in outbreaks of gastroenteritis, usually following contamination by sewage. Strongly seasonal. Link to shellfish contaminated from infected faeces. Coastal water contamination linked to rainfall.  | Freshwater/seawater – Strong                      | Waterborne                          | OI  |

| Pathogen  | How climate might affect disease occurrence  | Strength of evidence for water-related infections   | Type of infection route               | Type of study linking climate and infection |
|---|--|---|---------------------------------------|---|
| <i>Rotavirus</i>                                    | Rotaviruses are part of the Reovirus family and have a double-stranded RNA genome. Exposure is by contact with infected individuals or contaminated water or other materials. Group C rotaviruses have been identified throughout the world. Group B rotaviruses have caused large outbreaks of diarrhoeal illness in mainland China. The virus entered the population as a result of faecal contamination of water supplies drawn from rivers, and then spread through the population by person-to-person contact. Waterborne outbreaks in developing countries.  | Freshwater/seawater – Weak  | Waterborne                            |   |
| <i>Sapovirus</i>                                    | A calicivirus, formerly called 'Sapporo-like virus', is a classic or typical calicivirus and is associated with relatively mild gastroenteritis in children. Outbreak linked to flood-water contamination of shellfish with several viruses.   | Freshwater – Weak   | Unknown                               | POWE  |
| <i>Schistosoma</i> spp.                             | These are flukes (helminth) that are transmitted through the contamination of water with faeces containing the ova. Cases linked to flooding and land surface temperature.   | Freshwater – Strong   | Water based                           | OI; SA; RRM; MM; MLM                        |
| <i>Schistosoma japonicum</i>                        | Infection is found in Eastern Asia including Japan and Korea. Links to rainfall and temperature.   | Freshwater – Strong   | Water based                           | CSS   |
| <i>Schistosoma mansoni</i>                          | The life cycle involves the ova hatching and infecting specific snail species, and the cercaria infect people occupationally or recreationally exposed to contaminated water through the skin.   | Freshwater – Strong   | Water based                           | OI; SA                                      |
| <i>Toxoplasma gondii</i>                            | A protozoan parasite that occurs in a wide range of warm-blooded animals. The only definitive hosts in which the full sexual cycle has been observed are members of the cat family (Felidae), which excrete the oocysts which contaminate the environment and source waters. People can be infected from consuming food or water that is contaminated with oocysts or the consumption of undercooked meat which contains tissue cysts. Infection can be a particular problem for pregnant women and immunocompromised patients. Some evidence that heavy rainfall can precede outbreaks.   | Freshwater/seawater – Strong  | Waterborne                            | OI; SA                                      |
| <i>Vibrio cholerae</i>                              | Causes cholera, a disease that is characterised by acute and life-threatening diarrhoea and dehydration usually in epidemic outbreaks. Cholera is transmitted through drinking water, shellfish and contaminated food. The disease is usually restricted to less developed countries where drinking water and waste disposal are poor, and to migrant populations associated with drought, flood, famine and war. Evidence of links to rainfall over the past century.   | Freshwater/seawater – Strong  | Waterborne                            | GAMTS; EACO; POWE                           |
| <i>Vibrio parahaemolyticus</i>                      | Inhabits estuarine and marine environments. It can cause food poisoning through the contamination of seafood. <i>V. parahaemolyticus</i> associated with raised water temperature.   | Seawater – Moderate   | Foodborne through seafood             | RSE   |
| <i>Vibrio vulnificus</i>                            | <i>Vibrio vulnificus</i> can cause severe, soft tissue infections, septicaemia and deaths. Infection is through the consumption of contaminated seafood (particularly raw oysters). <i>V. vulnificus</i> infection increased following hurricane Katrina.  | Seawater – Strong   | Waterborne; Foodborne through seafood | MMF; HSM; MMST; OI                          |
| <i>Vibrio</i> spp. (other than <i>V. cholerae</i> ) | A variety of <i>Vibrio</i> spp. can cause human disease, including the halophilic <i>V. parahaemolyticus</i> , <i>V. fluvialis</i> and <i>V. cholerae</i> , and the non-halophilic vibrios non-O1 <i>V. cholerae</i> and <i>V. mimicus</i> . Cholera is a classical waterborne disease, and the water route is still important in developing countries. There is no evidence that vibrios are able to cause human disease by growing within water distribution systems. <i>Vibrio</i> spp. are part of normal marine flora and can be found in marine, estuarine and river water. These organisms proliferate during the summer months. People are infected through the consumption of raw or undercooked contaminated shellfish, other foods and faecally contaminated water. A large infective dose is required to initiate infection, and person-to-person transmission does not occur. Infections in the UK tend to be in travellers returning from developing countries. Non-cholera <i>V. cholera</i> in warmer Baltic waters. | Freshwater ( <i>V. cholerae</i> )/seawater ( <i>Vibrio</i> spp. including <i>V. cholerae</i> ) – Strong | Waterborne                            | MMO; OI; TSAT; WMR; POTA                    |

Notes:

CA: cost analysis; CEO: circumstantial evidence only; CSS: cross-sectional survey; EACI: ecological association between climate and infections; EACO: ecological association between climate and outbreaks; FTA: fault tree analysis; GAMTS: generalised additive model time series; GDSE: gastrointestinal and dermatological symptoms and exposure; HSM: hindsight suitability model; LR: literature review; MLM: multi-level modelling; MM: mathematical modelling; MMF: microbiological monitoring of flooding; MMO: microbiological monitoring of outbreaks; MMST: microbiological monitoring with salinity and temperature; NBM: negative binomial model; OI: outbreak investigation; PMCC: pairwise-matched case-control study; PORA: post-outbreak rainfall analysis; POTA: post-outbreak temperature analysis; POWE: post-outbreak water examination; RA: review of animal infections; RCS: retrospective cohort study; RILO: rodent investigation linked to outbreak; RRM: rainfall-runoff model; RSA: rainy-season association; RSE: recreational-swimming exposure; SA: spatial analysis; SCS: sporadic case series; SFA: seasonal-factor analysis; TSAT: time-series analysis of temperature; QMRA: quantitative microbial risk assessment; WMR: water microbiology and rainfall

(a) Mode, type of evidence and strength assessment of climate influence are based on Nichols et al. (2018).

## SM3.5 Section 3.6

### SM3.5.1 Assessment of the Potential of Marine and Coastal Adaptation Solutions

The text below supports Figure 3.23 and Tables 3.28, 3.29 and 3.30 in Section 3.6.2.

#### SM3.5.1.1 Socio-Institutional Adaptation

##### *Knowledge diversity*

There is *high agreement* that successful ocean and coastal adaptation needs to incorporate the diversity of knowledge systems, moving beyond technical and scientific knowledge to also incorporate Indigenous knowledge and local knowledge (IKLK) (Norström et al., 2020; Petzold et al., 2020; Gianelli et al., 2021; Schlingmann et al., 2021). Knowledge diversity guarantees an enriching understanding of ecological, technical and political aspects of adaptation (Norström et al., 2020) while conciliating nature world views (Coscieme et al., 2020). Considering IKLK systems is more beneficial for the communities (*high confidence*) (Nalau et al., 2018; Sultana et al., 2019; Owen, 2020; Gianelli et al., 2021), increases their resilience (*robust evidence*) (Ford et al., 2020) and is relevant and transferable beyond the local scale (*medium evidence*) (Schlingmann et al., 2021). Implementing knowledge diversity in marine and coastal systems adaptation ranges from ecotourism (Section 3.6.3.1.3) to conservation (Section 3.6.3.2.1) and from Small Island Developing States (SIDS) to the Arctic (Section 3.6.3.4.1).

##### *Socially inclusive policies*

Socially inclusive policies that promote participation of all groups are able to address existing vulnerabilities in coastal communities, and promote adaptation and transformational change (*high agreement, low evidence*) (Brodie Rudolph et al., 2020; Ford et al., 2020; Friedman et al., 2020). Examples are described in Sections 3.6.3.4.1 and 3.6.3.4.2.

##### *Participation*

Participation in decision making and adaptation processes is recommended across a range of different hazards and contexts (Brodie Rudolph et al., 2020; Claudet et al., 2020; Sumaila et al., 2021) and has the potential to improve adaptation outcomes (*medium confidence*) (Hügel and Davies, 2020). Section 3.6.3 shows some examples of participation for fisheries and mariculture (Section 3.6.3.1.2), and in Indigenous communities (Section 3.6.3.4.1).

##### *Livelihood diversification*

Diversification of livelihoods is a common response strategy in coastal communities exposed to climate impacts such as coastal flooding, ocean extreme events, and changes in abundance and diversity of marine resources for food or income (*high confidence*) (Mohamed Shaffril et al., 2020; Owen, 2020; Biswas and Mallick, 2021). Livelihood diversification entails a transformative change (Barnes et al., 2020; Biswas and Mallick, 2021) whereby dependence on marine systems is alleviated by engaging in additional sources of

income, formally or informally, temporarily or permanently. Evidence shows that livelihood diversification in marine- and coastal-systems-dependent communities reduces climate risks (Mohamed Shaffril et al., 2020; Owen, 2020; Pinsky, 2021) and confers flexibility to individuals, which is key for adaptive capacity (*medium evidence*) (Blanchard et al., 2017; Cinner and Barnes, 2019; Taylor et al., 2021). However, diversification depends on the agency of the individuals, and existing vulnerabilities and inequities can cause diversification to result in maladaptive outcomes (*high agreement*) (Cinner and Barnes, 2019; Ford et al., 2020; Ojea et al., 2020). Therefore, we assess with *medium confidence* the ability of livelihood diversification alone to address the impacts of climate change in coastal communities. Livelihood diversification as a response to climate change is further assessed in fisheries and mariculture (Section 3.6.3.1.2), coastal communities (Cross-Chapter Box SLR in Chapter 3) and tourism (Section 3.6.3.1.3).

##### *Mobility*

Mobility of coastal livelihoods is a common practice in many regions of the world, such as the Pacific Islands (Chapter 15), that has been reported as a response to climate-change impacts such as coastal flooding and extreme events. When individuals are given the choice about mobility, they use this response in order to minimise climate risks and benefit their livelihoods (*medium evidence*) (Barnett and McMichael, 2018). An example of mobility includes fishing (Section 3.6.3.1.2).

##### *Migration*

Coastal livelihoods facing severe climate-change impacts often respond with migration as a critical livelihood diversification strategy (Maharjan et al., 2020; Biswas and Mallick, 2021; Zickgraf, 2021). Migration often involves different spatio-temporal scales than mobility (Barnett and McMichael, 2018), and it could be considered an adaptation solution for some coastal and island populations in the cases of extreme events, but also as a response to more gradual changes (e.g., coastal erosion from SLR) (Zickgraf, 2021). There is *low confidence* that migration alone can be a resilient response to climate-change impacts in ocean and coastal systems (Section 3.6.3.1.1). The maladaptive outcomes of mobility and migration (and relocation) (Section 3.6.3.1.1) are influenced by the mobility of vulnerable groups, the regions where the displacement occurs and the capacities that these individuals possess (Dandy et al., 2019; Maharjan et al., 2020). Examples of migration include island dwellers (Section 3.6.3.1.1).

##### *Finance and market mechanisms*

Financial mechanisms and credit provision for marine-dependent livelihoods are effective for overcoming impacts from SLR (Hinkel et al., 2018; Moser et al., 2019; Woodruff et al., 2020) and extreme events (*medium evidence*) (Shaffril et al., 2017; Dunstan et al., 2018; Sainz et al., 2019). Market mechanisms include payment for ecosystem services that can directly or indirectly, through mitigation, contribute to adaptation outcomes in marine and coastal systems (Cross-Chapter Box NATURAL in Chapter 2; Himes-Cornell et al., 2018; Brathwaite et al., 2021). There is *high confidence* in the potential of improved financial and market mechanisms for ocean adaptation, as they are key

for transitioning into future ocean sustainability (Chapter 18; Sumaila et al., 2021). Examples of implementation of finance and economic mechanisms are further assessed in Section 3.6.3.4.2.

#### *Disaster response programmes*

In the occurrence of coastal and ocean extreme events, coastal communities and marine-dependent livelihoods can respond following existing disaster response programmes, which confer resilience to communities and contribute to adaptation when designed to be inclusive, participatory and adaptive (*high confidence*) (Nurhidayah and McLgorm, 2019). Disaster response programmes need adequate finance that combines viable economic investments and risk-financing instruments (Jongman, 2018). Recent evidence suggests that the analysis and understanding of communications data during disaster responses is key to avoid maladaptation outcomes (*high agreement*) (Nichols et al., 2019). Disaster response programmes can be implemented with climate services (Section 3.6.3.4.3), and examples include the tourism cruise-ship sector (Section 3.6.3.1.3).

#### *Multi-level ocean governance*

The multi-scale nature of ocean and coastal climate-change risk demands adaptation solutions at multiple levels of governance (*high confidence*) (Miller et al., 2018; Gilfillan, 2019; Holsman et al., 2019; Obura et al., 2021), which consider the objectives and perceptions of all stakeholders to support local implementation of broad strategies (Miller et al., 2018). However, current ocean governance is complex and fragmented (Scobie, 2019b; Haas et al., 2021), and faces challenges that intersect with climate adaptation. Therefore, there is *high agreement* in the need to transition into multi-level governance to respond to the challenges of climate change in the oceans (Chapter 18; Haas et al., 2021).

#### *Institutional transboundary agreements*

At the international level, institutional agreements for the management of transboundary marine resources are key for a sustainable future given current impacts on marine species distribution due to climate change (*high agreement*) (Mason et al., 2020; Melbourne-Thomas et al., 2021). Existing climate-adaptive transboundary agreements are scarce (Melbourne-Thomas et al., 2021; Sumbay et al., 2021) and need to be redesigned in order to address the challenges of climate-induced marine species distributional changes (*medium confidence*) (Engler, 2020; Oremus et al., 2020). Despite the need for new instruments, adhering to existing ocean-conservation and resource-management international agreements contributes to sustainable ocean futures and climate-change adaptation (Haas et al., 2021). Examples are implemented in fisheries (Section 3.6.3.1.2; Cross-Chapter Box MOVING SPECIES in Chapter 5).

### SM3.5.1.2 Built Infrastructure and Technology

#### *Accommodation and relocation*

Asset accommodation and relocation in the marine and coastal environment are among the most commonly discussed adaptations to climate hazards such as SLR and coastal extreme events (Hanson and Nicholls, 2020; Monios and Wilmsmeier, 2020). Planned relocation is a response to extreme events and SLR in coastal regions, and it has the advantage of maintaining community and social structures (Zickgraf, 2021) but can lead to maladaptation in cases where individuals are not included in decision making and where no monitoring exists (Zickgraf, 2021). While there is *high confidence* that relocation reduces coastal risk, there are important social and economic costs linked to such interventions (Cross-Chapter Box SLR in Chapter 3) and they require transformative changes in the longer run (Magnan et al., 2020).

#### *Protection and beach-and-shore nourishment*

As an alternative to hard structures, a common response to beach erosion around the world is beach nourishment (Barbier, 2014; Oppenheimer et al., 2019; Pinto et al., 2020; Toimil et al., 2020; Elko et al., 2021). This practice involves supplementing existing beach sediments with sediment sourced from adjacent ecosystems. Part of the appeal of beach nourishment is that it provides relatively rapid results, but it invariably also entails poorly quantified trade-offs between efficacy, long-term cost, utility to beach users and ecological damage (de Schipper et al., 2021). Protection with seawalls and beach-and-shore nourishment constrains the development of ecosystem-based adaptation and the associated co-benefits, and can have negative consequences on coastal ecosystems (Sections 3.4.2.4–3.4.2.6; Cross-Chapter Box SLR in Chapter 3). However, protection may be a feasible alternative for densely populated and urbanised coastal areas (Barnard et al., 2021; Neijnens et al., 2021). Therefore, there is *medium confidence* on the ability of protection with beach and shore nourishment to reduce climate-change impacts in coastal systems. Combined solutions, including soft or nature-based infrastructure (Section 3.6.2.3) and socio-institutional approaches (Section 3.6.2.2), achieve better outcomes when implemented together (Cross-Chapter Box SLR in Chapter 3;Gattuso et al., 2018; Foti et al., 2020; Reguero et al., 2020; Seddon et al., 2020). Examples of implementation of these solutions are further assessed in Section 3.6.3.1.1.

#### *Early warning systems*

AR5 and SROCC mention that early warning systems can support decision making, decrease economic losses from extreme events, and aid in the enterprises and development of adaptive management systems for coastal systems and fisheries (Hoegh-Guldberg et al., 2014; Bindoff et al., 2019; Collins et al., 2019). Such systems have potential risks in the form of erroneous forecasts, resulting in unintended consequences (e.g., forecast leading to increased exploitation of a resource or decrease in tourism) and exacerbation of inequality due to geographic unevenness of development and access (Section 3.6.3.2.4; Soares et al., 2018). Early warning systems may also be at risk of disruption by non-climate events as demonstrated by the recent

interruptions of maintenance and monitoring of ocean- and coastal-observing systems owing to COVID-19 (Northrop et al., 2020).

### *Seasonal and dynamic forecasts*

Rapid expansion of hindcast data, remote-sensing data and computational power have led to the proliferation of real-time and seasonal forecasts of temperature extremes, MHWs and their impacts (Liu et al., 2018; Holbrook et al., 2020; Spillman and Smith, 2021), storm surges and wave-driven flooding (Fernández-Montblanc et al., 2019; Winter et al., 2020), water quality and HABs (Bever et al., 2021; Davidson et al., 2021), and the distribution of living marine resources (Payne et al., 2017; Tommasi et al., 2017; Hazen et al., 2018). These tools have the capacity to contribute to monitoring, early warning systems, adaptive management and ecosystem-based management, as improvements in the spatial scale for management are being achieved (*high confidence*) (Tommasi et al., 2017). There is *high confidence* that capacity-building and broad engagement of stakeholders from communities, governments and industries are critical to creating early warning systems with forecasts that can be properly interpreted by users and be effectively incorporated into monitoring, management and decision making (Section 3.6.3.2.4).

### *Monitoring*

Monitoring systems that address climate-change hazards, ecosystem impacts and social vulnerabilities in marine social–ecological systems are a key priority for adaptation to climate hazards in coastal areas (*high agreement*) (Nichols et al., 2019; Claudet et al., 2020; Wilson et al., 2020). For ocean extreme events, specific event-based monitoring can help managers and stakeholders in decision making in real time (Holbrook et al., 2020). Monitoring and detection of marine species range shifts is key to adaptation (Melbourne-Thomas et al., 2021). However, monitoring efforts and costs differ across regions, where the most remote ocean systems regularly lack such actions (Claudet et al., 2020) and there is a lack of standardised methods and open-access information in global terms (Rilov et al., 2020; A. Maureaud et al., 2021; Melbourne-Thomas et al., 2021); therefore, there is *medium confidence* on the potential of monitoring systems alone in supporting adaptation in marine and ocean systems. Examples of implemented monitoring systems are shown for MPAs (Section 3.6.3.2.1), climate services (Section 3.6.3.2.4) and fisheries (Section 3.6.3.1.2).

### *Ecological interventions: habitat development, active restoration and assisted evolution*

The near-term nature of climate risk to coastal systems has led to increased research and investment in technological interventions to support and enhance the capacity of species and ecosystems to adapt to climate change (Jones et al., 2018; Boström-Einarsson et al., 2020; Kleypas et al., 2021). Unlike traditional marine conservation ('passive') approaches (Section 3.6.2.3), which aim to allow ecosystems to naturally recover from disturbance, ecological interventions and engineering approaches aim to accelerate recovery of damaged systems and promote ecological or biological adaptation to future climate change (Jones et al., 2018; Kleypas et al., 2021). Artificial habitat development, for example, has the potential to stabilise shorelines and increase fisheries productivity in rocky reef systems (Gilby et al., 2018). Active restoration involves reintroducing

species or augmenting existing populations, for example, propagating and transplanting heat-tolerant coral species (Boström-Einarsson et al., 2020; Rinkevich, 2021). More controversial interventions like assisted evolution, in which genes are manipulated to accelerate natural selection, has been investigated for corals (National Academies of Sciences, 2019), kelp (Morris et al., 2020) and other habitat-forming species (Bulleri et al., 2018). The effectiveness and feasibility of these solutions are further assessed for existing restoration efforts in Section 3.6.3.2.2.

### SM3.5.1.3 Marine and Coastal Nature-Based Solutions

#### *Habitat restoration*

Marine habitat restoration increases biodiversity (*high confidence*) (McLeod et al., 2019) and protects shorelines and coastal livelihoods from climate oceanic hazards in the near term (*high confidence*) (Colls et al., 2009; Arkema et al., 2017; Espeland and Kettnerring, 2018; McLeod et al., 2019). To date, restoration efforts have predominantly focused on typically productive coastal habitats, including coral reefs, seagrasses, mangroves, salt marshes, oyster reefs and kelp forests (Bayraktarov et al., 2016; Espeland and Kettnerring, 2018; Gilby et al., 2018; Reguero et al., 2018; McLeod et al., 2019; Duarte et al., 2020a) that provide critical nursery or foraging habitats for commercially or culturally important species. Although habitat restoration can enhance fish-stock production (Ermgassen et al., 2016; McLeod et al., 2019), bring sociocultural benefits by promoting stewardship and community involvement (Hein et al., 2019), and benefit recreational use and tourism (Section 3.6.3.2; Weatherdon et al., 2016; Hoegh-Guldberg et al., 2018; Agyeman, 2019), there is *limited evidence* that it can safeguard fish-stock production in future climate conditions (McLeod et al., 2019) or restore coastal habitats after conversion to alternate states (Sections 3.4.2.1, 3.4.2.3, 3.6.2.3; Hein et al., 2017; Fox et al., 2019; Hein et al., 2021). There is also substantial evidence that simply restoring habitats to mid-20th century states will not enable them to persist in the medium term without substantial emissions reductions (*medium to high confidence*) (Sections 3.4.2.1, 3.4.2.3, 3.4.2.5, 3.4.2.6), because benefits are challenging to quantify and include time-delayed responses associated with the rebuilding of biological communities (McLeod et al., 2019). Finally, habitat restoration can limit loss of ecosystem services related to recreational use and traditional tourism (*medium confidence*) (Weatherdon et al., 2016; Hoegh-Guldberg et al., 2018) while benefiting ecotourism (Agyeman, 2019). (See an assessment of implemented restoration efforts in marine systems in Section 3.6.3.2.2.)

#### *Marine protected areas and OECMs*

Marine protected areas (MPAs) are the most widely implemented approach to conserving marine biodiversity and have long provided the foundation for initiatives at local to international scales (Rilov et al., 2020; Arafeh-Dalmau et al., 2021). These MPAs and networks of MPAs that are well designed and enforced provide well-known conservation and sociocultural benefits to people, because they protect biodiversity and ecosystem functioning that support delivery of important ecosystem services, including food supply, recreation, scenic beauty and water regulation (Section 3.5; Edgar et al., 2014; Gill et al., 2017; Wilson et al., 2020; Ovando et al., 2021; Sala et al., 2021). However, the effectiveness of current MPA networks to

assist in climate-change adaptation is equivocal (Tittensor et al., 2019; Wilson et al., 2020), as climate-change impacts are rarely incorporated in management (*high confidence*) (Section 3.6.3.2.1; Rilov et al., 2020; Arafah-Dalmau et al., 2021). If carefully designed to address climate change, strategically placed and well enforced, they hold great potential to deliver better adaptation outcomes (*high confidence*) (e.g., Queirós et al., 2016; Roberts et al., 2017; Maxwell et al., 2020a; Arafah-Dalmau et al., 2021; Sala et al., 2021). An additional spatial-conservation instrument that contributes to ocean protection and adaptation of coastal livelihoods are the other effective area-based conservation measures (OECDMs) (Gurney et al., 2021). These are areas managed by ocean-dependent communities that are recognised for the contribution of such management and interaction to marine conservation (Maxwell et al., 2020b; Gurney et al., 2021). Recognising these areas can benefit adaptation through the increased ecological and social resilience that such management regimes confer (Section 3.6.3.2.1).

#### *Conservation of climate refugia*

Marine regions that retain climate and biodiversity conditions for longer periods of time under climate-change impacts are considered climate refugia (Wilson et al., 2020; Arafah-Dalmau et al., 2021). There is *low evidence* but *high agreement* (*medium confidence*) that protecting these areas can increase the resilience of marine ecosystems in the face of ocean warming and MHW (Rilov et al., 2020; Arafah-Dalmau et al., 2021), facilitate marine species shifts (Cross-Chapter Box MOVING SPECIES in Chapter 5; Wilson et al., 2020) and help avoid species extinctions and extirpations. But it is not recommended as the only climate adaptation solution for marine conservation (Tittensor et al., 2019; Wilson et al., 2020). Recent evidence points that the selection of marine climate-refugia areas needs to be guided by both socioeconomic criteria and broadly applicable climate-change metrics such as climate velocity (Brito-Morales et al., 2018; Arafah-Dalmau et al., 2021).

#### *Transboundary MSP and ICZM*

Marine spatial planning (MSP) is an often participatory process to allocate spatio-temporal distribution of human uses in the ocean, with the aim to reduce conflicts and reach sustainability (Gissi et al., 2019; Frazão Santos et al., 2020). After several decades of development of MSP initiatives and MSP processes in more than half of EEZs, 25 countries have already implemented or have government-approved plans, most of them in economically developed nations (Frazão Santos et al., 2020). The potential of MSP to incorporate adaptation goals is important but limited so far by the lack of consideration of climate change in the plans (*robust evidence*) (Gissi et al., 2019; Frazão Santos et al., 2020), and the lack of consideration of sociocultural goals (Pennino et al., 2021). MSP that incorporates climate-change impacts and adaptation in the design can contribute to support climate adaptation from a multi-sector perspective and through existing policy frameworks (*low evidence*) (Tittensor et al., 2019; Frazão Santos et al., 2020; Rilov et al., 2020). However, climate-resilient MSP would require a transformation in current policy systems, as plans would need to be dynamic and incorporate new jurisdictions, climatic predictions and novel expertise (Frazão Santos et al., 2020; Pennino et al., 2021). In this context, transboundary marine spatial planning (TMSP), a process

of international cooperation in the marine space to resolve conflicts across nations (Li and Jay, 2020), holds promise to adapt to the shifting nature of climate-change impacts in the oceans and make MSP robust to climate-change impacts (Pinsky et al., 2021). Integrated coastal zone management (ICZM or ICM) differs from MSP in that it focuses on the land–sea interface (Frazão Santos et al., 2020). Recent evidence supports the need to integrate disaster response programmes and adaptation goals in ICZM (*low evidence*) (Rosendo et al., 2018), and to overcome the existing implementation challenges for governments to be able to use ICZM for climate-change adaptation (Rosendo et al., 2018). Examples of MSP and ICZM implementation are further assessed in Section 3.6.3.1.3 (tourism) and Section 3.6.3.2.1 (conservation).

#### *Sustainable harvesting*

Harvesting marine resources at rates that are ecologically sustainable increases the resilience of marine systems to climate change, while providing important ecosystem services to marine-dependent communities (Chapter 5). Sustainable harvesting is recognised as a nature-based solution that contributes to adaptation by safeguarding the provision of marine food services (Section 3.5.3), related cultural services (Section 3.5.6) and food security, while reducing marine systems' ecological vulnerability (*high confidence*) (Gattuso et al., 2018; Burden and Fujita, 2019; Duarte et al., 2020a). Further assessment is performed for fisheries and mariculture practices (Section 3.6.3.1.2).

#### *Climate-adaptive management*

Climate-adaptive management of ocean and coastal resources allows to iteratively update management with climate knowledge and information available for the system, with observed and projected changes in the environment and with the experience of 'learning by doing' (Rilov et al., 2020; Wilson et al., 2020). There is *high agreement* on the need to incorporate climate-adaptive management in marine resources in order to adapt to the impacts of species distribution (Cross-Chapter Box MOVING SPECIES in Chapter 5; Wilson et al., 2020; Melbourne-Thomas et al., 2021) and other climate-change multiple hazards (Rilov et al., 2019). There are examples of implementation of adaptive management in many contexts such as MPAs (Nickols et al., 2019), fisheries management and mangroves (Sections 3.6.3.1.2, 3.6.3.2.1, 3.6.3.2.2; Ellison et al., 2020).

#### *Ecosystem-based management*

Ecosystem-based management is an approach to manage ocean and coastal systems that focuses on the habitats and ecosystems as management units, where initiatives can follow a set of key principles (Long et al., 2015). It incorporates many of the above-mentioned tools and processes (Harvey et al., 2018), such as participatory processes, conservation tools, MSP and ICZM, adaptive management and sustainable harvesting, among others. By incorporating climate-adaptive measures and focusing on the ecosystems, this approach benefits the adaptation of marine ecosystems and supports the provision of ecosystem services under climate change (*high confidence*) (Fernandino et al., 2018; Lowerre-Barbieri et al., 2019). When developing nature-based adaptation measures, there is an increase in the ecosystem resilience and a decrease in socioeconomic

vulnerability to climate change (Miller et al., 2018; Holsman et al., 2019; McLeod et al., 2019; Seddon et al., 2020). An increasing number of vulnerability and risk assessments of climate change and impacts of multiple stressors on species, habitats and natural communities (e.g., Holsman et al., 2017; Pinsky et al., 2019; Samhouri et al., 2019; Gissi et al., 2021), on ecosystem services (Kurniawan et al., 2016; Stewart-Sinclair et al., 2020a), and on associated vulnerabilities and risks to social systems (Gaichas et al., 2018; Thiault et al., 2019), support the development of NbS for adaptation.

#### EBA hotspots

Mangroves and coral reefs have been identified as EbA hotspots (Figure 3.25) because they line shores that are highly vulnerable to tropical storms and SLR, and they protect at least 5.3 and 3.4 million people living within 2 km of the coast, respectively (Section 3.4.2.5; Cross-Chapter Box SLR in Chapter 3; Jones et al., 2020; Menéndez et al., 2020; Van Coppenolle and Temmerman, 2020). At least 38% of mangroves that intersect with EbA hotspots are under some level of protection (Jones et al., 2020), which is especially important considering recent analyses which show global yearly emissions of 23.5–38.7 Tg yr<sup>-1</sup> due to losses of mangroves (Ouyang and Lee, 2020). Greater levels of protection would improve the potential of these EbA, especially in countries with low adaptive capacity (Friess et al., 2019), and this potential could be further enhanced by incorporating mangrove restoration (Jones et al., 2020; Menéndez et al., 2020).

#### SM3.5.2 Fisheries Adaptation to Climate Change

Adaptation to climate impacts in fisheries and mariculture most commonly begins with harvesters moving. Mobility allows fishing fleets and fishers to adapt to shifting marine species distributions (*high agreement*) (Section 3.5.3, 3.4.3.1; Peck and Pinnegar, 2018; Pinsky et al., 2018; Frazão Santos et al., 2020). Following preferred species is challenging: well-developed, regionally coordinated fisheries can fail to match climate-driven range shifts of target species (Pinsky and Fogarty, 2012); in some cases, target species may vacate designated fishing grounds (Bell et al., 2021). An alternative is diversifying fishing locations, as occurred when the squid fishery in Baja California Sur (Mexico) faced drastic decreases in catches before collapsing (Gonzalez-Mon et al., 2021); however, fishery relocation can disrupt cultural associations or worsen scarce employment opportunities (e.g., Jarre et al., 2013). Even mobile oceanic fisheries, like the industrial tuna fleet, face falling revenues from tracking shifting target species (Bell et al., 2021). Overall, there is *limited evidence* on the positive or maladaptive implications of relocation (Magnan et al., 2020; Islam et al., 2021).

Diversification of harvests, tactics and livelihoods is a common adaptation strategy that can help address climate impacts to fisheries and mariculture (*high confidence*) (Galappaththi et al., 2017; Blair and Momtaz, 2018; Miller et al., 2018; Peck and Pinnegar, 2018; Robinson et al., 2020; Gonzalez-Mon et al., 2021). Harvest diversification increases yields and livelihood stability in commercial and small-scale fisheries (*high confidence*) (Cline et al., 2017; Young et al., 2019; Barnes et al., 2020; Robinson et al., 2020). Tactics may include changing fishing gear or vessel power, or combining different income-generating

activities within the fishing and mariculture sectors (Ojea et al., 2020). Livelihood diversification includes transitioning from wild fisheries to mariculture (Gephart et al., 2020; Ojea et al., 2020), or to other marine sectors like ecotourism, or leaving marine sectors entirely. Coastal and offshore mariculture is expected to continue growing to meet increasing seafood demand (Costello et al., 2020a) despite constraints from climate change (Froehlich et al., 2018a; Froehlich et al., 2020) and sustainability (Corten et al., 2017; Theuerkauf et al., 2019; Costello et al., 2020a), and it requires its own set of adaptations, including technological (Froehlich et al., 2018b; Cottrell et al., 2020) and socio-institutional changes (Galappaththi et al., 2020). (A full assessment of mariculture and fisheries within food systems is found in Chapter 5.) Increasing tourism is a commonly reported diversification strategy for fishers (Wood et al., 2013; Cinner, 2014), but this action has a limited ability to improve or maintain the well-being of fishing communities (*high confidence*) (Lasso and Dahles, 2018; Fabinyi, 2020; Pham, 2020) as they become dependent on tourism flows and seasonality. Social capital and land tenure can support successful transitions among sectors (*low evidence*) (Diedrich et al., 2019; Fabinyi, 2020), and livelihoods also benefit from participating in co-management systems to confront the impacts of climate change (*medium evidence*) (Voorberg and Van der Veer, 2020; Gianelli et al., 2021).

Technology and infrastructure adaptations can improve marine-harvest efficiency, reduce risk and support resource-management goals (Friedman et al., 2020; Bell et al., 2021; Melbourne-Thomas et al., 2021), but their ability to overcome climate-change impacts remains uncertain (Bell et al., 2020). For example, the tropical tuna industry uses aggregation devices to increase efficiency, but the ability of these devices to offset climatic impacts is uncertain (Rubio et al., 2020). Technologies like satellite-tracked vessel-monitoring systems facilitate enforcement of marine reserves (Bradley et al., 2019; White et al., 2020) and identify illegal, unreported and unregulated fishing activity (Lubchenco and Grorud-Colvert, 2015; Cabral et al., 2018; Kurekin et al., 2019), thereby helping alleviate overfishing, a major detrimental interacting driver (Section 3.5.3); however, the degree to which these technologies can overcome expected climate-change impacts (Bell et al., 2013; Erauskin-Extramiana et al., 2019) is unquantified, as are any associated ecological consequences.

Improving capacity to predict anomalous conditions in coastal and marine ecosystems (Jacox et al., 2019; Holbrook et al., 2020; Jacox et al., 2020), storm-driven flooding in reef-lined coasts (Scott et al., 2020; Winter et al., 2020) and fisheries-stock recruitment (Payne et al., 2017; Tommasi et al., 2017; Muhling et al., 2018) can improve forecasts of coastal and marine resources. These can enhance sustainability of wild-capture fisheries (*high confidence*) (Blanchard et al., 2017; Tommasi et al., 2017) and inform fisheries and mariculture decision making at management-relevant time scales (Payne et al., 2017; Tommasi et al., 2017; Hobday et al., 2018). Combining seasonal and multi-year climate projections in dynamic-model forecasts helps marine industries consider longer-term adaptations such as changing practices or relocating (*medium confidence*) (Tommasi et al., 2017; Hobday et al., 2018; Merryfield et al., 2020). These tools will be most effective when they are accessible to decision makers and local communities (see Section 3.6.3.3; Tommasi et al., 2017; A. Maureaud et al., 2021).

Table SM3.3 | Background materials and data for Figures 3.23 and 3.24 (includes levels of feasibility and effectiveness in Figures 3.23 and 3.24; includes the full feasibility assessment in Figure 3.24)

| Adaptation solution         | Technical and economic feasibility   | Supporting references (Technical and economic feasibility)              | Institutional and geophysical feasibility   | Supporting references (Institutional and geophysical feasibility) | Social–ecological feasibility   | Supporting references (Social–ecological feasibility)   | Feasibility (general) | Effectiveness   | Supporting references (Effectiveness)  |
|-----------------------------|--|---|---|---|---|---|-----------------------|---|--|
| Knowledge diversity         | High: Incorporation of various knowledge systems is at the early stages of implementation but is transversal to other systems and sectors (i.e., agriculture, biodiversity conservation, etc.). It is in the phase of adaptation planning and early implementation. There are no high associated economic costs; unknown productivity; low technical requirements. | Pasquini and Cowling (2015); Dawson et al. (2020)                       | Medium: IKLK are transferable across regions. Institutions are not currently designed to incorporate knowledge diversity and act as a barrier, as do laws and regulations in many places of the world. Global institutions greatly support knowledge diversity, also the SDGs.        | Stephenson et al. (2016); Schlingmann et al. (2021)               | High: Environmental feasibility is high as knowledge diversity is related to higher ecological resilience. Socially, diversity of knowledge systems is also increasing social–ecological resilience and supporting ecosystem services related to culture and identity.  | FAO (2015a); Ross et al. (2019); Terra Stori et al. (2019); Ford et al. (2020)                            | High                  | Low: Needs combination with other solutions. Facilitates the implementation and success of adaptation tools such as EbA.                  | Nalau et al. (2018); Peñaherrera-Palma et al. (2018); Raymond-Yakoubian and Daniel (2018); Coscione et al. (2020); Owen (2020) |
| Socially inclusive policies | High: Technical readiness is medium based at an adaptation planning and early implementation stage; however, socially inclusive policies are transversal to systems and sectors, and their implementation is well known. Economic feasibility is high as no high costs are expected.   | Archer et al. (2014)  | Medium: Not applicable in the geographical context. Institutions can support at all levels social inclusion, and applications differ across countries while supported by the SDGs; however, current governance system may limit inclusive policies.                                   | Devereux (2016); Bennett (2018)                                   | High: Ecological feasibility is high as there are no known trade-offs between inclusive processes and biodiversity ecosystem services, and there is a win–win situation. Social feasibility is high as social aspects, including equity in access to resources and recognition and inclusion of all stakeholders within policy planning and implementation, are key to success. | Anderson (2015); Basel et al. (2020); McNamara et al. (2020); Ogier et al. (2020); Williams et al. (2020) | High                  | Low: Needs combination with other solutions. Facilitates finance and market mechanisms, monitoring systems, among others, for adaptation. | Tommasi et al. (2017); Claudet et al. (2020); A. Maureaud et al. (2021); Sumalla et al. (2021)                                 |
| Participation               | High: broadly implemented approaches, trans-sectorial, with low economic and technological constraints. New successful approaches to co-generate adaptation solutions.   | van der Voorn et al. (2017); Flood et al. (2018); Johnson et al. (2020) | Low: Unequal opportunities for co-management and participation across institutions, low co-management in industrial fisheries, tourism or marine conservation and restoration. Not largely implemented in existing governance across sectors. Geophysical feasibility not applicable. | Nursey-Bray et al. (2018); Brodie Rudolph et al. (2020)           | High: Increases environmental resilience and reduces inequities due to secure access to resources. Allows for co-management systems for marine resources. Improves education.   | Ojea et al. (2017); Koengstein et al. (2020); Voorberg and Van der Veer (2020); Gianelli et al. (2021)    | Medium                | High: Potential to improve adaptation outcomes and identify impacts and adaptation needs.   | Rumore et al. (2016); Hügel and Davies (2020)  |

| Adaptation solution        | Technical and economic feasibility  | Institutional and geophysical feasibility  | Supporting references (Institutional and geophysical feasibility)                      | Social–ecological feasibility   | Supporting references (Social–ecological feasibility)                        | Feasibility (general) | Effectiveness  | Supporting references (Effectiveness)   |
|----------------------------|---|--|--|---|--|-----------------------|--|---|
| Livelihood diversification | Medium: Implementation expanding, low technical constraints; however, economic constraints given the available income deriving alternatives in place, which will require public and private investments (i.e., aquaculture, tourism, blue energy, etc.).  | Low: Rigid institutions and systems to allow livelihood diversification, unknown political acceptability, existing legal and regulatory actions (i.e., licenses). Requires transformational change.  | Peck and Pinnegar (2018); Ojea et al. (2020); Biwas and Mallick (2021)                 | Low: Environmentally has the potential of increasing pressure in the marine environment with other uses (if conducted non-sustainably). Socially it is constrained by the learning capacity of people and it has the risks of resource detachment, and loss of cultural identity and professional pride.  | Hossain et al. (2018); Cinner and Barnes (2019); Fabinyi (2020); Pham (2020) | Low                   | Medium: Reduces climate risks and provides adaptive capacity, but does not necessarily engage all individuals due to agency. | Cinner and Barnes (2019); Ford et al. (2020); Mohamed Shaffri et al. (2020); Ojea et al. (2020); Owen (2020); Pinsky (2021); Taylor et al. (2021) |
| Mobility                   | Medium: Early planning stage and little implementation. Mobility is a tradition for livelihoods in some specific regions. In fisheries, industrial fleets are able to move, but economic costs are often subsidised. Small-scale fleet do not have many mobility options, or they come at high costs.                                     | Medium: Mixed regulations and governance to favour mobility of livelihoods or within industries. Current regulations allow for high fish mobility in high-capacity fishing countries, but management regulations often do not match shifting fishing grounds.  | Cross-Chapter Box MOVING SPECIES in Chapter 5; Young et al. (2019); Bell et al. (2021) | Low: Environmental feasibility is low as it can diminish resource availability elsewhere. Social feasibility is low as conflicts can arise between communities, countries and industries, and increase vulnerabilities of specific groups (women, Indigenous Peoples, migrants). This is particularly problematic where the communities have long cultural associations with fisheries and/or where few other employment opportunities exist. | Jarre et al. (2013); Ojea et al. (2020); Gonzalez-Mon et al. (2021)          | Medium                | Medium: Given the choice and under specific circumstances, livelihoods engage in mobility to reduce risk.                    | Barnett and McMichael (2018)  |
| Migration                  | Medium: Adaptation planning and early implementation. Technical feasibility is moderate, but economic feasibility is low due to high costs of relocation, building new infrastructures, lost infrastructure spaces, etc. However, in some specific settings, migration (planned relocation) may be a lower cost solution than protection. | Low: Not always physically feasible due to borders and international regulations, institutions globally and at the national or regional level are not ready for transboundary relocation and migration, and can interfere with international agreements on human rights. Few countries have currently included migration in national climate-change commitments. | Wilkinson et al. (2016); Scoble (2019a)  | Low: Although migration can alleviate risks (i.e., SLR for human populations), it has substantial political, social and economic costs, and sometimes it is simply impractical, as in the case of coastal megacities.   | Gibbs (2015); Bordin et al. (2020)   | Low                   | Medium: Is a common response for incremental impacts and hazards, but often results in maladaptation outcomes.               | Biswas and Mallick (2021); Zickgraf (2021)  |

| Adaptation solution           | Technical and economic feasibility   | Institutional and geophysical feasibility  | Supporting references (Institutional and geophysical feasibility)                | Social–ecological feasibility                    | Supporting references (Social–ecological feasibility)  | Feasibility (general)   | Effectiveness | Supporting references (Effectiveness)  |
|-------------------------------|--|--|--|--|--|---|---------------|--|
| Finance and market mechanisms | High: Known technical feasibility from adaptation planning and early implementation. High economic feasibility from cost-effective market mechanisms. Finance mechanisms require private/public investment and innovative solutions that have been already tested. | Medium: Multi-scale reach. Political acceptability varies across countries; existing regulations support mechanisms that are already functioning in other sectors and can be ready to transfer to the marine realm; varying transparency of processes. | Bott and Braun (2019); Ware and Banhami-Zakar et al. (2021)                      | Lowe et al. (2019)                               | Medium: Environmental feasibility is medium as these mechanisms can promote one ecosystem service and hinder others. Although they can be designed for win-win solutions (i.e. carbon and biodiversity), it is not always possible. Socially, these instruments usually do not recognise traditional livelihoods, minorities without land tenure and other social facets, and can exacerbate existing inequalities. Design in the mechanisms and equitable access can increase their feasibility, but more development of the instruments is needed. Barriers are greater in low- to mid-income countries. | Donner and Webber (2014); Hinkel et al. (2018); Klöck and Nunn (2019); Claudet et al. (2020); Ojea et al. (2020); Tompkins et al. (2020); Sumaila et al. (2021) | Medium        | High: Effective for reducing impacts of SLE and extreme events.  |
| Disaster response programmes  | High: Technical feasibility is high given the widespread implementation. Economic costs and investments can be significant, but benefits are high. It requires viable economic investments and risk-financing instruments.   | High: Scales of risk can match the scales of responses. Institutions allow the implementation and maintenance of disaster response programmes and are common in every nation and in international treaties.  | Stewart et al. (2015); Dawson et al. (2016); Jongman (2018); Quinn et al. (2019) | Rosendo et al. (2018)                            | High: High social and environmental feasibility as they address impacts in livelihoods, and they could also benefit ecosystem services provision. When inclusive, participatory and adaptive, they confer resilience to communities.   | Cvitancic et al. (2016); Nurindayah and McGlomm (2019)  | High          | Low: Solution reduces impacts; however, it depends on design and innovations, needs coordination with climate adaptation and generally has a science policy gap. |
| Multi-level ocean governance  | High: Allowing for multiple bodies and a polycentric governance system in the oceans is not necessarily constrained by economic costs and benefits of technology.  | High: Broad scalability, existing instruments and institutions; feasible in most countries where high governance and transparent decision making is present. Globally high feasibility. Required for land–sea interactions.                            | Armitage (2007)  | Mahon and Fanning (2019); Schlüter et al. (2020) | Medium: Social feasibility is medium as an asymmetrical distribution of power to make decisions for resource control or uses has impeded inter-agency collaboration. This is not necessarily always the case. Environmental feasibility is high as it is recognised to increase social–ecological resilience.  | Ho et al. (2012); Ojea et al. (2017); Miller et al. (2018)  | High          | High: Allows for integration of policymaking across levels for sectors such as fisheries. Polycentric governance is effective in adaptation.                     |

| Adaptation solution                        | Technical and economic feasibility   | Supporting references (Technical and economic feasibility)                       | Institutional and geophysical feasibility  | Supporting references (Institutional and geophysical feasibility) | Social–ecological feasibility  | Supporting references (Social–ecological feasibility)  | Feasibility (general) | Effectiveness  | Supporting references (Effectiveness)  |
|--|--|--|--|---|--|--|-----------------------|--|--|
| Institutional transboundary agreements     | Medium: Early planning and little implementation. Transboundary agreements can be constrained by economic negotiations over quotas and/or compensations. Technologically, they may require adaptive management or other systems that require a degree of technical capacity. | Pinsky et al. (2018); Melbourne-Thomas et al. (2021); Sunby et al. (2021)        | Low: Lack of an international regulatory framework or convention for climate-induced transboundary agreements. Existing regional fisheries management platforms that are transboundary, but countries have a strong role. Existing fishing agreements and policies are not considering climate-change impacts and may need to be redesigned. | Gaines et al. (2018); Engler (2020); Oremus et al. (2020)         | High: High feasibility in environmental and social terms as trade-offs are not expected if agreements address sustainability and equity issues. Risk of denying access to resources to communities, and of shifting pressure to other regions.   | Mason et al. (2020); Palacios-Abrantes et al. (2020); Melbourne-Thomas et al. (2021)           | Medium                | High: Institutional agreements and cooperation in sectors, such as fisheries, contributes to adaptation and offsetting climate-change impacts.   | Oremus et al. (2020); Haas et al. (2021)   |
| Accommodation and relocation               | Low: At the stage of adaptation planning and early implementation. Technology is advanced, but economic costs are high.  | Marzia et al. (2015); Hanson and Nicholls (2020); Monios and Willemsmeier (2020) | Medium: Diverse political support for this option, shift in international institutions towards nature-based solutions. Geophysically constrained by the environment and urbanisation of coastlines.  | Vilkolainen et al. (2017)   | Low: Low environmental feasibility due to irreversible environmental change. Medium social feasibility as sometimes these are socially accepted but other times can cause conflicts. Effects to communities outside of the immediate community need to be considered as well as local impacts. It is important that stakeholders be involved in decision making to ensure that impacts are understood and agreed upon. | Shelton (2014); Zickgraf (2021)  | Low                   | High: Relocation can be planned with participatory processes to achieve higher effectiveness, and may be required in the long run.   | Cross-Chapter Box SLR in Chapter 3; Magnan et al. (2020); Zickgraf (2021)                      |
| Protection and beach-and-shore nourishment | Medium: At the stage of adaptation planning and early implementation; high economic returns and technological ready; includes, however, large economic costs.  | Bayraktarov et al. (2016); Samora-Anvela et al. (2017); Pinto et al. (2018)      | High: High political support at the international level and increasing support nationally. Geophysical possible although limited by raw materials and footprint.   | Nicholls (2018)   | Medium: Low environmental feasibility as it can involve interventions that transform the natural habitats, although with the objective of maintaining ecosystem services. Medium social feasibility as these measures are generally accepted, although if not well designed, they can burden certain groups. Protection may be a feasible option for highly populated coastal areas.                                   | Gattuso et al. (2018); Barnard et al. (2021); de Schipper et al. (2021); Neijens et al. (2021) | Medium                | Medium: Protection and soft infrastructure is effective in the short term and required under highly developed coastlines, but is ineffective in the longer term as it is a barrier to Nbs. | Gattuso et al. (2018); Bevacqua et al. (2020); Kirezi et al. (2020); de Schipper et al. (2021) |

| Adaptation solution            | Technical and economic feasibility  | Supporting references (Technical and economic feasibility)                            | Institutional and geophysical feasibility  | Supporting references (Institutional and geophysical feasibility)              | Social–ecological feasibility  | Supporting references (Social–ecological feasibility)   | Feasibility (general) | Effectiveness   | Supporting references (Effectiveness)  |
|--------------------------------|---|---|--|--|--|---|-----------------------|---|--|
| Early warning systems          | High: Technology is mature and in widespread use. Economic barriers are linked to access to technology and information. Implementation continues to expand.   | Demele et al. (2019)  | High: High political acceptability and supporting institutions.  | Leal Filho et al. (2018)   | Medium: Early warning systems can aid decision making, diversification of enterprises and development of adaptable management systems for coastal systems and fisheries; however, it has implementation risks such as increasing inequalities due to access.               | Soares et al. (2018); Bindoff et al. (2019)   | High                  | High: Allows for a planned response to disasters and for policy integration (i.e., health, nutrition, etc.).                                  | Hoegh-Guldberg et al. (2014); Bindoff et al. (2019); Collins et al. (2019)                     |
| Seasonal and dynamic forecasts | Medium: Implementation is widespread, technologically ready, however, technology requires high capacity and economic costs are significant. Combining seasonal forecasting and climate projections in dynamic model forecasts might provide a pragmatic option for marine industries, including fisheries, aquaculture and tourism to adapt to future climate risk by changing practices or relocating. | Recha et al. (2015); Payne et al. (2017); Tommasi et al. (2017); Hobday et al. (2018) | High: High political acceptability and supporting institutions. Need to share information and data across jurisdictions.   | Hobday et al. (2016); A. Maureaud et al. (2021)                                | High: Rapid expansion of remote-sensing data, computational ability and ocean modelling have led to the proliferation of real-time and seasonal forecasts of marine heatwaves and associated impacts as well as the abundance and distribution of living marine resources. | Payne et al. (2017); Hobday et al. (2018)   | High                  | High: Improved technologies match management scales and can be incorporated into many adaptation solutions for different hazards and sectors. | Tommasi et al. (2017); Winter et al. (2020); Davidson et al. (2021); Spillman and Smith (2021) |
| Monitoring systems             | Medium: Technology is widely implemented and currently available and used. Economic quotas vary across regions where there are large gaps in ocean and coastal monitoring across the oceans.  | Kurekin et al. (2019); Claudet et al. (2020)  | Medium: Diverse political acceptability such that institutions differ at the national and local levels on the engagement of these technologies and investing in climate-change monitoring. There is uneven access and lack of shared information and open data in general. | Rilov et al. (2020); A. Maureaud et al. (2021); Melbourne-Thomas et al. (2021) | Medium: High feasibility for environmental monitoring in protected areas and management agencies. Medium social acceptance for social monitoring in sectors such as fishing. Benefits of monitoring vessel activity to avoid illegal harvesting.                           | Bel et al. (2013); Lubchenko and Grorud-Colvert (2015); Cabral et al. (2018); Kurekin et al. (2019) | Medium                | Medium: Monitoring is effective for climate-adaptive management; however, there are many barriers to implementation.                          | Rilov et al. (2020); A. Maureaud et al. (2021); Melbourne-Thomas et al. (2021)                 |

| Adaptation solution | Technical and economic feasibility  | Supporting references (Technical and economic feasibility)              | Institutional and geophysical feasibility  | Supporting references (Institutional and geophysical feasibility) | Social–ecological feasibility  | Supporting references (Social–ecological feasibility)   | Feasibility (general) | Effectiveness   | Supporting references (Effectiveness)  |
|---------------------|---|---|--|---|--|---|-----------------------|---|--|
| Habitat development | Low: Low stage of implementation with vulnerability assessment and early planning. High economic costs are related to technology, which is at the earlier stages.         | Gilby et al. (2018)   | Medium: Geographic feasibility may be moderately limited by marine spatial planning and uses; institutional feasibility has the regulations and support for these actions.   | Boerema and Meire (2017); von Haaren et al. (2019)                | Medium: Medium ecological and social feasibility as it has the potential to alleviate some of the risks posed by climate change. Possible risks in implementing assisted migration: invasive species, mortality and investment loss. Restoration activities may be ineffective or may lead to undesirable social impacts with endangered species or impacts from increased recreational use of restored habitat.   | Buckley and Crane (2008); Bindoff et al. (2019)   | Medium                | Medium to high: Has the potential to stabilise shorelines and increase fisheries productivity in rocky reef systems, but there is little evidence on effectiveness. | Gilby et al. (2018)  |
| Active restoration  | High: This restoration is expanding over marine systems and needs high investments, but it is economically productive. Technology is accessible and feasible as of today. | Bayraktarov et al. (2016); Basconi et al. (2020); Duarte et al. (2020a) | High: The geographic feasibility is high, does not require extensive areas limiting other uses and may reduce hazards. Regulations and policies are in place to regulate stressors and allow for broad restoration and conservation of ecosystems. | Larkin et al. (2019)  | Medium: High feasibility to recover ecosystem services provision that benefits livelihoods and their resilience. There can be conflicting uses, but these can be addressed with inclusive processes, and the benefits are supposed to outweigh the losses in most cases.   | Fadli et al. (2012); Gattuso et al. (2018); Bostrom-Einarsson et al. (2020); Hafezi et al. (2021) | High                  | High: Achieves species recovery and reintroduction.   | Bostrom-Einarsson et al. (2020); Rinkevich (2021)  |
| Assisted evolution  | Medium: Implementation is widespread; however, technical capacity can be a barrier and at the early stages; moderate to high cost.  | van Oppen et al. (2015)   | Medium: No significant barriers in geographical feasibility, as these interventions are small scale. Institutional feasibility is medium due to political acceptability and the lack of an existing regulatory and legal framework.                | Thiele (2020)   | Medium: Medium ecological and social feasibility as it has the potential to alleviate some of the risks posed by climate change. Translocated plants and animals may carry pathogens or parasites affecting the health of native populations, be maladapted to other non-climate-related changes or may cause a change in genetic composition or population structure of native organisms, a loss of genetic diversity or a breakdown of co-adapted gene complexes. A counter-argument here may be that with climate change and severe declines on the horizon, the spread and dominance of selected keystone species might be a better outcome than total loss. | Laikre et al. (2010); van Oppen et al. (2015); Anthony et al. (2017); Gattuso et al. (2018)       | Medium                | Medium: Potential to conserve species and habitats from future change.  | Bulleri et al. (2018); National Academies of Sciences (2019); Bostrom-Einarsson et al. (2020); Fredriksen et al. (2020); Morris et al. (2020); Kleypas et al. (2021) |

| Adaptation solution             | Technical and economic feasibility   | Supporting references (Technical and economic feasibility)  | Institutional and geophysical feasibility  | Supporting references (Institutional and geophysical feasibility)                   | Social–ecological feasibility   | Supporting references (Social–ecological feasibility)   | Feasibility (general) | Effectiveness   | Supporting references (Effectiveness)  |
|---------------------------------|--|---|--|---|---|---|-----------------------|---|--|
| Habitat restoration             | Medium: Adaptation planning and early to mid-implementation, high costs.   | Williams et al. (2015); Bayraktarov et al. (2016); Pinto et al. (2018); Duarte et al. (2020a)                   | Medium: Medium scope; not economies of scale, scalability limited and depends on habitats (i.e., mangroves more scalable than seagrasses); helps hazard-risk reduction (mitigation).   | Duarte et al. (2020a); Bertolini and da Motta (2021); Braun de Torrez et al. (2021) | High: Supports ecosystem services and biodiversity reducing their vulnerability to climate change; provides ecological resilience and social co-benefits. May have varying social acceptability given other habitat uses (fishing, infrastructure, etc.)  | Shelton (2014); Gattuso et al. (2018)   | Medium                | High: Has proven effective from local to regional to global scales; has associated co-benefits from mitigation to ecosystem services to coastal livelihoods.  | Silver et al. (2019); Duarte et al. (2020a); Gordon et al. (2020); Braun de Torrez et al. (2021) |
| MPAs and OECMs                  | High: MPAs already in place, widely implemented. There are additional costs of increasing enforcement, spill-over economic effects. OECMs have high technical feasibility as they only require recognition of an existing management system. | Takasaki (2016); Maxwell et al. (2020b); Rilov et al. (2020); Arafah-Dalmau et al. (2021); Guiméy et al. (2021) | High: High geophysical feasibility given the current network of MPAs and OECMs potential initiatives already in place, high institutional readiness given the regulatory framework of marine protection.   | Roberts et al. (2018); Tittensor et al. (2019)                                      | High: Support ecosystem services and biodiversity, although these can be impacted by climate change significantly. Provide ecological resilience. Some MPAs have associated social conflicts and acceptability issues, but these already exist. This can be minimised if inclusiveness and participatory processes are incorporated.  | Edgar et al. (2014); Gill et al. (2017); Wilson et al. (2020); Ovando et al. (2021); Sala et al. (2021) | High                  | Medium: As current MPAs are not expected to be effective against climate-change impacts in the middle term, where adaptive management, dynamic conservation and/or conservation of climate refugia may be needed. | Tittensor et al. (2019); Wilson et al. (2020)  |
| Conservation of climate refugia | Low: Adaptation planning stage, no implementation yet, extent of new MPAs unknown.   | Roberts et al. (2017); Rilov et al. (2020); Arafah-Dalmau et al. (2021)   | Low: Physically feasible and in a large-scale application but probably dynamic over time. Institutional bodies and agreements are not prepared for these dynamic conservation tools. Climate-change not yet incorporated in conservation planning. | Tittensor et al. (2019); Rilov et al. (2020); Arafah-Dalmau et al. (2021)           | High: Supports ecosystem services under climate change more efficiently and increases ecological and social resilience in the longer term. Limits social acceptability as access rights and marine uses can be limited. Loss of access to natural resources may result in burdens on communities and livelihood shifts, and may result in inequitable distribution of benefits. Design and implementation should be inclusive and participatory, combining participation with climate-change metrics. | Britto-Morales et al. (2018); Wilson et al. (2020); Arafah-Dalmau et al. (2021)                         | Medium                | High: Models and experiments show potential effectiveness of conserving climate refugia.  | Rilov et al. (2020); Wilson et al. (2020); Arafah-Dalmau et al. (2021)                           |

| Adaptation solution         | Technical and economic feasibility  | Supporting references (Technical and economic feasibility)                             | Institutional and geophysical feasibility   | Supporting references (Institutional and geophysical feasibility)     | Social–ecological feasibility  | Supporting references (Social–ecological feasibility)  | Feasibility (general) | Effectiveness   | Supporting references (Effectiveness)  |
|-----------------------------|---|--|---|---|--|--|-----------------------|---|--|
| Transboundary MSP and ICZM  | High: Large implementation processes of MSP and ICZM in most countries in the world; potential for transboundary MSP  | Fraião Santos et al. (2020); Li and Jay (2020)   | Medium: Coastal-use change and planning is feasible and broad in scope, but institutional needs for transboundary integrated coastal and ocean spatial management are at the infancy (i.e., ABNJ treaty). Additionally, existing MSP and ICZM lack considerations of climate-change adaptation. | Gissi et al. (2019); Fraião Santos et al. (2020)                      | High: Well-designed ICZM and MSP across jurisdictions can guarantee access rights and sustainable resource use that generates social and ecological co-benefits.   | Free et al. (2020)   | High                  | High: MSP that incorporates climate-change impacts and adaptation in the design can contribute to support climate adaptation from a multi-sector perspective and through existing policy frameworks. Little evidence exists for transboundary MSP and ICZM. | Tittensor et al. (2019); Fraião Santos et al. (2020); Rilov et al. (2020)            |
| Sustainable harvesting      | Medium: 30% of stocks not managed sustainably, problems of enforcement, access to assets, capacity shortfalls and perverse subsidies. High potential for sustainable mariculture. Tools known and implemented broadly, requires fleet adjustments in some cases and ending illegal fishing. Sustainable fisheries and mariculture stabilise income and create opportunities for value addition. | International Council for Science (2017); Khan et al. (2018a); Costello et al. (2020b) | High: Global institutions are ready and advocating for sustainable fisheries management, including regional fisheries management organizations and SDGs. National institutions vary in readiness, but overall global feasibility is high.   | Miller et al. (2010); Burden and Fujita (2019); Duarte et al. (2020a) | High: Increases provisioning ecosystem services in different systems and allows for social co-benefits if access is guaranteed and regulations are inclusive. Reduces the number of species at risk from climate change. Supports intergenerational equity and inclusive and participatory policies (i.e., co-management). | International Council for Science (2017); Le Blanc et al. (2017); Cheung et al. (2018); Allison (2011) | High                  | Medium: Sustainable harvesting effectively reduced the impacts of climate change; however, it may not be sufficient for specific systems (i.e., tropical coral reefs) and/or regions that expect large maximum-catch potential losses.                      | Gaines et al. (2018); Free et al. (2020); Lam et al. (2020); Sumaila and Tai (2020)  |
| Climate-adaptive management | Low: Requires data-rich assessments and ecological and climate monitoring. Adaptive management is effective, but with high costs for countries with scientific shortfalls. Very limited implementation so far.  | Huntington et al. (2017); Paulus et al. (2019); Holzman et al. (2020)                  | High: Scope still is small, but existing institutions can incorporate adaptive fisheries management without broad transformations; it is physically feasible and helps hazard-risk reduction  | Pentz et al. (2018)   | High: There are no significant environmental barriers, as it increases ecosystem services (food provisioning), SDG2. It can have social co-benefits or differ in acceptability. In order to increase acceptability, design should be inclusive and distribution across the globe.  | Pinsky et al. (2018)   | Medium                | High: Scarce examples of implementation, but models and evidence show high effectiveness to reduce risks and foster adaptation in fisheries and conservation.   | Asch (2015); Levin and Mellmann (2015); Fulton et al. (2019a); Nickols et al. (2019) |

| Adaptation solution        | Technical and economic feasibility   | Institutional and geophysical feasibility   | Supporting references (Institutional and geophysical feasibility) | Social–ecological feasibility  | Supporting references (Social–ecological feasibility) | Feasibility (general) | Effectiveness  | Supporting references (Effectiveness)                           |
|----------------------------|--|---|---|--|---|-----------------------|--|---|
| Ecosystem-based management | Medium: Implementation is expanding, requires economic resources but less than adaptive fisheries management; data-poor management is possible. Increasing implementation in fisheries management. | Medium: Scope so far is small but can be scaled up to regional management. Institutions are not ready, but it may not require important transformations. It contributes to risk reduction and resilience. | Wamsler et al. (2016); Bryndum-Buchholz et al. (2021)             | High: High ecological feasibility due to the co-benefits and the support to biodiversity and ecosystem services. High acceptability supports intergenerational equity and allows for participation, co-management and inclusive processes. | Alexander et al. (2019)                               | Medium                | High: Ecosystem-based management can incorporate many adaptation solutions, and can reduce climate impacts in fisheries in the near term and under low-emission scenarios. | Harvey et al. (2018); Karp et al. (2019); Holsman et al. (2020) |

At the global level, eliminating overexploitation *very likely* benefits fisheries adaptation to climate change (Burden and Fujita, 2019; Free et al., 2019; Sumaila and Tai, 2020). Regulating landing sizes is a simple strategy that addresses both climate-change and overexploitation impacts (Queirós et al., 2018; Thompson et al., 2020). Controlling overfishing may also decrease methylmercury bioaccumulation in pelagic Atlantic finfish (Schartup et al., 2019). But more sophisticated approaches like adaptive management, which anticipates and responds to changes in both fishery productivity and distribution, cannot only reduce fisheries impacts but also improve fisheries yields and profits (*high confidence*) (Costello et al., 2016; Gaines et al., 2018; Pinsky et al., 2018; IPBES, 2019; Karp et al., 2019). Management that adjusts the timing of fishery closures and uses timed stock-assessment surveys or hatchery releases might also effectively address climate-driven changes in phenology (Section 3.4.3.2) that alter the seasonality of harvests (*low evidence*) (Asch, 2015) and decrease the dependability of seasonal employment and cultural activities (Section 3.5.6). Achieving sustainable fishing practices together with strong mitigation (RCP2.6) is projected to reduce the number of fisheries target species at risk in 2100 by 63% (Cheung et al., 2018), but the implementation of climate-adaptive management in global fisheries remains limited (Holsman et al., 2020).

Ecosystem-based fisheries management is widely legislated (Bryndum-Buchholz et al., 2021) and can reduce climate impacts in fisheries in the near term, especially under low-emission scenarios (Karp et al., 2019; Holsman et al., 2020). Multi-species ecosystem-based management outperforms single-species management (Fulton et al., 2019a) and has been proposed as an effective tool for addressing regime shifts (Section 3.4.3.3; Levin and Möllmann, 2015). Ecosystem-based management, however, poses substantial challenges for nations that lack the necessary resources or information (Fernandino et al., 2018). Awareness and integration of social–ecological contexts in policy and management promotes equitable strategies for sustainable fisheries and mariculture, and for reducing risks of unintended negative social impacts on regions or sectors (Cochrane, 2021).

Transboundary agreements on shifting fisheries will reduce the risk of overharvesting and depletion of resources at the trailing edges of their distributions (*high agreement, medium confidence*) (Gaines et al., 2018). Permits tradable across political boundaries could also address this challenge, but *limited evidence* is available regarding their efficacy (Pinsky et al., 2018). One promising approach for migratory species is the ‘Vessel Day Scheme’, a cap-and-trade system (Agorau et al., 2018) that allows allocated fishing days to be exchanged among countries as El Niño shifts the tropical Pacific skipjack tuna distribution while preserving license revenue for all participating countries; however, this approach is not expected to fully accommodate projected tuna distributional shifts, requiring additional management adjustment (Bell et al., 2021). Other proposed adaptation strategies incorporate fisheries management and climate-smart conservation under the negotiations on areas beyond national jurisdiction (Pinsky et al., 2018; Tittensor et al., 2019; Frazão Santos et al., 2020), and in the Convention on Biological Diversity (CBD) areas designed as OECMs (Tittensor et al., 2019).

Despite the potential for adaptive management to achieve sustainable fisheries, outcomes will *very likely* be inequitable (Gaines et al., 2018;

Free et al., 2020; Lam et al., 2020). Many tropical and low-income countries will not be able to offset all climate-change impacts on fisheries or mariculture with management and policy reforms (*high confidence*) (Frazão Santos et al., 2020; Free et al., 2020; Bell et al., 2021), owing to the unequal geographic distribution of climate impacts and hazards (*high confidence*). In addition, fisheries reforms and adaptive management are less developed in the most climate-vulnerable and fisheries-dependent nations (*high confidence*) (Thiault et al., 2019; Lam et al., 2020; Bell et al., 2021). Human adaptations that reduce climate-driven risks to food provision can have positive or negative effects on marine resource management and socioeconomic systems (*limited evidence, high agreement*) (see also Section 3.6, Chapter 5 and detailed information in Barange et al., 2018). To overcome these limitations, community-level analyses that account for environmental and social conditions, as well as policy history, are required to support adaptation planning (*high confidence*) (Dubik et al., 2019; Rogers et al., 2019) so that climate risk of individual fishing communities and responses by local ecosystems and social systems can be considered in the context of neighbouring communities. Flexible and polycentric governance approaches have facilitated some short-term successes in achieving equitable, sustainable fisheries practices, but these may be challenging to implement where other governance systems, especially hierarchical systems, are well established (Cvitanovic et al., 2018; Bell et al., 2020).

### SM3.5.3 Multilateral Environmental Agreements and Climate Change

Several established global agreements and regional, sectoral or scientific bodies do already address climate adaptation and resilience, including the UN Framework Convention on Climate Change (UNFCCC), the UN Convention on Biological Diversity (CBD), Sustainable Development Goal 14 (Section 3.6.4), Regional Seas Conventions and Action Plans, and the Convention on Wetlands (Ramsar). All of them, except Ramsar, also address reducing other stressors, and regional fisheries management organizations and agreements (RFMOs/RFMAs), and the International Seabed Authority (ISA), seek to reduce non-climate stressors, also. UNFCCC and CBD also address finance mechanisms necessary for climate action (Conservation International and IUCN, 2021; Sumaila et al., 2021).

Reduction of non-climate stressors and adaptation to climate impacts has advanced slowly and unevenly under existing agreements and established bodies (*medium evidence, high agreement*). Parties to the UNFCCC's Paris Agreement have developed nationally determined contributions (NDCs) and adaptation communications detailing their plans to reduce greenhouse gas emissions and adapt to climate change; in 2017, only 9% of coastal nations' NDCs lacked marine considerations (Gallo et al., 2017). By 2020, more than 50 nations presented blue carbon strategies (Duarte et al., 2020a) intended to contribute to both mitigation and adaptation, and 29% of parties proposed restoring and protecting marine habitats for adaptation (Seddon et al., 2020). National adaptation communications include a diverse suite of products (Christiansen et al., 2020) whose ocean focus has not been assessed. Under the CBD, in 2011, nations pledged to protect 10% of the global ocean by 2020 (the Aichi targets) by achieving sustainable harvest and management of marine resources, alleviating other anthropogenic pressures on marine ecosystems, and expanding coverage and effectiveness of

MPAs and OECMs (CBD, 2020; CBD Secretariat, 2020). But as of August 2021, these targets were not met as only 7.74% of marine areas are protected (UNEP-WCMC, 2021). The 18 Regional Seas Conventions under the UN Environmental Programme, encompassing 146 nations, have advanced regional action on common marine environmental concerns, such as marine pollution, biodiversity, area-based measures, monitoring and climate-change adaptation (Johnson et al., 2021), all of which influence, and are influenced by, climate impacts. Since 1971, the Ramsar Convention has advanced coordinated action to monitor and sustainably manage wetlands, contributing to greater resilience, but this work has not been explicitly coupled to climate considerations (Finlayson et al., 2017). The RFMOs and other regional fisheries bodies facilitate international cooperation on high-seas fishing, but opinions vary on whether they have adequately prevented overfishing and marine ecosystem degradation (Lodge et al., 2007), two major drivers interacting with climate impacts on ocean and coastal systems (Sections 3.4–3.6). The ISA organises and controls mineral-resources-related activities in areas beyond national jurisdiction (International Seabed Authority, 2021), and although mineral extraction has not begun, vigorous debate exists about the potential short-term climate mitigation advances from accessing critical minerals to support sustainable technology versus potential long-term mitigation and adaptation impacts from influencing marine biodiversity and other deep-sea ecosystem functions (Koschinsky et al., 2018; Niner et al., 2018; Weaver et al., 2018; Carver et al., 2020; Kung et al., 2021; Tilot et al., 2021).

Many opportunities exist for policies being implemented or developed now to support climate adaptation and mitigation as well as plan for probable climate impacts. Some of these opportunities exist in the growing implementation of national and international marine spatial planning (especially in South American and African coastal countries) (Ehler and Fanny, 2009; Wright et al., 2019; Frazão Santos et al., 2020), and others are associated with the marine biological diversity beyond national jurisdictions (BBNJ) treaty being negotiated now by the UN, which will set targets for protection and sustainable use of the high seas (including the deep sea) and integrate planning, use and environmental-impact assessment of activities (Leary, 2019; Levin et al., 2020; Orejas et al., 2020). Opportunities also exist to update long-standing agreements, such as Regional Seas and RFMOs (Billé et al., 2017; Pentz et al., 2018; Johnson et al., 2021), and Ramsar (Hettiarachchi et al., 2015), with climate-informed targets and actions to achieve interacting objectives of climate mitigation and adaptation, reduction of non-climate impacts and protection of biodiversity.



## SM3.5.4 Data Supporting Figure 3.25

Table SM3.4 | Background materials and data for past implementation of marine nature-based solutions in Figure 3.25a

| Adapta-tion solution          | Final indicators                                 | Description   | Years     | Refer-ence  | Limitations   | Units           | 1970 | 1971       | 1972       | 1973       |
|-------------------------------|--|---|-----------|---|---|-----------------|------|------------|------------|------------|
| Habitat restoration           | Number of restoration projects: Salt marshes     | Number of restoration projects over time (per habitat type)                     | 1972–2018 | Duarte et al. (2020b)   | The global deployment of restoration projects for coastal ecosystems (kelps, coral reefs, mangroves, salt marshes, seagrass meadows and oyster reefs) were compiled for existing resources, including the published literature and databases, from which location, year of initiation, latitude and longitude of individual restoration sites were compiled. Where the year of initiation was not provided, it was estimated by subtracting 5 years from the year of publication, which was the mean time lag between project initiation and reporting. | n               | NA   | NA         | 1          | 1          |
|                               | Number of restoration projects: Seagrass meadows | Number of restoration projects over time (per habitat type)                     | 1972–2018 | Duarte et al. (2020b)   | The global deployment of restoration projects for coastal ecosystems (kelps, coral reefs, mangroves, salt marshes, seagrass meadows and oyster reefs) were compiled for existing resources, including the published literature and databases, from which location, year of initiation, latitude and longitude of individual restoration sites were compiled. Where the year of initiation was not provided, it was estimated by subtracting 5 years from the year of publication, which was the mean time lag between project initiation and reporting. | n               | NA   | NA         | 9          | 12         |
|                               | Number of restoration projects: Mangroves        | Number of restoration projects over time (per habitat type)                     | 1972–2018 | Duarte et al. (2020b)   | The global deployment of restoration projects for coastal ecosystems (kelps, coral reefs, mangroves, salt marshes, seagrass meadows and oyster reefs) were compiled for existing resources, including the published literature and databases, from which location, year of initiation, latitude and longitude of individual restoration sites were compiled. Where the year of initiation was not provided, it was estimated by subtracting 5 years from the year of publication, which was the mean time lag between project initiation and reporting. | n               | NA   | NA         | 13         | 13         |
|                               | Number of restoration projects: Oyster reef      | Number of restoration projects over time (per habitat type)                     | 1972–2018 | Duarte et al. (2020b)   | The global deployment of restoration projects for coastal ecosystems (kelps, coral reefs, mangroves, salt marshes, seagrass meadows and oyster reefs) were compiled for existing resources, including the published literature and databases, from which location, year of initiation, latitude and longitude of individual restoration sites were compiled. Where the year of initiation was not provided, it was estimated by subtracting 5 years from the year of publication, which was the mean time lag between project initiation and reporting. | n               | NA   | NA         | 2          | 2          |
|                               | Number of restoration projects: Coral reef       | Number of restoration projects over time (per habitat type)                     | 1972–2018 | Duarte et al. (2020b)   | The global deployment of restoration projects for coastal ecosystems (kelps, coral reefs, mangroves, salt marshes, seagrass meadows and oyster reefs) were compiled for existing resources, including the published literature and databases, from which location, year of initiation, latitude and longitude of individual restoration sites were compiled. Where the year of initiation was not provided, it was estimated by subtracting 5 years from the year of publication, which was the mean time lag between project initiation and reporting. | n               | NA   | NA         | 0          | 0          |
| Current MPAs                  | Coastal and marine MPAs (km <sup>2</sup> )       | MPA coverage (in km <sup>2</sup> over time, per type of MPA: total and no take) | 1971–2020 | Data on location, size and year of declaration of MPAs from UNEP-WCMC and IUCN (2020)                     | This dataset has been filtered to only include coastal and marine ecosystems  | km <sup>2</sup> | NA   | 224960.338 | 551639.784 | 663301.317 |
|                               | No-take MPAs                                     | MPA coverage (in km <sup>2</sup> over time, per type of MPA: total and no take) | 1971–2020 | Data on location, size and year of declaration of MPAs from UNEP-WCMC and IUCN (2020)                     | This dataset has been filtered to only include coastal and marine ecosystems  | km <sup>2</sup> | NA   | 636.911852 | 647.541472 | 686.337148 |
| Transboundary MSP and ICM     | High-sea MPAs (km <sup>2</sup> )                 | High-seas MPAs (coverage in km <sup>2</sup> over time)                          | 2006–2018 | Data on location, size and year of declaration of high-sea MPAs (i.e., ABNJ) from OSPAR Commission (2019) | Older data not available in sources.  | km <sup>2</sup> | NA   | NA         | NA         | NA         |
| sustainable fisheries harvest | Rebuilt fish stocks (%)                          | Overexploited and collapsed fish stocks in the process of rebuilding            | 1971–2014 | Kleisner et al. (2013); Gattuso et al. (2018)   | Relative to existing fisheries assessments and statistics   | %               | NA   | 0.07       | 0.07       | 0.14       |

| 1974       | 1975       | 1976       | 1977       | 1978       | 1979       | 1980       | 1981       | 1982       | 1983       | 1984       | 1985       | 1986       | 1987      | 1988       | 1989       |
|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-----------|------------|------------|
| 3          | 5          | 6          | 6          | 8          | 9          | 11         | 15         | 15         | 18         | 21         | 24         | 27         | 29        | 32         | 33         |
| 13         | 14         | 15         | 17         | 18         | 19         | 22         | 24         | 27         | 28         | 32         | 34         | 36         | 46        | 52         | 58         |
| 13         | 13         | 15         | 16         | 16         | 18         | 24         | 26         | 28         | 29         | 33         | 36         | 44         | 48        | 51         | 57         |
| 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 3         | 3          | 3          |
| 0          | 0          | 0          | 0          | 0          | 0          | 0          | 0          | 1          | 1          | 1          | 1          | 1          | 3         | 3          | 3          |
| 672626.794 | 683341.533 | 793865.791 | 1802607.41 | 1986538.67 | 1994272.78 | 2141194.34 | 2516441.45 | 2554939.12 | 2572375.06 | 2747849.85 | 2777075.94 | 2793954.65 | 2823355.9 | 2869392.06 | 2894654.26 |
| 690.224648 | 722.747882 | 1359.39753 | 1410.67753 | 1542.13753 | 1914.67085 | 1919.61947 | 4556.83278 | 5307.07417 | 5573.23417 | 5575.53417 | 23070      | 5803.17723 | 5839.9455 | 5862.21423 | 5866.01055 |
| NA         | NA        | NA         | NA         |
| 0.21       | 0.07       | 0.27       | 0.55       | 0.47       | 0.61       | 0.54       | 0.54       | 0.54       | 0.67       | 0.87       | 0.74       | 0.93       | 1.06      | 1.26       | 1.33       |

**Table SM3.5 |** Background materials and data for coral reef adaptation pathways in Figure 3.25b, based on assessment of effectiveness at maintaining >20% coral cover

| Coral reefs          | 1°     | 1–1.5°                 | 1.5–2°                 | 2–2.5°                 | 2.5–3°                 | 3–4°                 | Confidence level | Rationale   | References   |
|----------------------|--------|------------------------|------------------------|------------------------|------------------------|----------------------|------------------|---|--|
| Best management      | Starts | Ongoing <sup>a</sup>   | Uncertain <sup>c</sup> |                        |                        |                      | Very high        | Consensus of model results for reduction of grazing fish pressure (Caribbean) and crown-of-thorns starfish removal (Australia)  | Anthony et al. (2019); National Academies of Sciences Engineering and Medicine (2019)                  |
| Coral restoration    | Starts | Uncertain <sup>b</sup> | Uncertain <sup>c</sup> |                        |                        |                      | High             | Restoration without coral enhancement (e.g., assisted evolution, assisted gene flow) ineffective beyond 1.5°C warming, following SR1.5, SROCC and Section 3.4.2.2   | Section 3.4.2.1 (drawn from text on restoration)   |
| Enhanced coral       |        | Starts                 | Ongoing <sup>a</sup>   | Uncertain <sup>b</sup> | Uncertain <sup>c</sup> |                      | Medium           | Consensus of model results combining best management and introduction of more heat-tolerant corals 'outplanting warm-adapted coral juveniles' in Anthony et al. (2019)  | Anthony et al. (2019); National Academies of Sciences Engineering and Medicine (2019)                  |
| Reef shading         |        | Starts                 | Ongoing <sup>a</sup>   | Uncertain <sup>b</sup> | Uncertain <sup>c</sup> |                      | Medium           | Consensus of model results combining best management, introduction of more heat-tolerant corals and artificial reef shading   | Anthony et al. (2019); National Academies of Sciences Engineering and Medicine (2019)                  |
| Changing livelihoods |        | Starts                 | Ongoing <sup>a</sup>   | Ongoing <sup>a</sup>   | Ongoing <sup>a</sup>   | Ongoing <sup>a</sup> | Very high        | Drawn from <i>very high confidence</i> in reef degradation with >1.5°C warming and no management; supported consensus of the two model results with no management applied, and by other available projection studies which do not simulate management (see Section 3.4.2.1, no evolution case) (e.g., Logan et al., 2021) | Section 3.4.2.1; Anthony et al. (2019); National Academies of Sciences Engineering and Medicine (2019) |

Notes:

'Best management' refers to reducing fishing pressure and elimination of non-climate stressors (e.g., removal of crown-of-thorns starfish). 'Coral restoration' refers to restoring damaged reefs via methods such as transplanting corals grown in nurseries. Enhanced interventions that would increase coral thermal tolerance (e.g., assisted evolution or assisted gene flow). 'Reef shading' refers to efforts to decrease incident radiation. 'Changing livelihoods' refers to reduction in community reliance on coral reef services.

(a) *Likely* (>66%)(b) *More likely than not to likely* (50–66%)(c) *Unlikely to more likely than not* (33–50%)**Table SM3.6 |** Background materials and data for mangrove adaptation pathways in Figure 3.25c

| Mangrove                 | 1°     | 1–1.5°  | 1.5–2°  | 2–2.5°    | 2.5–3°    | 3–4°      | Confidence level | Rationale  | References                                  |
|--------------------------|--------|---------|---------|-----------|-----------|-----------|------------------|--|---|
| Restoration/revegetation | Starts | Ongoing | Ongoing | Ongoing   | Uncertain | Uncertain | Very high        | Mangroves are resistant to increase in temperature, though it is uncertain how much temperature may affect the recruitment of new plants and the availability of suitable space.   | Duarte et al. (2020a); Friess et al. (2020) |
| Conservation             | Starts | Ongoing | Ongoing | Uncertain |           |           | High             | Without interventions that allow mangroves to migrate landward, mangroves will likely suffer significant losses by mid-century, even under SSP1-2.6. By the end of the century, the risk of severe mangrove losses resulting from SLR ranges from <i>very likely</i> under SSP1-2.6 to <i>extremely likely</i> under SSP5-8.5. | Section 3.4.2.5; Duarte et al. (2020a)      |

| Mangrove                                    | 1° | 1–1.5° | 1.5–2°  | 2–2.5°  | 2.5–3°    | 3–4°      | Confidence level | Rationale  | References   |
|---|----|--------|---------|---------|-----------|-----------|------------------|--|--|
| Payment for ecosystem services and C market |    | Starts | Ongoing | Ongoing | Ongoing   | Uncertain | High             | The C market and PES strategies will depend on the success of other interventions to provide C stock in the long term. | Macreadie et al. (2019)                                |
| Diversify livelihoods                       |    | Starts | Ongoing | Ongoing | Uncertain | Uncertain | Medium           |  | Duarte et al. (2020a); Stewart-Sinclair et al. (2020b) |
| Migration and relocation (people)           |    | Starts | Ongoing | Ongoing | Uncertain |           | Low              |  | Duarte et al. (2020a); Lovelock and Reef (2020)        |
| Hard infrastructure/retreat                 |    | Starts | Ongoing | Ongoing | Ongoing   | Uncertain | Low              |  | Duarte et al. (2020a); Lovelock and Reef 2020)         |

### SM3.5.5 Data Supporting Figure 3.26

Table SM3.7 | Background materials and data for Figure 3.26

| Group   | From                                  | To                                | Inter-action | Agree-ment | Author scores (-3 to 3) |   |   |   |   | References  |
|---------|---------------------------------------|-----------------------------------|--------------|------------|-------------------------|---|---|---|---|---|
|         | SDG13                                 | 14.1 – Reduce Pollution           | 2            | 100        | 2                       | 2 | 2 | 2 | 2 | International Council for Science (2017); Le Blanc et al. (2017); Nilsson et al. (2018)                       |
|         | SDG13                                 | 14.2 – Protection and Restoration | 3            | 100        | 3                       | 3 | 3 | 3 | 3 | Hoegh-Guldberg and Bruno (2010); Le Blanc et al. (2017); Pecl et al. (2017)                                   |
|         | SDG13                                 | 14.3 – Reduce OA                  | 3            | 100        | 3                       | 3 | 3 | 3 | 3 | Hoegh-Guldberg and Bruno (2010); Le Blanc et al. (2017); Nilsson et al. (2018)                                |
|         | SDG13                                 | 14.4 – Sustainable Fishing        | 2            | 80         | 3                       | 2 | 2 | 2 | 2 | Le Blanc et al. (2017); Pecl et al. (2017)  |
|         | SDG13                                 | 14.5 – Conservation               | 2            | 100        | 2                       | 2 | 2 | 2 | 2 | Le Blanc et al. (2017)  |
|         | SDG13                                 | 14.6 – No Overfishing Subsidies   | 0            | 80         | 0                       | 1 | 0 | 0 | 0 | Sumaila et al. (2010); Pinsky et al. (2018)   |
|         | SDG13                                 | 14.7 – Sustainable Resources      | 3            | 100        | 3                       | 3 | 3 | 3 | 3 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Wabnitz et al. (2018) |
|         | SDG13                                 | 14.A – Knowledge                  | 3            | 80         | 2                       | 3 | 3 | 3 | 3 | International Council for Science (2017); Pecl et al. (2017)  |
|         | SDG13                                 | 14.B – Small-Scale Fisheries      | 1            | 60         | 1                       | 1 | 1 | 0 | 2 | FAO (2015b)   |
|         | SDG13                                 | 14.C – Sea Law                    | 3            | 100        | 3                       | 3 | 3 | 3 | 3 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018)   |
| Society | 14.1 – Reduce Marine Pollution        | SDG1 – No Poverty                 | 2            | 80         | 2                       | 1 | 2 | 2 | 2 | Shahidul Islam and Tanaka (2004); International Council for Science (2017); Le Blanc et al. (2017)            |
| Society | 14.2 – Protect and Restore Ecosystems | SDG1 – No Poverty                 | 2            | 60         | 2                       | 2 | 2 | 3 | 1 | International Council for Science (2017); Le Blanc et al. (2017)  |
| Society | 14.3 – Reduce Ocean Acidification     | SDG1 – No Poverty                 | 2            | 60         | 1                       | 3 | 1 | 2 | 1 | International Council for Science (2017)  |
| Society | 14.4 – Sustainable Fishing            | SDG1 – No Poverty                 | 2            | 60         | 2                       | 2 | 2 | 3 | 2 | Allison (2011); International Council for Science (2017); Le Blanc et al. (2017)                              |

| Group   | From   | To                 | Inter-action | Agree-ment | Author scores<br>(-3 to 3) |    |    |   |   | References  |
|---------|--|--------------------|--------------|------------|----------------------------|----|----|---|---|---|
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG1 – No Poverty  | 1            | 40         | -1                         | 1  | 1  | 3 | 2 | International Council for Science (2017); Le Blanc et al. (2017); Singh et al. (2018)                       |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG1 – No Poverty  | 0            | 60         | -1                         | -1 | -1 | 2 | 2 | Allison (2011); Le Blanc et al. (2017)  |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG1 – No Poverty  | 3            | 60         | 3                          | 2  | 2  | 3 | 2 | International Council for Science (2017); Le Blanc et al. (2017); Singh et al. (2018)                       |
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG1 – No Poverty  | 2            | 40         | 1                          | 0  | 2  | 2 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.B – Support Small-Scale Fisheries   | SDG1 – No Poverty  | 3            | 100        | 3                          | 3  | 3  | 3 | 2 | Le Blanc et al. (2017)  |
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG1 – No Poverty  | 1            | 40         | 1                          | 0  | 2  | 1 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.1 – Reduce Marine Pollution   | SDG2 – Zero Hunger | 2            | 80         | 3                          | 1  | 3  | 1 | 3 | Shahidul Islam and Tanaka (2004)  |
| Society | 14.2 – Protect and Restore Ecosystems  | SDG2 – Zero Hunger | 2            | 60         | 2                          | 2  | 2  | 2 | 2 | Kawarazuka and Béné (2011); Béné et al. (2016); International Council for Science (2017)                    |
| Society | 14.3 – Reduce Ocean Acidification  | SDG2 – Zero Hunger | 2            | 60         | 1                          | 1  | 2  | 3 | 1 | Hoegh-Guldberg et al. (2017); Le Blanc et al. (2017)  |
| Society | 14.4 – Sustainable Fishing   | SDG2 – Zero Hunger | 2            | 60         | 3                          | 1  | 2  | 3 | 2 | Allison (2011); International Council for Science (2017); Le Blanc et al. (2017)                            |
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG2 – Zero Hunger | 1            | 40         | 1                          | -1 | 1  | 2 | 1 | International Council for Science (2017); Le Blanc et al. (2017); Singh et al. (2018)                       |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG2 – Zero Hunger | 0            | 60         | -1                         | -1 | 1  | 0 | 1 | Allison (2011); Le Blanc et al. (2017)  |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG2 – Zero Hunger | 2            | 60         | 3                          | 1  | 2  | 3 | 1 | Béné et al. (2016); FAO (2016); Le Blanc et al. (2017)  |
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG2 – Zero Hunger | 2            | 40         | 1                          | 2  | 2  | 2 | 1 | Le Blanc et al. (2017)  |
| Society | 14.B – Support Small-Scale Fisheries   | SDG2 – Zero Hunger | 3            | 100        | 2                          | 3  | 3  | 3 | 2 | Le Blanc et al. (2017)  |
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG2 – Zero Hunger | 1            | 40         | 1                          | 0  | 2  | 1 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |

| Group   | From   | To                                | Inter-action | Agreement | Author scores<br>(-3 to 3) |   |   |   |   | References  |
|---------|--|-----------------------------------|--------------|-----------|----------------------------|---|---|---|---|---|
| Society | 14.1 – Reduce Marine Pollution   | SDG3 – Good Health and Well-being | 3            | 80        | 3                          | 3 | 3 | 2 | 3 | Le Blanc et al. (2017)  |
| Society | 14.2 – Protect and Restore Ecosystems  | SDG3 – Good Health and Well-being | 2            | 40        | 1                          | 1 | 2 | 2 | 3 | HLPE (2014); IPCC (2014a); IPCC (2014b); Béné et al. (2016)   |
| Society | 14.3 – Reduce Ocean Acidification  | SDG3 – Good Health and Well-being | 1            | 80        | 1                          | 1 | 2 | 1 | 1 | Vezzulli et al. (2012)  |
| Society | 14.4 – Sustainable Fishing   | SDG3 – Good Health and Well-being | 2            | 40        | 1                          | 2 | 2 | 3 | 3 | Le Blanc et al. (2017)  |
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG3 – Good Health and Well-being | 1            | 40        | 0                          | 0 | 1 | 2 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG3 – Good Health and Well-being | 0            | 80        | 0                          | 0 | 0 | 0 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG3 – Good Health and Well-being | 2            | 60        | 1                          | 1 | 2 | 3 | 1 | Perry (2010)  |
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG3 – Good Health and Well-being | 2            | 60        | 1                          | 2 | 1 | 2 | 2 | Calcabrini et al. (2017); Le Blanc et al. (2017)  |
| Society | 14.B – Support Small-Scale Fisheries   | SDG3 – Good Health and Well-being | 2            | 40        | 1                          | 1 | 2 | 2 | 3 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG3 – Good Health and Well-being | 2            | 60        | 1                          | 1 | 2 | 1 | 3 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.1 – Reduce Marine Pollution   | SDG4 – Quality Education          | 0            | 80        | 0                          | 0 | 1 | 0 | 0 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.2 – Protect and Restore Ecosystems  | SDG4 – Quality Education          | 0            | 60        | 0                          | 0 | 1 | 0 | 1 | Vladimirova and Le Blanc (2016)   |
| Society | 14.3 – Reduce Ocean Acidification  | SDG4 – Quality Education          | 0            | 80        | 0                          | 0 | 1 | 0 | 0 | NA <sup>a</sup>   |
| Society | 14.4 – Sustainable Fishing   | SDG4 – Quality Education          | 1            | 60        | 0                          | 1 | 2 | 0 | 0 | NA <sup>a</sup>   |
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG4 – Quality Education          | 0            | 100       | 0                          | 0 | 0 | 0 | 0 | NA <sup>a</sup>   |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG4 – Quality Education          | 0            | 80        | 0                          | 1 | 0 | 0 | 0 | NA <sup>a</sup>   |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG4 – Quality Education          | 1            | 80        | 1                          | 1 | 1 | 1 | 0 | NA <sup>a</sup>   |

| Group   | From   | To                                | Inter-action | Agree-ment | Author scores<br>(-3 to 3) |   |   |   |   | References  |
|---------|--|-----------------------------------|--------------|------------|----------------------------|---|---|---|---|---|
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG4 – Quality Education          | 1            | 80         | 1                          | 1 | 1 | 1 | 0 | NA <sup>a</sup>   |
| Society | 14.B – Support Small-Scale Fisheries   | SDG4 – Quality Education          | 1            | 60         | 0                          | 0 | 2 | 0 | 1 | NA <sup>a</sup>   |
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG4 – Quality Education          | 0            | 60         | 0                          |   | 1 | 0 | 0 | NA <sup>a</sup>   |
| Society | 14.1 – Reduce Marine Pollution   | SDG5 – Gender Equality            | 2            | 60         | 1                          | 2 | 1 | 2 | 2 | Harper et al. (2013); Bellante et al. (2016)  |
| Society | 14.2 – Protect and Restore Ecosystems  | SDG5 – Gender Equality            | 1            | 40         | 0                          | 0 | 1 | 3 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.3 – Reduce Ocean Acidification  | SDG5 – Gender Equality            | 1            | 40         | 1                          | 0 | 1 | 3 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.4 – Sustainable Fishing   | SDG5 – Gender Equality            | 1            | 60         | 1                          | 0 | 1 | 3 | 1 | Allison (2011)  |
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG5 – Gender Equality            | 1            | 40         | 0                          | 0 | 1 | 3 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG5 – Gender Equality            | 1            | 60         | 0                          | 0 | 0 | 2 | 1 | Schuhbauer et al. (2017)  |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG5 – Gender Equality            | 1            | 80         | 1                          | 1 | 1 | 3 | 1 | Le Blanc et al. (2017)  |
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG5 – Gender Equality            | 2            | 60         | 1                          | 1 | 1 | 3 | 2 | Le Blanc et al. (2017)  |
| Society | 14.B – Support Small-Scale Fisheries   | SDG5 – Gender Equality            | 2            | 40         | 1                          | 3 | 1 | 3 | 2 | Harper et al. (2013); Le Blanc et al. (2017)  |
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG5 – Gender Equality            | 1            | 40         | 1                          | 0 | 1 | 3 | 1 | McLeod et al. (2018); Michalena et al. (2020)   |
| Society | 14.1 – Reduce Marine Pollution   | SDG6 – Clean Water and Sanitation | 3            | 80         | 3                          | 3 | 3 | 3 | 2 | Ferrario et al. (2014)  |
| Society | 14.2 – Protect and Restore Ecosystems  | SDG6 – Clean Water and Sanitation | 2            | 60         | 2                          | 2 | 2 | 3 | 1 | Luh et al. (2017); Pecl et al. (2017)   |
| Society | 14.3 – Reduce Ocean Acidification  | SDG6 – Clean Water and Sanitation | 1            | 40         | 1                          | 0 | 1 | 2 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.4 – Sustainable Fishing   | SDG6 – Clean Water and Sanitation | 1            | 60         | 0                          | 0 | 0 | 2 | 1 | Hassan et al. (2008)  |

| Group   | From   | To                                 | Inter-action | Agreement | Author scores<br>(-3 to 3) |    |    |   |    | References  |
|---------|--|------------------------------------|--------------|-----------|----------------------------|----|----|---|----|---|
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG6 – Clean Water and Sanitation  | 2            | 60        | 1                          | 2  | 2  | 2 | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG6 – Clean Water and Sanitation  | 0            | 80        | 0                          | 0  | 0  | 0 | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG6 – Clean Water and Sanitation  | 1            | 60        | 1                          | -1 | 1  | 3 | 1  | Holding et al. (2016); United Nations (2019)  |
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG6 – Clean Water and Sanitation  | 1            | 60        | 1                          | 0  | 1  | 2 | 1  | Racault et al. (2019)   |
| Society | 14.B – Support Small-Scale Fisheries   | SDG6 – Clean Water and Sanitation  | 1            | 60        | 0                          | 0  | 1  | 1 | 1  | Rangel Soares et al. (2002); FAO (2009)   |
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG6 – Clean Water and Sanitation  | 1            | 80        | 1                          | 0  | 1  | 1 | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.1 – Reduce Marine Pollution   | SDG7 – Affordable and Clean Energy | 2            | 40        | 3                          | 2  | 2  | 3 | -1 | European Commission (2012); Coping et al. (2014); Ellabban et al. (2014); Rilov et al. (2020)               |
| Society | 14.2 – Protect and Restore Ecosystems  | SDG7 – Affordable and Clean Energy | 0            | 40        | 0                          | -2 | 2  | 2 | -1 | Wright (2015); Fuso Nerini et al. (2018)  |
| Society | 14.3 – Reduce Ocean Acidification  | SDG7 – Affordable and Clean Energy | 0            | 80        | 0                          | 0  | 1  | 0 | 0  | NA <sup>a</sup>   |
| Society | 14.4 – Sustainable Fishing   | SDG7 – Affordable and Clean Energy | -1           | 60        | 0                          | -2 | 0  | 0 | -1 | Copping et al. (2014)   |
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG7 – Affordable and Clean Energy | -1           | 40        | 0                          | -2 | -1 |   | -1 | Wright (2015); Rilov et al. (2020)  |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG7 – Affordable and Clean Energy | 0            | 100       | 0                          | 0  | 0  | 0 | 0  | NA <sup>a</sup>   |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG7 – Affordable and Clean Energy | 2            | 60        | 1                          | 2  | 2  | 2 | 1  | Blechinger et al. (2016)  |
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG7 – Affordable and Clean Energy | 3            | 60        | 3                          | 2  | 3  | 3 | 2  | Gegg and Wells (2019)   |
| Society | 14.B – Support Small-Scale Fisheries   | SDG7 – Affordable and Clean Energy | 0            | 100       | 0                          | 0  | 0  | 0 | 0  | NA <sup>a</sup>   |

| Group   | From   | To   | Inter-action | Agree-ment | Author scores<br>(-3 to 3) |    |   |    |   | References  |
|---------|--|--|--------------|------------|----------------------------|----|---|----|---|---|
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG7 – Affordable and Clean Energy             | 1            | 60         | 2                          | 1  | 1 | 2  | 1 | Wright (2014)   |
| Economy | 14.1 – Reduce Marine Pollution   | SDG8 – Decent Work and Economic Growth         | 1            | 40         | 1                          | -1 | 2 | 2  | 1 | Jang et al. (2014); International Council for Science (2017); Krelling et al. (2017)                        |
| Economy | 14.2 – Protect and Restore Ecosystems  | SDG8 – Decent Work and Economic Growth         | 1            | 40         | -1                         | -1 | 3 | 1  | 1 | Gillett and Lightfoot (2001); Alison (2011); Béné et al. (2016); International Council for Science (2017)   |
| Economy | 14.3 – Reduce Ocean Acidification  | SDG8 – Decent Work and Economic Growth         | 1            | 80         | 1                          | 0  | 1 | 1  | 1 | Gillett and Lightfoot (2001); Alison (2011); Béné et al. (2016); International Council for Science (2017)   |
| Economy | 14.4 – Sustainable Fishing   | SDG8 – Decent Work and Economic Growth         | 2            | 80         | 2                          | -1 | 2 | 2  | 2 | Allison (2011); International Council for Science (2017); Le Blanc et al. (2017)                            |
| Economy | 14.5 – Conserve Coastal and Marine Areas                                       | SDG8 – Decent Work and Economic Growth         | 1            | 60         | -1                         | 1  | 1 | 2  | 1 | Russi et al. (2016); International Council for Science (2017); Le Blanc et al. (2017)                       |
| Economy | 14.6 – End Subsidies Contributing to Overfishing                               | SDG8 – Decent Work and Economic Growth         | 0            | 40         | -1                         | -1 | 0 | 0  | 1 | Le Blanc et al. (2017)  |
| Economy | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG8 – Decent Work and Economic Growth         | 3            | 60         | 3                          | 2  |   | 3  | 3 | International Council for Science (2017); Le Blanc et al. (2017)  |
| Economy | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG8 – Decent Work and Economic Growth         | 2            | 40         | 1                          | 3  | 2 | 3  | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.B – Support Small-Scale Fisheries   | SDG8 – Decent Work and Economic Growth         | 2            | 40         | 2                          | 1  | 3 | 2  | 3 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.C – Implement and Enforce International Sea Law                             | SDG8 – Decent Work and Economic Growth         | 1            | 40         | 1                          | 0  | 3 | 1  | 2 | Russi et al. (2016)   |
| Economy | 14.1 – Reduce Marine Pollution   | SDG9 – Industry, Innovation and Infrastructure | 0            | 40         | 0                          | -1 | 2 | -1 | 1 | Anderson et al. (2017)  |
| Economy | 14.2 – Protect and Restore Ecosystems  | SDG9 – Industry, Innovation and Infrastructure | 0            | 60         | 0                          | 0  | 2 | -1 | 0 | NA <sup>a</sup>   |
| Economy | 14.3 – Reduce Ocean Acidification  | SDG9 – Industry, Innovation and Infrastructure | 0            | 80         | 0                          | 0  | 1 | 0  | 0 | NA <sup>a</sup>   |
| Economy | 14.4 – Sustainable Fishing   | SDG9 – Industry, Innovation and Infrastructure | 1            | 60         | 0                          | 1  | 2 | 0  | 0 | NA <sup>a</sup>   |

| Group   | From   | To   | Inter-action | Agreement | Author scores<br>(-3 to 3) |    |   |    |   | References  |
|---------|--|--|--------------|-----------|----------------------------|----|---|----|---|---|
| Economy | 14.5 – Conserve Coastal and Marine Areas                                       | SDG9 – Industry, Innovation and Infrastructure | 0            | 60        | 0                          | 0  | 1 | -1 | 0 | NA <sup>a</sup>   |
| Economy | 14.6 – End Subsidies Contributing to Overfishing                               | SDG9 – Industry, Innovation and Infrastructure | 0            | 40        | 0                          | 1  | 1 | -1 | 0 | NA <sup>a</sup>   |
| Economy | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG9 – Industry, Innovation and Infrastructure | 1            | 60        | 0                          | 0  | 2 | 0  | 1 | Blechinger et al. (2016)  |
| Economy | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG9 – Industry, Innovation and Infrastructure | 2            | 60        | 0                          | 2  | 2 | 3  | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.B – Support Small-Scale Fisheries   | SDG9 – Industry, Innovation and Infrastructure | 1            | 40        | 0                          | 0  | 2 | 3  | 2 | NA <sup>a</sup>   |
| Economy | 14.C – Implement and Enforce International Sea Law                             | SDG9 – Industry, Innovation and Infrastructure | 0            | 40        | 0                          | 0  | 2 | -1 | 1 | NA <sup>a</sup>   |
| Economy | 14.1 – Reduce Marine Pollution   | SDG10 – Reduced Inequalities                   | 1            | 40        | 1                          | 0  | 2 | 0  | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.2 – Protect and Restore Ecosystems  | SDG10 – Reduced Inequalities                   | 1            | 60        | 1                          | 0  | 2 | 1  | 1 | Beck et al. (2018); Naidoo et al. (2019)  |
| Economy | 14.3 – Reduce Ocean Acidification  | SDG10 – Reduced Inequalities                   | 1            | 80        | 1                          | 1  | 1 | 0  | 1 | White et al. (2000)   |
| Economy | 14.4 – Sustainable Fishing   | SDG10 – Reduced Inequalities                   | 1            | 40        | 2                          | 1  | 2 | 0  | 1 | Allison (2011)  |
| Economy | 14.5 – Conserve Coastal and Marine Areas                                       | SDG10 – Reduced Inequalities                   | 0            | 60        | -1                         | -1 | 1 | -1 | 1 | Le Blanc et al. (2017); Singh et al. (2018)   |
| Economy | 14.6 – End Subsidies Contributing to Overfishing                               | SDG10 – Reduced Inequalities                   | 1            | 60        | 1                          | 2  | 1 | 2  | 1 | Le Blanc et al. (2017); Schuhbauer et al. (2017)  |
| Economy | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG10 – Reduced Inequalities                   | 3            | 100       | 3                          | 3  | 3 | 3  | 3 | Le Blanc et al. (2017)  |
| Economy | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG10 – Reduced Inequalities                   | 1            | 80        | 1                          | 1  | 2 | 1  | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.B – Support Small-Scale Fisheries   | SDG10 – Reduced Inequalities                   | 2            | 60        | 2                          | 2  | 3 | 2  | 3 | Le Blanc et al. (2017)  |

| Group   | From   | To   | Inter-action | Agree-ment | Author scores<br>(-3 to 3) |   |   |   |   | References  |
|---------|--|--|--------------|------------|----------------------------|---|---|---|---|---|
| Economy | 14.C – Implement and Enforce International Sea Law                             | SDG10 – Reduced Inequalities                   | 1            | 80         | 1                          | 1 | 2 | 1 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.1 – Reduce Marine Pollution   | SDG11 – Sustainable Cities and Communities     | 2            | 60         | 1                          | 3 | 2 | 2 | 2 | International Council for Science (2017)  |
| Society | 14.2 – Protect and Restore Ecosystems  | SDG11 – Sustainable Cities and Communities     | 2            | 40         | 1                          | 1 | 2 | 3 | 2 | Marzeion and Levermann (2014); International Council for Science (2017); Reimann et al. (2018)              |
| Society | 14.3 – Reduce Ocean Acidification  | SDG11 – Sustainable Cities and Communities     | 1            | 60         | 1                          | 2 | 1 | 2 | 1 | Heron et al. (2017); International Council for Science (2017)   |
| Society | 14.4 – Sustainable Fishing   | SDG11 – Sustainable Cities and Communities     | 1            | 60         | 1                          | 1 | 2 | 1 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG11 – Sustainable Cities and Communities     | 1            | 60         | 1                          | 2 | 1 | 2 | 1 | International Council for Science (2017)  |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG11 – Sustainable Cities and Communities     | 1            | 40         | 1                          | 0 | 1 | 2 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG11 – Sustainable Cities and Communities     | 1            | 40         | 1                          | 0 | 1 | 3 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG11 – Sustainable Cities and Communities     | 1            | 60         | 1                          | 1 | 2 | 2 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.B – Support Small-Scale Fisheries   | SDG11 – Sustainable Cities and Communities     | 1            | 60         | 1                          | 0 | 1 | 1 | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG11 – Sustainable Cities and Communities     | 1            | 80         | 1                          | 0 | 1 | 1 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.1 – Reduce Marine Pollution   | SDG12 – Responsible Consumption and Production | 2            | 60         | 3                          | 1 | 3 | 3 | 1 | Grizzetti et al. (2013); International Council for Science (2017)   |
| Economy | 14.2 – Protect and Restore Ecosystems  | SDG12 – Responsible Consumption and Production | 2            | 60         | 1                          | 3 | 3 | 3 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.3 – Reduce Ocean Acidification  | SDG12 – Responsible Consumption and Production | 1            | 80         | 1                          | 1 | 1 | 2 | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |

| Group   | From   | To   | Inter-action | Agreement | Author scores<br>(-3 to 3) |   |   |    |    | References  |
|---------|--|--|--------------|-----------|----------------------------|---|---|----|----|---|
| Economy | 14.4 – Sustainable Fishing   | SDG12 – Responsible Consumption and Production | 2            | 60        | 2                          | 3 | 3 | 3  | 1  | International Council for Science (2017)  |
| Economy | 14.5 – Conserve Coastal and Marine Areas                                       | SDG12 – Responsible Consumption and Production | 1            | 80        | 1                          | 1 | 1 | 2  | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.6 – End Subsidies Contributing to Overfishing                               | SDG12 – Responsible Consumption and Production | 1            | 40        | 0                          | 3 | 1 | 2  | 1  | Le Blanc et al. (2017)  |
| Economy | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG12 – Responsible Consumption and Production | 2            | 60        | 1                          | 2 | 1 | 3  | 1  | International Council for Science (2017)  |
| Economy | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG12 – Responsible Consumption and Production | 1            | 80        | 1                          | 1 | 2 | 1  | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.B – Support Small-Scale Fisheries   | SDG12 – Responsible Consumption and Production | 1            | 100       | 1                          | 1 | 1 | 1  | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.C – Implement and Enforce International Sea Law                             | SDG12 – Responsible Consumption and Production | 1            | 60        | 1                          | 1 | 2 | 2  | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.1 – Reduce Marine Pollution   | SDG16 – Peace and Justice                      | 1            | 60        | 1                          | 0 | 1 | 0  | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.2 – Protect and Restore Ecosystems  | SDG16 – Peace and Justice                      | 1            | 60        | 1                          | 0 | 2 | 1  | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.3 – Reduce Ocean Acidification  | SDG16 – Peace and Justice                      | 1            | 60        | 1                          | 0 | 1 | 0  | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.4 – Sustainable Fishing   | SDG16 – Peace and Justice                      | 1            | 60        | -1                         | 2 | 2 | 1  | 2  | Brashares et al. (2014)   |
| Society | 14.5 – Conserve Coastal and Marine Areas                                       | SDG16 – Peace and Justice                      | -1           | 60        | -1                         | 0 | 0 | -1 | -1 | Singh et al. (2018)   |
| Society | 14.6 – End Subsidies Contributing to Overfishing                               | SDG16 – Peace and Justice                      | 0            | 60        | 0                          | 1 | 0 | 0  | 1  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG16 – Peace and Justice                      | 1            | 60        | 1                          | 0 | 1 | 1  | 2  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG16 – Peace and Justice                      | 1            | 60        | 1                          | 0 | 1 | 1  | 2  | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |

| Group   | From   | To                                 | Inter-action | Agree-ment | Author scores<br>(-3 to 3) |   |   |    |   | References  |
|---------|--|------------------------------------|--------------|------------|----------------------------|---|---|----|---|---|
| Society | 14.B – Support Small-Scale Fisheries   | SDG16 – Peace and Justice          | 1            | 60         | 1                          | 1 | 1 | 2  | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Society | 14.C – Implement and Enforce International Sea Law                             | SDG16 – Peace and Justice          | 1            | 60         | 1                          | 2 | 1 | 1  | 2 | Pinsky et al. (2018)  |
| Economy | 14.1 – Reduce Marine Pollution   | SDG17 – Partnerships for the Goals | 0            | 60         | 1                          | 0 | 1 | -1 | 1 | Nilsson et al. (2018)   |
| Economy | 14.2 – Protect and Restore Ecosystems  | SDG17 – Partnerships for the Goals | 1            | 60         | 1                          | 0 | 1 | 1  | 2 | Unger et al. (2016)   |
| Economy | 14.3 – Reduce Ocean Acidification  | SDG17 – Partnerships for the Goals | 1            | 60         | 1                          | 0 | 1 | 0  | 1 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.4 – Sustainable Fishing   | SDG17 – Partnerships for the Goals | 1            | 40         | 1                          | 0 | 1 | -1 | 2 | Unger et al. (2016)   |
| Economy | 14.5 – Conserve Coastal and Marine Areas                                       | SDG17 – Partnerships for the Goals | 0            | 100        | 0                          | 0 | 0 | 0  | 0 | NA <sup>a</sup>   |
| Economy | 14.6 – End Subsidies Contributing to Overfishing                               | SDG17 – Partnerships for the Goals | 1            | 60         | 0                          | 1 | 0 | 0  | 2 | NA <sup>a</sup>   |
| Economy | 14.7 – Increase the Economic Benefits from Sustainable Use of Marine Resources | SDG17 – Partnerships for the Goals | 1            | 40         | 1                          | 0 | 1 | 3  | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.A – Increase Scientific Knowledge, Research and Technology for Ocean Health | SDG17 – Partnerships for the Goals | 1            | 60         | 1                          | 1 | 1 | 2  | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.B – Support Small-Scale Fisheries   | SDG17 – Partnerships for the Goals | 1            | 60         | 1                          | 0 | 1 | 1  | 2 | Nilsson et al. (2016); International Council for Science (2017); Nilsson et al. (2018); Singh et al. (2018) |
| Economy | 14.C – Implement and Enforce International Sea Law                             | SDG17 – Partnerships for the Goals | 2            | 40         | 3                          | 1 | 3 | 1  | 2 | Unger et al. (2016)   |

Note:

(a) The information is from authors' discussion.

## References

- Adams, D.H., D.M. Tremain, R. Paperno and C. Sonne, 2019: Florida lagoon at risk of ecosystem collapse. *Science*, **365**(6457), 991–992, doi:10.1126/science.aaz0175.
- Adloff, F., et al., 2015: Mediterranean Sea response to climate change in an ensemble of twenty first century scenarios. *Clim. Dyn.*, **45**(9), 2775–2802, doi:10.1007/s00382-015-2507-3.
- Agyeman, Y.B., 2019: Ecotourism as an adaptation strategy for mitigating climate change impacts on local communities around protected areas in Ghana. In: *Handbook of Climate Change Resilience* [Leal Filho, W.(ed.)]. Springer International Publishing, Cham, pp. 1–19. ISBN 978-3319710259.
- Ahas, R. and A. Aasa, 2006: The effects of climate change on the phenology of selected Estonian plant, bird and fish populations. *Int. J. Biometeorol.*, **51**(1), 17–26, doi:10.1007/s00484-006-0041-z.
- Al-Said, T., et al., 2018: High total organic carbon in surface waters of the northern Arabian Gulf: implications for the oxygen minimum zone of the Arabian Sea. *Mar. Pollut. Bull.*, **129**(1), 35–42, doi:10.1016/j.marpolbul.2018.02.013.
- Al-Yamani, F. and S.W.A. Naqvi, 2019: Chemical oceanography of the Arabian Gulf. *Deep Sea Res. Part II Top. Stud. Oceanogr.*, **161**, 72–80, doi:10.1016/j.dsr2.2018.10.003.
- Alexander, K. A., et al., 2019: Progress in integrating natural and social science in marine ecosystem-based management research. *Mar. Freshw. Res.*, **70**(1), 71–83, doi:10.1071/MF17248.
- Allison, E.H., 2011: *Aquaculture, Fisheries, Poverty and Food Security*. Working Paper 2011-65. The WorldFish Center, Penang, Malaysia, Accessed 31st December 2021, [http://pubs.iwarm.net/resource\\_centre/WF\\_2971.pdf](http://pubs.iwarm.net/resource_centre/WF_2971.pdf). (60 pp).
- Anderson, D.M., S.F.E. Boerlage and M.B. Dixon (eds.), 2017: *Harmful Algal Blooms (HABs) and Desalination: a Guide to Impacts, Monitoring and Management*. IOC Manuals and Guides, Vol. 78. Intergovernmental Oceanographic Commission of UNESCO, Paris, France. 539 pp.
- Anderson, R., 2015: Sustainability, ideology, and the politics of development in Cabo Pulmo, Baja California Sur, Mexico. *J. Polit. Ecol.*, **22**(1), 239–254, doi:10.2458/v22i1.21107.
- Anthony, K., et al., 2017: New interventions are needed to save coral reefs. *Nat. Ecol. Evol.*, **1**(10), 1420–1422, doi:10.1038/s41559-017-0313-5.
- Anthony, K.R.N., et al., 2019: *Reef Restoration and Adaptation Program: Modelling Methods and Findings*. A report provided to the Australian Government by the Reef Restoration and Adaptation Program, Australian Institute of Marine Science. 112 pp.
- Appelqvist, C. and J.N. Havenhand, 2016: A phenological shift in the time of recruitment of the shipworm, *Teredo navalis* L., mirrors marine climate change. *Ecol. Evol.*, **6**(12), 3862–3870, doi:10.1002/ece3.2126.
- Aqorau, T., J. Bell and J.N. Kittinger, 2018: Good governance for migratory species. *Science*, **361**(6408), 1208–1209, doi:10.1126/science.aav2051.
- Arafeh-Dalmau, N., et al., 2021: Incorporating climate velocity into the design of climate-smart networks of marine protected areas. *Methods Ecol. Evol.*, **12**(10), 1969–1983, doi:10.1111/2041-210X.13675.
- Archer, D., et al., 2014: Moving towards inclusive urban adaptation: approaches to integrating community-based adaptation to climate change at city and national scale. *Clim. Dev.*, **6**(4), 345–356, doi:10.1080/17565529.2014.918868.
- Arkema, K.K., et al., 2017: Linking social, ecological, and physical science to advance natural and nature-based protection for coastal communities. *Ann. N.Y. Acad. Sci.*, **1399**(1), 5–26, doi:10.1111/nyas.13322.
- Armitage, A.R., C.A. Weaver, J.S. Kominoski and S.C. Pennings, 2019: Resistance to hurricane effects varies among wetland vegetation types in the marsh-mangrove ecotone. *Estuaries Coasts*, **43**, 960–970, doi:10.1007/s12237-019-00577-3.
- Armitage, D., 2007: Governance and the commons in a multi-level world. *Int. J. Commons*, **2**(1), 7–32, doi:10.18352/ijc.28.
- Asch, R.G., 2015: Climate change and decadal shifts in the phenology of larval fishes in the California Current ecosystem. *Proc. Natl. Acad. Sci. U.S.A.*, **112**(30), E4065–E4074, doi:10.1073/pnas.1421946112.
- Atkinson, A., et al., 2015: Questioning the role of phenology shifts and trophic mismatching in a planktonic food web. *Prog. Oceanogr.*, **137**, 498–512, doi:10.1016/j.pocean.2015.04.023.
- Barange, M., et al., 2018: *Impacts of Climate Change on Fisheries and Aquaculture: Synthesis of Current Knowledge, Adaptation and Mitigation Options*. FAO Technical Report, Vol. 627. Food and Agriculture Organisation of the United Nations, Rome, Italy. 628 pp.
- Barbier, E.B., 2014: A global strategy for protecting vulnerable coastal populations. *Science*, **345**(6202), 1250–1251, doi:10.1126/science.1254629.
- Barbraud, C. and H. Weimerskirch, 2006: Antarctic birds breed later in response to climate change. *Proc. Natl. Acad. Sci. U.S.A.*, **103**(16), 6248–6251, doi:10.1073/pnas.0510397103.
- Barnard, P.L., et al., 2021: Multiple climate change-driven tipping points for coastal systems. *Sci. Rep.*, **11**(1), 15560, doi:10.1038/s41598-021-94942-7.
- Barnes, M.L., et al., 2020: Social determinants of adaptive and transformative responses to climate change. *Nat. Clim. Change*, **10**(9), 823–828, doi:10.1038/s41558-020-0871-4.
- Barnett, J. and C. McMichael, 2018: The effects of climate change on the geography and timing of human mobility. *Popul. Environ.*, **39**(4), 339–356, doi:10.1007/s11111-018-0295-5.
- Basconi, L., C. Cadier and G. Guerrero-Limón, 2020: Challenges in marine restoration ecology: how techniques, assessment metrics, and ecosystem valuation can lead to improved restoration success. In: *YOUNG MARES 9 – The Oceans: Our Research, Our Future: Proceedings of the 2018 conference for YOUNG MARINE RESEARCHER in Oldenburg, Germany* [Jungblut, S., V. Liebich and M. Bode-Dalby(eds.)]. Springer International Publishing, Cham, pp. 83–99. ISBN 978-3030203894.
- Basel, B., G. Goby and J. Johnson, 2020: Community-based adaptation to climate change in villages of Western Province, Solomon Islands. *Mar. Pollut. Bull.*, **156**, 111266, doi:10.1016/j.marpolbul.2020.111266.
- Bayraktarov, E., et al., 2016: The cost and feasibility of marine coastal restoration. *Ecol. Appl.*, **26**(4), 1055–1074, doi:10.1890/15-1077.
- Beck, M.W., et al., 2018: The global flood protection savings provided by coral reefs. *Nat. Commun.*, **9**(1), 2186, doi:10.1038/s41467-018-04568-z.
- Bell, J.D., et al., 2013: Mixed responses of tropical Pacific fisheries and aquaculture to climate change. *Nat. Clim. Change*, **3**, 591–599, doi:10.1038/nclimate1838.
- Bell, J.D., et al., 2021: Pathways to sustaining tuna-dependent Pacific Island economies during climate change. *Nat. Sustain.*, **4**, 900–910, doi:10.1038/s41893-021-00745-z.
- Bell, R.J., J. Odell, G. Kirchner and S. Lomonico, 2020: Actions to promote and achieve climate-ready fisheries: summary of current practice. *Mar. Coast. Fish.*, **12**(3), 166–190, doi:10.1002/mcf.210112.
- Bellante, A., et al., 2016: Evaluation and comparison of trace metal accumulation in different tissues of potential bioindicator organisms: macrobenthic filter feeders *Styela plicata*, *Sabella spallanzanii*, and *Mytilus galloprovincialis*. *Environ. Toxicol. Chem.*, **35**(12), 3062–3070, doi:10.1002/etc.3494.
- Ben-Hasan, A. and V. Christensen, 2019: Vulnerability of the marine ecosystem to climate change impacts in the Arabian Gulf—an urgent need for more research. *Glob. Ecol. Conserv.*, **17**, e00556, doi:10.1016/j.gecco.2019.e00556.
- Ben-Hasan, A., et al., 2018: Is reduced freshwater flow in Tigris-Euphrates rivers driving fish recruitment changes in the Northwestern Arabian Gulf? *Mar. Pollut. Bull.*, **129**(1), 1–7, doi:10.1016/j.marpolbul.2018.02.012.
- Béné, C., et al., 2016: Contribution of fisheries and aquaculture to food security and poverty reduction: assessing the current evidence. *World Dev.*, **79**, 177–196, doi:10.1016/j.worlddev.2015.11.007.

- Bennett, N.J., 2018: Navigating a just and inclusive path towards sustainable oceans. *Mar. Policy*, **97**, 139–146, doi:10.1016/j.marpol.2018.06.001.
- Bertolini, C. and J. da Mosto, 2021: Restoring for the climate: a review of coastal wetland restoration research in the last 30 years. *Restor. Ecol.*, **29**(7), e13438, doi:10.1111/rec.13438.
- Bertram, D.F., D.L. Mackas and S.M. McKinnell, 2001: The seasonal cycle revisited: interannual variation and ecosystem consequences. *Prog. Oceanogr.*, **49**(1), 27–21, doi:10.1016/S0079-6611.
- Beukema, J.J., R. Dekker and J.M. Jansen, 2009: Some like it cold: populations of the tellinid bivalve *Macoma balthica* (L.) suffer in various ways from a warming climate. *Mar. Ecol. Prog. Ser.*, **384**, 135–145, doi:10.3354/meps07952.
- Bevacqua, E., et al., 2020: More meteorological events that drive compound coastal flooding are projected under climate change. *Commun. Earth Environ.*, **1**(1), 47, doi:10.1038/s43247-020-00044-z.
- Bever, A.J., M.A.M. Friedrichs and P. St-Laurent, 2021: Real-time environmental forecasts of the Chesapeake Bay: model setup, improvements, and online visualization. *Environ. Model. Softw.*, **140**, 105036, doi:10.1016/j.envsoft.2021.105036.
- Billé, R., et al., 2017: Regional oceans governance: making regional seas programmes, regional fishery bodies and large marine ecosystem mechanisms work better together. In: *Handbook on the Economics and Management of Sustainable Oceans*. Edward Elgar Publishing, Cheltenham, UK. ISBN 978-1786430717.
- Bindoff, N.L., W.W.L. Cheung, J.G. Kairo, J. Aristegui, V.A. Guinder, R. Hallberg, N. Hilmi, N. Jiao, M.S. Karim, L. Levin, S. O'Donoghue, S.R. Purca Cuicapusa, B. Rinkevich, T. Suga, A. Tagliabue and P. Williamson, 2019: Changing Ocean, Marine Ecosystems, and Dependent Communities. In: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. [Pörtner, H.-O., D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E.S. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama and N. Weyer (eds.)] (In press).
- Birk, T. and K. Rasmussen, 2014: Migration from atolls as climate change adaptation: current practices, barriers and options in Solomon Islands. *Nat. Resour. Forum.*, **38**(1), 1–13, doi:10.1111/1477-8947.12038.
- Biswas, B. and B. Mallick, 2021: Livelihood diversification as key to long-term non-migration: evidence from coastal Bangladesh. *Environ. Dev. Sustain.*, **23**(6), 8924–8948, doi:10.1007/s10668-020-01005-4.
- Blair, A.A.C. and S. Momtaz, 2018: Climate change perception and response: case studies of fishers from Antigua and Efate. *Ocean Coast. Manag.*, **157**, 86–94, doi:10.1016/j.ocecoaman.2018.02.015.
- Blanchard, J.L., et al., 2017: Linked sustainability challenges and trade-offs among fisheries, aquaculture and agriculture. *Nat. Ecol. Evol.*, **1**(9), 1240–1249, doi:10.1038/s41559-017-0258-8.
- Blechinger, P., et al., 2016: Global analysis of the techno-economic potential of renewable energy hybrid systems on small islands. *Energy Policy*, **98**, 674–687, doi:10.1016/j.enpol.2016.03.043.
- Boerema, A. and P. Meire, 2017: Management for estuarine ecosystem services: a review. *Ecol. Eng.*, **98**, 172–182, doi:10.1016/j.ecoleng.2016.10.051.
- Bordner, A.S., C.E. Ferguson and L. Ortolano, 2020: Colonial dynamics limit climate adaptation in Oceania: Perspectives from the Marshall Islands. *Glob. Environ. Chang.*, **61**, 102054, doi:10.1016/j.gloenvcha.2020.102054.
- Boström-Einarsson, L., et al., 2020: Coral restoration – a systematic review of current methods, successes, failures and future directions. *PLoS ONE*, **15**(1), e0226631, doi:10.1371/journal.pone.0226631.
- Bott, L.-M. and B. Braun, 2019: How do households respond to coastal hazards? A framework for accommodating strategies using the example of Semarang Bay, Indonesia. *Int. J. Disaster Risk Reduct.*, **37**, 101177, doi:10.1016/j.ijdr.2019.101177.
- Bradley, D., et al., 2019: Leveraging satellite technology to create true shark sanctuaries. *Conserv. Lett.*, **12**(2), e12610, doi:10.1111/conl.12610.
- Branoff, B.L., 2020: Mangrove disturbance and response following the 2017 hurricane season in Puerto Rico. *Estuaries Coasts*, **43**(5), 1248–1262, doi:10.1007/s12237-019-00585-3.
- Brashares, J.S., et al., 2014: Wildlife decline and social conflict. *Science*, **345**(6195), 376–378, doi:10.1126/science.1256734.
- Brathwaite, A., N. Pascal and E. Clua, 2021: When are payment for ecosystems services suitable for coral reef derived coastal protection? A review of scientific requirements. *Ecosyst. Serv.*, **49**, 101261, doi:10.1016/j.ecoser.2021.101261.
- Braun de Torrez, E.C., et al., 2021: Seasick: why value ecosystems severely threatened by sea-level rise? *Estuaries Coasts*, **44**(4), 899–910, doi:10.1007/s12237-020-00850-w.
- Brito-Morales, I., et al., 2018: Climate velocity can inform conservation in a warming world. *Trends Ecol. Evol.*, **33**(6), 441–457, doi:10.1016/j.tree.2018.03.009.
- Bryndum-Buchholz, A., D.P. Tittensor and H.K. Lotze, 2021: The status of climate change adaptation in fisheries management: policy, legislation and implementation. *Fish Fish.*, **22**(6), 1248–1273, doi:10.1111/faf.12586.
- Bucci, A.F., A.C. Thomas and I. Cetinić, 2020: Interannual variability in the thermal habitat of *Alexandrium catenella* in the Bay of Fundy and the implications of climate change. *Front. Mar. Sci.*, **7**(1060), 587990, doi:10.3389/fmars.2020.587990.
- Buckley, M.C. and E.E. Crone, 2008: Negative off-site impacts of ecological restoration: understanding and addressing the conflict. *Conserv. Biol.*, **22**(5), 1118–1124, doi:10.1111/j.1523-1739.2008.01027.x.
- Bulleri, F., et al., 2018: Harnessing positive species interactions as a tool against climate-driven loss of coastal biodiversity. *PLoS Biol.*, **16**(9), e2006852, doi:10.1371/journal.pbio.2006852.
- Burden, M. and R. Fujita, 2019: Better fisheries management can help reduce conflict, improve food security, and increase economic productivity in the face of climate change. *Mar. Policy*, **108**, 103610, doi:10.1016/j.marpol.2019.103610.
- Burthe, S., et al., 2012: Phenological trends and trophic mismatch across multiple levels of a North Sea pelagic food web. *Mar. Ecol. Prog. Ser.*, **454**, 119–134, doi:10.3354/meps09520.
- Busayo, E.T. and A.M. Kalumba, 2021: Recommendations for linking climate change adaptation and disaster risk reduction in urban coastal zones: lessons from East London, South Africa. *Ocean Coast. Manag.*, **203**, 105454, doi:10.1016/j.ocecoaman.2020.105454.
- Byrd, G.V., W.J. Syderman, H.M. Renner and S. Minobe, 2008: Responses of piscivorous seabirds at the Pribilof Islands to ocean climate. *Deep Sea Res. Part II Top. Stud. Oceanogr.*, **55**(16), 1856–1867, doi:10.1016/j.dsr2.2008.04.015.
- Cabral, R.B., et al., 2018: Rapid and lasting gains from solving illegal fishing. *Nat. Ecol. Evol.*, **2**(4), 650–658, doi:10.1038/s41559-018-0499-1.
- Calcabrini, C., et al., 2017: Marine sponge natural products with anticancer potential: an updated review. *Mar. Drugs*, **15**(10), 310, doi:10.3390/md15100310.
- Cannaby, H., B.A. Fach, S.S. Arkin and B. Salihoglu, 2015: Climatic controls on biophysical interactions in the Black Sea under present day conditions and a potential future (A1B) climate scenario. *J. Mar. Syst.*, **141**, 149–166, doi:10.1016/j.jmarsys.2014.08.005.
- Carrasco Navas-Parejo, J.C., A. Corzo and S. Papaspouli, 2020: Seasonal cycles of phytoplankton biomass and primary production in a tropical temporarily open-closed estuarine lagoon—The effect of an extreme climatic event. *Sci. Total Environ.*, **723**, 138014, doi:10.1016/j.scitotenv.2020.138014.
- Carstensen, J., J.H. Andersen, B.G. Gustafsson and D.J. Conley, 2014: Deoxygenation of the Baltic Sea during the last century. *Proc. Natl. Acad. Sci. U.S.A.*, **111**(15), 5628–5633, doi:10.1073/pnas.1323156111.
- Carver, R., et al., 2020: A critical social perspective on deep sea mining: Lessons from the emergent industry in Japan. *Ocean Coast. Manag.*, **193**, 105242, doi:10.1016/j.ocecoaman.2020.105242.
- Castañeda-Moya, E., et al., 2020: Hurricanes fertilize mangrove forests in the Gulf of Mexico (Florida Everglades, USA). *Proc. Natl. Acad. Sci. U.S.A.*, **117**(9), 4831–4841, doi:10.1073/pnas.1908597117.

- CBD, 2010: *X/2.Strategic Plan for Biodiversity 2011–2020*. Accessed 31st December 2021, <https://www.cbd.int/decision/cop/?id=12268>.
- CBD Secretariat, *Strategic Plan for Biodiversity 2011 – 2020 and the Aichi Biodiversity Targets*. Sustainable Ocean Initiative, Accessed 31st December 2021, <https://www.cbd.int/soi/about/strategic-plan>.
- Chambers, L.E., P. Dann, B. Cannell and E.J. Woehler, 2014: Climate as a driver of phenological change in southern seabirds. *Int. J. Biometeorol.*, **58**(4), 603–612, doi:10.1007/s00484-013-0711-6.
- Chambers, L.G., H.E. Steinmuller and J.L. Breithaupt, 2019: Toward a mechanistic understanding of “peat collapse” and its potential contribution to coastal wetland loss. *Ecology*, **100**(7), e02720, doi:10.1002/ecy.2720.
- Cherkiss, M., et al., 2020: Shifts in hatching date of American crocodile (*Crocodylus acutus*) in southern Florida. *J. Therm. Biol.*, **88**, 102521, doi:10.1016/j.jtherbio.2020.102521.
- Cherry, S.G., A.E. Derocher, G.W. Thiemann and N.J. Lunn, 2013: Migration phenology and seasonal fidelity of an Arctic marine predator in relation to sea ice dynamics. *J. Anim. Ecol.*, **82**(4), 912–921, doi:10.1111/1365-2656.12050.
- Cheung, W.W.L., et al., 2016a: Building confidence in projections of the responses of living marine resources to climate change. *ICES J. Mar. Sci.*, **73**(5), 1283–1296, doi:10.1093/icesjms/fsv250.
- Cheung, W.W.L., M.C. Jones, G. Reygondeau and T.L. Frölicher, 2018: Opportunities for climate-risk reduction through effective fisheries management. *Glob. Change Biol.*, **24**(11), 5149–5163, doi:10.1111/gcb.14390.
- Cheung, W.W.L., et al., 2016b: Structural uncertainty in projecting global fisheries catches under climate change. *Ecol. Model.*, **325**, 57–66, doi:10.1016/j.ecolmodel.2015.12.018.
- Chevillot, X., et al., 2017: Toward a phenological mismatch in estuarine pelagic food web? *PLoS ONE*, **12**(3), e0173752, doi:10.1371/journal.pone.0173752.
- Chivers, W.J., M. Edwards and G.C. Hays, 2020: Phenological shuffling of major marine phytoplankton groups over the last six decades. *Divers. Distrib.*, **26**(5), 536–548, doi:10.1111/ddi.13028.
- Christiansen, L., A. Olhoff and T. Dale, 2020: *Understanding Adaptation in the Global Stocktake*. UNEP DTU Partnership. Part of the iGST Designing a Robust Stocktake Discussion Series. Accessed 31st December 2021, [https://www.climateworks.org/wp-content/uploads/2020/05/Understanding-Adaptation-in-the-Global-Stocktake\\_iGST\\_UNEP-DTU.pdf](https://www.climateworks.org/wp-content/uploads/2020/05/Understanding-Adaptation-in-the-Global-Stocktake_iGST_UNEP-DTU.pdf). (1-24 pp).
- Cinner, J., 2014: Coral reef livelihoods. *Curr. Opin. Environ. Sustain.*, **7**, 65–71, doi:10.1016/j.cosust.2013.11.025.
- Cinner, J.E. and M.L. Barnes, 2019: Social dimensions of resilience in social-ecological systems. *One Earth*, **1**(1), 51–56, doi:10.1016/j.oneear.2019.08.003.
- Claudet, J., et al., 2020: A roadmap for using the UN decade of ocean science for sustainable development in support of science, policy, and action. *One Earth*, **2**(1), 34–42, doi:10.1016/j.oneear.2019.10.012.
- Cline, T.J., D.E. Schindler and R. Hilborn, 2017: Fisheries portfolio diversification and turnover buffer Alaskan fishing communities from abrupt resource and market changes. *Nat. Commun.*, **8**(1), 14042, doi:10.1038/ncomms14042.
- Cobb, C.K., 2020: *The Impact of Climate Change on the Migration Phenology of New England’s Anadromous River Herring and American Shad*. Colby College, Maine, USA.
- Cochrane, K.L., 2021: Reconciling sustainability, economic efficiency and equity in marine fisheries: Has there been progress in the last 20 years? *Fish Fish.*, **22**(2), 298–323, doi:10.1111/faf.12521.
- Collins M., M. Sutherland, L. Bouwer, S.-M. Cheong, T. Frölicher, H. Jacot Des Combes, M. Koll Roxy, I. Losada, K. McInnes, B. Ratter, E. Rivera-Arriaga, R.D. Susanto, D. Swingedouw, and L. Tibig, 2019: Extremes, Abrupt Changes and Managing Risks. In: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. [Pörtner, H.-O., D. C. Roberts, V. Masson-Delmonte, P. Zhai, M. Tignor, E. S. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama and N. Weyer (eds.)] (In press).
- Colls, A., N. Ash and N. Ikkala, 2009: *Ecosystem-based Adaptation: a Natural Response to Climate Change*. IUCN, Gland, Switzerland, Accessed 31st December 2021, <https://www.iucn.org/content/ecosystem-based-adaptation-a-natural-response-climate-change>. (16 pp).
- Conservation International and IUCN, 2021: *Addressing Ocean and Climate Issues Across Relevant Multilateral Environmental Agreements*. Accessed 31st December 2021, [https://www.iucn.org/sites/dev/files/content/documents/policybrief\\_climatechangeandmultilateralagreements.pdf](https://www.iucn.org/sites/dev/files/content/documents/policybrief_climatechangeandmultilateralagreements.pdf). (1-8 pp).
- Copping, A., et al., 2014: An international assessment of the environmental effects of marine energy development. *Ocean Coast. Manag.*, **99**, 3–13, doi:10.1016/j.ocecoaman.2014.04.002.
- Corten, A., C.-B. Braham and A.S. Sadegh, 2017: The development of a fishmeal industry in Mauritania and its impact on the regional stocks of sardinella and other small pelagics in Northwest Africa. *Fish. Res.*, **186**, 328–336, doi:10.1016/j.fishres.2016.10.009.
- Coscieme, L., et al., 2020: Multiple conceptualizations of nature are key to inclusivity and legitimacy in global environmental governance. *Environ. Sci. Policy*, **104**, 36–42, doi:10.1016/j.envsci.2019.10.018.
- Costello, C., et al., 2020a: The future of food from the sea. *Nature*, **588**, 95–100, doi:10.1038/s41586-020-2616-y.
- Costello, C., et al., 2020b: Ambitious subsidy reform by the WTO presents opportunities for ocean health restoration. *Sustain. Sci.*, **16**, 1391–1396, doi:10.1007/s11625-020-00865-z.
- Costello, C., et al., 2016: Global fishery prospects under contrasting management regimes. *Proc. Natl. Acad. Sci. U.S.A.*, **113**(18), 5125–5129, doi:10.1073/pnas.1520420113.
- Costello, J.H., B.K. Sullivan and D.J. Gifford, 2006: A physical–biological interaction underlying variable phenological responses to climate change by coastal zooplankton. *J. Plankton Res.*, **28**(11), 1099–1105, doi:10.1093/plankt/fbl042.
- Cottrell, R.S., et al., 2020: Global adoption of novel aquaculture feeds could substantially reduce forage fish demand by 2030. *Nat. Food*, **1**(5), 301–308, doi:10.1038/s43016-020-0078-x.
- Cramer, W., et al., 2018: Climate change and interconnected risks to sustainable development in the Mediterranean. *Nat. Clim. Change*, **8**(11), 972–980, doi:10.1038/s41558-018-0299-2.
- Crozier, L.G., M.D. Scheuerell and R.W. Zabel, 2011: Using time series analysis to characterize evolutionary and plastic responses to environmental change: a case study of a shift toward earlier migration date in sockeye salmon. *Am. Nat.*, **178**(6), 755–773, doi:10.1086/662669.
- Cullen, J.M., L.E. Chambers, P.C. Coutin and P. Dann, 2009: Predicting onset and success of breeding in little penguins *Eudyptula minor* from ocean temperatures. *Mar. Ecol. Prog. Ser.*, **378**, 269–278, doi:10.3354/meps07881.
- Cvitanovic, C., et al., 2016: Linking adaptation science to action to build food secure Pacific Island communities. *Clim. Risk Manag.*, **11**, 53–62, doi:10.1016/j.crm.2016.01.003.
- Cvitanovic, C., et al., 2018: Governing fisheries through the critical decade: the role and utility of polycentric systems. *Rev. Fish Biol. Fish.*, **28**(1), 1–18, doi:10.1007/s11160-017-9495-9.
- D’Alba, L., P.A.T. Monaghan and R.G. Nager, 2010: Advances in laying date and increasing population size suggest positive responses to climate change in Common Eiders *Somateria mollissima* in Iceland. *Ibis*, **152**(1), 19–28, doi:10.1111/j.1474-919X.2009.00978.x.
- Dandy, J., et al., 2019: Leaving home: place attachment and decisions to move in the face of environmental change. *Reg. Environ. Change*, **19**(2), 615–620, doi:10.1007/s10113-019-01463-1.
- Darmaraki, S., S. Somot, F. Sevault and P. Nabat, 2019: Past variability of Mediterranean Sea marine heatwaves. *Geophys. Res. Lett.*, **46**(16), 9813–9823, doi:10.1029/2019GL082933.
- Davidson, K., et al., 2021: HABreports: online early warning of harmful algal and biotoxin risk for the scottish shellfish and finfish aquaculture industries. *Front. Mar. Sci.*, **8**, 350, doi:10.3389/fmars.2021.631732.
- Dawson, J., et al., 2020: Infusing Inuit and local knowledge into the Low Impact Shipping Corridors: an adaptation to increased shipping activity and climate

- change in Arctic Canada. *Environ. Sci. Policy*, **105**, 19–36, doi:10.1016/j.envsci.2019.11.013.
- Dawson, J., E.J. Stewart, M. E. Johnston and C.J. Lemieux, 2016: Identifying and evaluating adaptation strategies for cruise tourism in Arctic Canada. *J. Sustain. Tour.*, **24**(10), 1425–1441, doi:10.1080/09669582.2015.1125358.
- de Schipper, M. A., et al., 2021: Beach nourishment has complex implications for the future of sandy shores. *Nat. Rev. Earth Environ.*, **2**(1), 70–84, doi:10.1038/s43017-020-00109-9.
- Deb, A.K. and C.E. Haque, 2016: Livelihood diversification as a climate change coping strategy adopted by small-scale fishers of Bangladesh. In: *Climate Change Adaptation, Resilience and Hazards* [Leal Filho, W., H. Musa, G. Cavan, P. O'Hare and J. Seixas(eds.)]. Springer International Publishing, Cham, pp. 345–368. ISBN 978-3319398808.
- Dembele, Y.M., L.A. Akinbile and O.O. Aminu, 2019: Adaptation strategies to climate change among cereal crop farmers in Kita, Kayes region of Mali. *J. Agric. Ext.*, **23**(3), 107–116, doi:10.4314/jae.v23i3.9.
- Descamps, S., et al., 2019: Diverging phenological responses of Arctic seabirds to an earlier spring. *Glob. Change Biol.*, **25**(12), 4081–4091, doi:10.1111/gcb.14780.
- Devereux, S., 2016: Social protection for enhanced food security in sub-Saharan Africa. *Food Policy*, **60**, 52–62, doi:10.1016/j.foodpol.2015.03.009.
- Díaz-Almela, E., N. Marbà and C.M. Duarte, 2007: Consequences of Mediterranean warming events in seagrass (*Posidonia oceanica*) flowering records. *Glob. Change Biol.*, **13**(1), 224–235, doi:10.1111/j.1365-2486.2006.01260.x.
- Diedrich, A., C. Benham, L. Pandihau and M. Sheaves, 2019: Social capital plays a central role in transitions to sportfishing tourism in small-scale fishing communities in Papua New Guinea. *Ambio*, **48**(4), 385–396, doi:10.1007/s13280-018-1081-4.
- Donner, S.D. and S. Webber, 2014: Obstacles to climate change adaptation decisions: a case study of sea-level rise and coastal protection measures in Kiribati. *Sustain. Sci.*, **9**(3), 331–345, doi:10.1007/s11625-014-0242-z.
- Duarte, C.M., et al., 2020a: Rebuilding marine life. *Nature*, **580**(7801), 39–51, doi:10.1038/s41586-020-2146-7.
- Duarte, C.M., et al., 2020b: Data set on restoration projects of coastal marine habitats reported worldwide. *PANGAEA*, doi:10.1594/PANGAEA.912232.
- Dubik, B.A., et al., 2019: Governing fisheries in the face of change: social responses to long-term geographic shifts in a U.S. fishery. *Mar. Policy*, **99**, 243–251, doi:10.1016/j.marpol.2018.10.032.
- Dufour, F., H. Arrizabalaga, X. Irigoien and J. Santiago, 2010: Climate impacts on albacore and bluefin tunas migrations phenology and spatial distribution. *Prog. Oceanogr.*, **86**(1–2), 283–290, doi:10.1016/j.pocean.2010.04.007.
- Dunstan, P.K., et al., 2018: How can climate predictions improve sustainability of coastal fisheries in Pacific small-island developing states? *Mar. Policy*, **88**, 295–302, doi:10.1016/j.marpol.2017.09.033.
- Edgar, G.J., et al., 2014: Global conservation outcomes depend on marine protected areas with five key features. *Nature*, **506**(7487), 216–220, doi:10.1038/nature13022.
- Edwards, M., et al., 2006: Ecological status report: results from the CPR survey 2004/2005. *SAHFOS Tech. Rep.*, **3**, 1–8. Accessed 31st December 2021, Available at: [http://plymsea.ac.uk/id/eprint/1903/1/ecological\\_status\\_%28ebook%29.pdf](http://plymsea.ac.uk/id/eprint/1903/1/ecological_status_%28ebook%29.pdf).
- Edwards, M. and A.J. Richardson, 2004: Impact of climate change on marine pelagic phenology and trophic mismatch. *Nature*, **430**(7002), 881–884, doi:10.1038/nature02808.
- Ehler, C. and D. Fanny, 2009: *Marine Spatial Planning: a Step-by-Step Approach Toward Ecosystem-based Management*. [Intergovernmental Oceanographic Commission and Man and the Biosphere Programme (ed.)]. IOC Manual and Guides No. 53, ICAM Dossier No. 6. UNESCO, Paris, Accessed 31st December 2021, <https://repository.oceanbestpractices.org/bitstream/handle/11329/459/186559e.pdf?sequence=1&isAllowed=y>.
- Elko, N., et al., 2021: A century of U.S. beach nourishment. *Ocean Coast. Manag.*, **199**, 105406, doi:10.1016/j.ocgeoaman.2020.105406.
- Ellabban, O., H. Abu-Rub and F. Blaabjerg, 2014: Renewable energy resources: current status, future prospects and their enabling technology. *Renew. Sustain. Energy Rev.*, **39**, 748–764, doi:10.1016/j.rser.2014.07.113.
- Ellison, A.M., A.J. Felson and D.A. Friess, 2020: Mangrove rehabilitation and restoration as experimental adaptive management. *Front. Mar. Sci.*, **7**, 327, doi:10.3389/fmars.2020.00327.
- Émond, K., B. Sainte-Marie and J. Béty, 2020: Long-term trends and drivers of larval phenology and abundance of dominant brachyuran crabs in the Gulf of St. Lawrence (Canada). *Fish. Oceanogr.*, **29**(2), 185–200, doi:10.1111/fog.12463.
- Engler, C., 2020: Transboundary fisheries, climate change, and the ecosystem approach: taking stock of the international law and policy seascape. *Ecol. Soc.*, **25**(4), 43, doi:10.5751/ES-11988-250443.
- Erauskin-Extramiana, M., et al., 2019: Large-scale distribution of tuna species in a warming ocean. *Glob. Change Biol.*, **25**(6), 2043–2060, doi:10.1111/gcb.14630.
- Ermgassen, P.S.E., J.H. Grabowski, J.R. Gair and S.P. Powers, 2016: Quantifying fish and mobile invertebrate production from a threatened nursery habitat. *J. Appl. Ecol.*, **53**(2), 596–606, doi:10.1111/1365-2664.12576.
- Espeland, E.K. and K.M. Kettenring, 2018: Strategic plant choices can alleviate climate change impacts: a review. *J. Environ. Manag.*, **222**, 316–324, doi:10.1016/j.jenvman.2018.05.042.
- European Commission, 2012: *Blue Growth: Opportunities for Marine and Maritime Sustainable Growth*. Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee Of The Regions. European Commission, Brussels, Belgium, Accessed 31 December 2021, [https://blackseablueconomy.eu/sites/default/files/blue\\_growth\\_opportunities\\_for\\_marine\\_and\\_maritime\\_sustainable\\_growth.pdf](https://blackseablueconomy.eu/sites/default/files/blue_growth_opportunities_for_marine_and_maritime_sustainable_growth.pdf). (12 pp).
- Fabinyi, M., 2020: The role of land tenure in livelihood transitions from fishing to tourism. *Marit. Stud.*, **19**(1), 29–39, doi:10.1007/s40152-019-00145-2.
- Fadli, N., et al., 2012: The role of habitat creation in coral reef conservation: a case study from Aceh, Indonesia. *Oryx*, **46**(4), 501–507, doi:10.1017/S0030605312000142.
- FAO, 2009: *The State of World Fisheries and Aquaculture 2008*. Food and Agriculture Organization of the United Nations, Rome, Italy, Accessed 31st December 2021, <http://www.fao.org/3/a-i0250e.pdf>. (176 pp).
- FAO, 2015a: *Fishers' Knowledge and the Ecosystem Approach to Fisheries: Applications, Experiences and Lessons in Latin America*. [Fischer, J., J. Jorgensen, H. Josupeit, D. Kalikoski and C. M. Lucas (eds.)]. FAO Fisheries and Aquaculture Technical Paper No. 591. Food and Agriculture Organization of the United Nations, Rome, Italy, Accessed 31st December 2021, <http://www.fao.org/3/a-i4664e.pdf>. (278 pp).
- FAO, 2015b: *Voluntary Guidelines for Securing Sustainable Small-scale Fisheries in the Context of Food Security and Poverty Eradication*. Nations, Food and Agriculture Organization of the United Nations, Rome, Italy, Accessed 31st December 2021, <http://www.fao.org/3/a-i4356en.pdf>. (19 pp).
- FAO, 2016: *The State of World Fisheries and Aquaculture 2016. Contributing to Food Security and Nutrition for All*. Food and Agriculture Organization of the United Nations, Rome, Italy, Accessed 31st December 2021, <http://www.fao.org/3/a-i5555e.pdf>. (200 pp).
- Feher, L.C., et al., 2020: The long-term effects of Hurricanes Wilma and Irma on soil elevation change in everglades mangrove forests. *Ecosystems*, **23**(5), 917–931, doi:10.1007/s10021-019-00446-x.
- Fernández-Montblanc, T., et al., 2019: Towards robust pan-European storm surge forecasting. *Ocean Model.*, **133**, 129–144, doi:10.1016/j.ocemod.2018.12.001.
- Fernandino, G., C.I. Elliff and I.R. Silva, 2018: Ecosystem-based management of coastal zones in face of climate change impacts: challenges and inequalities. *J. Environ. Manag.*, **215**, 32–39, doi:10.1016/j.jenvman.2018.03.034.
- Ferrario, F., et al., 2014: The effectiveness of coral reefs for coastal hazard risk reduction and adaptation. *Nat. Commun.*, **5**(1), 3794, doi:10.1038/ncomms4794.

- Field, C.R., et al., 2019: Framework for quantifying population responses to disturbance reveals that coastal birds are highly resilient to hurricanes. *Ecol. Lett.*, **22**(12), 2039–2048, doi:10.1111/ele.13384.
- Fincham, J.I., A.D. Rijnsdorp and G.H. Engelhard, 2013: Shifts in the timing of spawning in sole linked to warming sea temperatures. *J. Sea Res.*, **75**, 69–76, doi:10.1016/j.seares.2012.07.004.
- Finlayson, C.M., et al., 2017: Policy considerations for managing wetlands under a changing climate. *Mar. Freshw. Res.*, **68**(10), 1803–1815, doi:10.1071/MF16244.
- Fitchett, J.M., S.W. Grab and H. Portwig, 2019: Progressive delays in the timing of sardine migration in the southwest Indian Ocean. *S. Afr. J. Sci.*, **115**(7), 8, doi:10.17159/sajs.2019/5887.
- Flood, S., N.A. Cradock-Henry, P. Blackett and P. Edwards, 2018: Adaptive and interactive climate futures: systematic review of 'serious games' for engagement and decision-making. *Environ. Res. Lett.*, **13**(6), 063005, doi:10.1088/1748-9326/aac1c6.
- Ford, J.D., et al., 2020: The resilience of Indigenous peoples to environmental change. *One Earth*, **2**(6), 532–543, doi:10.1016/j.oneear.2020.05.014.
- Forsblom, L., et al., 2019: Environmental variables driving species and genus level changes in annual plankton biomass. *J. Plankton Res.*, **41**(6), 925–938, doi:10.1093/plankt/fbz063.
- Foti, E., R.E. Musumeci and M. Stagnitti, 2020: Coastal defence techniques and climate change: a review. *Rend. Lincei Sci. Fis. Nat.*, **31**(1), 123–138, doi:10.1007/s12210-020-00877-y.
- Fox, H.E., et al., 2019: Rebuilding coral reefs: success (and failure) 16 years after low-cost, low-tech restoration. *Restor. Ecol.*, **27**(4), 862–869, doi:10.1111/rec.12935.
- Frazão Santos, C., et al., 2020: Integrating climate change in ocean planning. *Nat. Sustain.*, **3**(7), 505–516, doi:10.1038/s41893-020-0513-x.
- Frederiksen, M., et al., 2004: Scale-dependent climate signals drive breeding phenology of three seabird species. *Glob. Change Biol.*, **10**(7), 1214–1221, doi:10.1111/j.1529-8817.2003.00794.x.
- Fredriksen, S., et al., 2020: Green gravel: a novel restoration tool to combat kelp forest decline. *Sci. Rep.*, **10**(1), 3983, doi:10.1038/s41598-020-60553-x.
- Free, C.M., et al., 2020: Realistic fisheries management reforms could mitigate the impacts of climate change in most countries. *PLoS ONE*, **15**(3), e0224347, doi:10.1371/journal.pone.0224347.
- Free, C.M., et al., 2019: Impacts of historical warming on marine fisheries production. *Science*, **363**(6430), 979–983, doi:10.1126/science.aau1758.
- Friedman, W.R., et al., 2020: Research priorities for achieving healthy marine ecosystems and human communities in a changing climate. *Front. Mar. Sci.*, **7**, 5, doi:10.3389/fmars.2020.00005.
- Friess, D.A., et al., 2020: Mangrove blue carbon in the face of deforestation, climate change, and restoration. *Annu. Plant Rev. Online*, **3**(3), 427–456, doi:10.1002/9781119312994.apr0752.
- Friess, D.A., et al., 2019: The state of the world's mangrove forests: past, present, and future. *Annu. Rev. Environ. Resour.*, **44**, 89–115, doi:10.1146/annurev-environ-101718-033302.
- Froehlich, H.E., et al., 2020: Mind the gap between ICES nations' future seafood consumption and aquaculture production. *ICES J. Mar. Sci.*, **78**(1), 468–477, doi:10.1093/icesjms/fsaa066.
- Froehlich, H.E., R.R. Gentry and B.S. Halpern, 2018a: Global change in marine aquaculture production potential under climate change. *Nat. Ecol. Evol.*, **2**(11), 1745–1750, doi:10.1038/s41559-018-0669-1.
- Froehlich, H.E., et al., 2018b: Avoiding the ecological limits of forage fish for fed aquaculture. *Nat. Sustain.*, **1**(6), 298–303, doi:10.1038/s41893-018-0077-1.
- Fulton, E.A., et al., 2019a: Ecosystems say good management pays off. *Fish Fish.*, **20**(1), 66–96, doi:10.1111/faf.12324.
- Fulton, S., et al., 2019b: Untapped potential of citizen science in Mexican small-scale fisheries. *Front. Mar. Sci.*, **6**, 517, doi:10.3389/fmars.2019.00517.
- Fuso Nerini, F., et al., 2018: Mapping synergies and trade-offs between energy and the Sustainable Development Goals. *Nat. Energy*, **3**(1), 10–15, doi:10.1038/s41560-017-0036-5.
- Gaichas, S.K., et al., 2018: Implementing ecosystem approaches to fishery management: risk assessment in the US Mid-Atlantic. *Front. Mar. Sci.*, **5**, 442, doi:10.3389/fmars.2018.00442.
- Gaines, S.D., et al., 2018: Improved fisheries management could offset many negative effects of climate change. *Sci. Adv.*, **4**(8), aao1378, doi:10.1126/sciadv.aao1378.
- Galappaththi, E.K., et al., 2020: Climate change adaptation in aquaculture. *Rev. Aquac.*, **12**(4), 2160–2176, doi:10.1111/raq.12427.
- Galappaththi, I.M., E.K. Galappaththi and S.S. Kodithuwakku, 2017: Can start-up motives influence social-ecological resilience in community-based entrepreneurship setting? Case of coastal shrimp farmers in Sri Lanka. *Mar. Policy*, **86**, 156–163, doi:10.1016/j.marpol.2017.09.024.
- Gallo, N.D., D.G. Victor and L.A. Levin, 2017: Ocean commitments under the Paris Agreement. *Nat. Clim. Change*, **7**(11), 833–838, doi:10.1038/nclimate3422.
- Gaston, A.J., H.G. Gilchrist and J.M. Hipfner, 2005: Climate change, ice conditions and reproduction in an Arctic nesting marine bird: Brunnich's guillemot (*Uria lomvia* L.). *J. Anim. Ecol.*, **74**(5), 832–841, doi:10.1111/j.1365-2656.2005.00982.x.
- Gattuso, J.-P., et al., 2018: Ocean solutions to address climate change and its effects on marine ecosystems. *Front. Mar. Sci.*, **5**, 337, doi:10.3389/fmars.2018.00337.
- Gegg, P. and V. Wells, 2019: The development of seaweed-derived fuels in the UK: an analysis of stakeholder issues and public perceptions. *Energy Policy*, **133**, 110924, doi:10.1016/j.enpol.2019.110924.
- Genevier, L.G.C., et al., 2019: Marine heatwaves reveal coral reef zones susceptible to bleaching in the Red Sea. *Glob. Change Biol.*, **25**(7), 2338–2351, doi:10.1111/gcb.14652.
- Gephart, J.A., et al., 2020: Scenarios for global aquaculture and its role in human nutrition. *Rev. Fish. Sci. Aquac.*, **29**(1), 122–138, doi:10.1080/23308249.2020.1782342.
- Gianelli, I., et al., 2021: Harnessing scientific and local knowledge to face climate change in small-scale fisheries. *Glob. Environ. Change*, **68**, 102253, doi:10.1016/j.gloenvcha.2021.102253.
- Gibbs, M.T., 2015: Pitfalls in developing coastal climate adaptation responses. *Clim. Risk Manag.*, **8**, 1–8, doi:10.1016/j.crm.2015.05.001.
- Gilby, B.L., et al., 2018: Maximizing the benefits of oyster reef restoration for finfish and their fisheries. *Fish Fish.*, **19**(5), 931–947, doi:10.1111/faf.12301.
- Gillfillan, D., 2019: The health sector's role in governance of climate change adaptation in Myanmar. *Clim. Dev.*, **11**(7), 574–584, doi:10.1080/17565529.2018.1510364.
- Gill, D.A., et al., 2017: Capacity shortfalls hinder the performance of marine protected areas globally. *Nature*, **543**(7647), 665–669, doi:10.1038/nature21708.
- Gillett, R. and C. Lightfoot, 2001: *The Contribution of Fisheries to the Economies of Pacific Island Countries*. Asian Development Bank, Manila, Philippines, Accessed 31st December 2021, [\(217 pp\)](https://think-asia.org/handle/11540/2649).
- Gissi, E., S. Fraschetti and F. Micheli, 2019: Incorporating change in marine spatial planning: a review. *Environ. Sci. Policy*, **92**, 191–200, doi:10.1016/j.envsci.2018.12.002.
- Gissi, E., et al., 2021: A review of the combined effects of climate change and other local human stressors on the marine environment. *Sci. Total Environ.*, **755**, 142564, doi:10.1016/j.scitotenv.2020.142564.
- Global Ocean Oxygen Network, 2018: *The Ocean is Losing Its Breath: Declining Oxygen in the World's Ocean and Coastal Waters*. [Breitburg, D., M. Gregoire and K. Isensee (eds.)]. IOC Technical Series. IOC-UNESCO, Paris, Accessed 31st December 2021, [\(40 pp\)](https://unesdoc.unesco.org/ark:/48223/pf0000265196).
- Gonzalez-Mon, B., et al., 2021: Spatial diversification as a mechanism to adapt to environmental changes in small-scale fisheries. *Environ. Sci. Policy*, **116**, 246–257, doi:10.1016/j.envsci.2020.11.006.

- Gordon, T.A.C., A.N. Radford, S.D. Simpson and M.G. Meekan, 2020: Marine restoration projects are undervalued. *Science*, **367**(6478), 635–636, doi:10.1126/science.aba9141.
- Greve, W., F. Reiners and J. Nast, 1996: Biocoenotic changes of the zooplankton in the German Bight: the possible effects of eutrophication and climate. *ICES J. Mar. Sci.*, **53**(6), 951–956, doi:10.1006/jmsc.1996.0117.
- Grizzetti, B., et al., 2013: The contribution of food waste to global and European nitrogen pollution. *Environ. Sci. Policy*, **33**, 186–195, doi:10.1016/j.envsci.2013.05.013.
- Guinder, V.A., C.A. Popovich, J.C. Molinero and G.M. E. Perillo, 2010: Long-term changes in phytoplankton phenology and community structure in the Bahía Blanca Estuary, Argentina. *Mar. Biol.*, **157**(12), 2703–2716, doi:10.1007/s00227-010-1530-5.
- Güraslan, C., B.A. Fach and T. Oguz, 2017: Understanding the impact of environmental variability on anchovy overwintering migration in the Black Sea and its implications for the fishing industry. *Front. Mar. Sci.*, **4**, 275, doi:10.3389/fmars.2017.00275.
- Gurney, G.G., et al., 2021: Biodiversity needs every tool in the box: use OECMs. *Nature*, **595**, 646–649, doi:10.1038/d41586-021-02041-4.
- Haas, B., et al., 2021: The future of ocean governance. *Rev. Fish Biol. Fish.*, doi:10.1007/s11160-020-09631-x.
- Hafezi, M., et al., 2021: Evaluating coral reef ecosystem services outcomes from climate change adaptation strategies using integrative system dynamics. *J. Environ. Manag.*, **285**, 112082, doi:10.1016/j.jenvman.2021.112082.
- Hanson, S.E. and R.J. Nicholls, 2020: Demand for ports to 2050: climate policy, growing trade and the impacts of sea-level rise. *Earth's Future*, **8**(8), e2020EF001543, doi:10.1029/2020EF001543.
- Harper, S., et al., 2013: Women and fisheries: contribution to food security and local economies. *Mar. Policy*, **39**, 56–63, doi:10.1016/j.marpol.2012.10.018.
- Harvey, B.J., K.L. Nash, J.L. Blanchard and D.P. Edwards, 2018: Ecosystem-based management of coral reefs under climate change. *Ecol. Evol.*, **8**(12), 6354–6368, doi:10.1002/ece3.4146.
- Hassan, M.A., et al., 2008: Salmon-driven bed load transport and bed morphology in mountain streams. *Geophys. Res. Lett.*, **35**(4), doi:10.1029/2007GL032997.
- Hauser, D.D.W., et al., 2017: Decadal shifts in autumn migration timing by Pacific Arctic beluga whales are related to delayed annual sea ice formation. *Glob. Change Biol.*, **23**(6), 2206–2217, doi:10.1111/gcb.13564.
- Hawkes, L.A., A.C. Broderick, M.H. Godfrey and B.J. Godley, 2007: Investigating the potential impacts of climate change on a marine turtle population. *Glob. Change Biol.*, **13**(5), 923–932, doi:10.1111/j.1365-2486.2007.01320.x.
- Hazen, E.L., et al., 2018: A dynamic ocean management tool to reduce bycatch and support sustainable fisheries. *Sci. Adv.*, **4**(5), eaar3001, doi:10.1126/sciadv.aar3001.
- Hein, M.Y., et al., 2019: Coral restoration: socio-ecological perspectives of benefits and limitations. *Biol. Conserv.*, **229**, 14–25, doi:10.1016/j.biocon.2018.11.014.
- Hein, M.Y., et al., 2021: Perspectives on the use of coral reef restoration as a strategy to support and improve reef ecosystem services. *Front. Mar. Sci.*, **8**(299), 618303, doi:10.3389/fmars.2021.618303.
- Hein, M.Y., B.L. Willis, R. Beeden and A. Birtles, 2017: The need for broader ecological and socioeconomic tools to evaluate the effectiveness of coral restoration programs. *Restor. Ecol.*, **25**(6), 873–883, doi:10.1111/rec.12580.
- Henderson, M. E., et al., 2017: Effects of spring onset and summer duration on fish species distribution and biomass along the Northeast United States continental shelf. *Rev. Fish Biol. Fish.*, **27**(2), 411–424, doi:10.1007/s11160-017-9487-9.
- Heneghan, R.F., et al., 2021: Disentangling diverse responses to climate change among global marine ecosystem models. *Prog. Oceanogr.*, **198**, 102659, doi:10.1016/j.pocean.2021.102659.
- Heron, S.F., et al., 2017: *Impacts of Climate Change on World Heritage Coral Reefs: A First Global Scientific Assessment*. UNESCO World Heritage Centre, Paris, France, Accessed 31st December 2021, <https://repository.library.noaa.gov/view/noaa/16386>.
- Hettiarachchi, M., T.H. Morrison and C. McAlpine, 2015: Forty-three years of Ramsar and urban wetlands. *Glob. Environ. Change*, **32**, 57–66, doi:10.1016/j.gloenvcha.2015.02.009.
- Himes-Cornell, A., L. Pendleton and P. Atiyah, 2018: Valuing ecosystem services from blue forests: a systematic review of the valuation of salt marshes, sea grass beds and mangrove forests. *Ecosyst. Serv.*, **30**, 36–48, doi:10.1016/j.ecoser.2018.01.006.
- Hindell, M. A., et al., 2012: Long-term breeding phenology shift in royal penguins. *Ecol. Evol.*, **2**(7), 1563–1571, doi:10.1002/ece3.281.
- Hinkel, J., et al., 2018: The ability of societies to adapt to twenty-first-century sea-level rise. *Nat. Clim. Change*, **8**(7), 570–578, doi:10.1038/s41558-018-0176-z.
- Hinrichsen, H.H., et al., 2016: Oxygen depletion in coastal seas and the effective spawning stock biomass of an exploited fish species. *R. Soc. Open Sci.*, **3**(1), 150338, doi:10.1098/rsos.150338.
- Hjerne, O., et al., 2019: Climate driven changes in timing, composition and magnitude of the baltic sea phytoplankton spring bloom. *Front. Mar. Sci.*, **6**, 482, doi:10.3389/fmars.2019.00482.
- HLPE, 2014: *Sustainable Fisheries and Aquaculture for Food Security and Nutrition*. HLPE Report 7.High Level Panel of Experts on Food Security and Nutrition of the Committee on World Food Security, Rome, Italy, Accessed 31st December 2021, <https://www.unscn.org/uploads/web/news/HLPE-Report-7-EN.pdf>. (118 pp).
- Ho, T.V.T., A. Cottrell, P. Valentine and S. Woodley, 2012: Perceived barriers to effective multilevel governance of human-natural systems: an analysis of Marine Protected Areas in Vietnam. *J. Polit. Ecol.*, **19**(1), 17–35, doi:10.2458/v19i1.21711.
- Hobday, A.J., et al., 2018: A framework for combining seasonal forecasts and climate projections to aid risk management for fisheries and aquaculture. *Front. Mar. Sci.*, **5**, 137, doi:10.3389/fmars.2018.00137.
- Hobday, A.J., C.M. Spillman, J. Paige Eveson and J.R. Hartog, 2016: Seasonal forecasting for decision support in marine fisheries and aquaculture. *Fish. Oceanogr.*, **25**(S1), 45–56, doi:10.1111/fog.12083.
- Hoegh-Guldberg, O. and J.F. Bruno, 2010: The impact of climate change on the world's marine ecosystems. *Science*, **328**(5985), 1523–1528, doi:10.1126/science.1189930.
- Hoegh-Guldberg, O., et al., 2014: The Ocean. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*. [Barros, V. R., C. B. Field, D. J. Dokken, M. D. Mastrandrea, K. J. Mach, T. E. Bilir, M. Chatterjee, K. L. Ebi, Y. O. Estrada, R. C. Genova, B. Girma, E. S. Kissel, A. N. Levy, S. MacCracken, P. R. Mastrandrea and L. L. White (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 1655–1731. ISBN 978-1107058163.
- Hoegh-Guldberg, O., D. Jacob, M. Taylor, M. Bind, S. Brown, I. Camilloni, A. Diedhiou, R. Djalante, K.L. Ebi, F. Engelbrecht, J. Guiot, Y. Hijikata, S. Mehrotra, A. Payne, S.I. Seneviratne, A. Thomas, R. Warren, and G. Zhou, et al., 2018: Impacts of 1.5°C Global Warming on Natural and Human Systems. In: *Global Warming of 1.5°C. An IPCC Special Report on the Impacts of Global Warming of 1.5°C Above Pre-industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*. [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P. R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J. B. R. Matthews, Y. Chen, X. Zhou, M. I. Gomis, E. Lonnoy, T. Maycock, M. Tignor and T. Waterfield (eds.)](In Press).
- Hoegh-Guldberg, O., E.S. Poloczanska, W. Skirving and S. Dove, 2017: Coral reef ecosystems under climate change and ocean acidification. *Front. Mar. Sci.*, **4**, 158, doi:10.3389/fmars.2017.00158.
- Holbrook, N.J., et al., 2020: Keeping pace with marine heatwaves. *Nat. Rev. Earth Environ.*, **1**(9), 482–493, doi:10.1038/s43017-020-0068-4.

- Holding, S., et al., 2016: Groundwater vulnerability on small islands. *Nat. Clim. Change*, **6**(12), 1100–1103, doi:10.1038/nclimate3128.
- Holsman, K., et al., 2017: An ecosystem-based approach to marine risk assessment. *Ecosyst. Health Sustain.*, **3**(1), e01256, doi:10.1002/ehs2.1256.
- Holsman, K.K., et al., 2020: Ecosystem-based fisheries management forestalls climate-driven collapse. *Nat. Commun.*, **11**(1), 4579, doi:10.1038/s41467-020-18300-3.
- Holsman, K.K., et al., 2019: Towards climate resiliency in fisheries management. *ICES J. Mar. Sci.*, **76**(5), 1368–1378, doi:10.1093/icesjms/fsz031.
- Hosia, A., T. Falkenhaug and L.J. Naustvoll, 2014: Trends in abundance and phenology of *Aurelia aurita* and *Cyanea* spp. at a Skagerrak location, 1992–2011. *Mar. Ecol. Prog. Ser.*, **498**, 103–115, doi:10.3354/meps10619.
- Hossain, M.A.R., M. Ahmed, E. Ojea and J.A. Fernandes, 2018: Impacts and responses to environmental change in coastal livelihoods of southwest Bangladesh. *Sci. Total. Environ.*, **637-638**, 954–970, doi:10.1016/j.scitotenv.2018.04.328.
- Hügel, S. and A.R. Davies, 2020: Public participation, engagement, and climate change adaptation: a review of the research literature. *WIREs Clim. Change*, **11**(4), e645, doi:10.1002/wcc.645.
- Huntington, H.P., et al., 2017: How small communities respond to environmental change: patterns from tropical to polar ecosystems. *Ecol. Soc.*, **22**(3), 9, doi:10.5751/ES-09171-220309.
- International Council for Science (ed.), 2017: *A Guide to SDG Interactions: from Science to Implementation*. International Council for Science, Paris, France. 236 pp.
- International Seabed Authority, *About ISA*. International Seabed Authority, Kingston, Jamaica, Accessed 31st December 2021, <https://www.isa.org.jm/about-isa>.
- IPBES, 2019: *Summary for Policymakers of the Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services* [Díaz, S., J. Settele, E. S. Brondizio, H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. Roy Chowdhury, Y.-J. Shin, I. J. Visserin-Hamakers, K. J. Willis and C. N. Zayas (eds.)]. IPBES Secretariat, Bonn, Germany. 56 pp.
- IPCC, 2014a: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field, C. B., V. R. Barros, D. J. Dokken, K. J. Mach, M. D. Mastrandrea, T. E. Bilir, M. Chatterjee, K. L. Ebi, Y. O. Estrada, R. C. Genova, B. Girma, E. S. Kissel, A. N. Levy, S. MacCracken, P. R. Mastrandrea and L. L. White (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, ISBN 978-1107058071. 1132 pp.
- IPCC, 2014b: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Barros, V. R., C. B. Field, D. J. Dokken, M. D. Mastrandrea, K. J. Mach, T. E. Bilir, M. Chatterjee, K. L. Ebi, Y. O. Estrada, R. C. Genova, B. Girma, E. S. Kissel, A. N. Levy, S. MacCracken, P. R. Mastrandrea and L. L. White (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, ISBN 978-1107058163. 688 pp.
- IPCC, 2019: Summary for Policymakers. In: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. [Pörtner, H.-O., D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. S. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama and N. Weyer (eds.)] (In press).
- Islam, M.M., et al., 2021: Transformational adaptations to climatic hazards: insights from mangroves-based coastal fisheries dependent communities of Bangladesh. *Mar. Policy*, **128**, 104475, doi:10.1016/j.marpol.2021.104475.
- Islam, M.M., S. Sallu, K. Hubacek and J. Paavola, 2014: Migrating to tackle climate variability and change? Insights from coastal fishing communities in Bangladesh. *Clim. Change*, **124**(4), 733–746, doi:10.1007/s10584-014-1135-y.
- Izumi, T., et al., 2019: Disaster risk reduction and innovations. *Prog. Disaster Sci.*, **2**, 100033, doi:10.1016/j.pdisas.2019.100033.
- Jacox, M.G., et al., 2020: Seasonal-to-interannual prediction of North American coastal marine ecosystems: forecast methods, mechanisms of predictability, and priority developments. *Prog. Oceanogr.*, **183**, 102307, doi:10.1016/j.pocean.2020.102307.
- Jacox, M.G., et al., 2019: Predicting the evolution of the 2014–2016 California Current system marine heatwave from an ensemble of coupled global climate forecasts. *Front. Mar. Sci.*, **6**, 497, doi:10.3389/fmars.2019.00497.
- Jang, Y.C., et al., 2014: Estimation of lost tourism revenue in Geoje Island from the 2011 marine debris pollution event in South Korea. *Mar. Pollut. Bull.*, **81**(1), 49–54, doi:10.1016/j.marpolbul.2014.02.021.
- Jansen, T. and H. Gislason, 2011: Temperature affects the timing of spawning and migration of North Sea mackerel. *Cont. Shelf Res.*, **31**(1), 64–72, doi:10.1016/j.csr.2010.11.003.
- Jarre, A., S.M. Ragaller and L. Hutchings, 2013: Long-term, ecosystem-scale changes in the southern Benguela marine pelagic social-ecological system: interaction of natural and human drivers. *Ecol. Soc.*, **18**(4), 55, doi:10.5751/ES-05917-180455.
- Johnson, D.E., M. A. Ferreira and C.B. Troján, 2021: *Regional Seas Biodiversity under the Post-2020 Global Biodiversity Framework*. [United Nations (ed.)]. UNEP Regional Seas Working Paper.Nairobi, Accessed 31st December 2021, [https://wedocs.unep.org/xmlui/bitstream/handle/20.500.11822/35102/RSB\\_Post2020GBF.pdf?sequence=3&isAllowed=y](https://wedocs.unep.org/xmlui/bitstream/handle/20.500.11822/35102/RSB_Post2020GBF.pdf?sequence=3&isAllowed=y).
- Johnson, N., et al., 2020: Knowledge co-production and co-management of Arctic wildlife. *Arct. Sci.*, **6**(3), 124–126, doi:10.1139/as-2020-0028.
- Jones, H.P., et al., 2018: Restoration and repair of Earth's damaged ecosystems. *Proc. R. Soc. B*, **285**(1873), 20172577, doi:10.1098/rspb.2017.2577.
- Jones, H.P., et al., 2020: Global hotspots for coastal ecosystem-based adaptation. *PLoS ONE*, **15**(5), e0233005, doi:10.1371/journal.pone.0233005.
- Jongman, B., 2018: Effective adaptation to rising flood risk. *Nat. Commun.*, **9**(1), 1986, doi:10.1038/s41467-018-04396-1.
- Juanes, F., S. Gephard and K.F. Beland, 2004: Long-term changes in migration timing of adult Atlantic salmon (*Salmo salar*) at the southern edge of the species distribution. *Can. J. Fish. Aquat. Sci.*, **61**(12), 2392–2400, doi:10.1139/f04-207.
- Kanamori, Y., A. Takasuka, S. Nishijima and H. Okamura, 2019: Climate change shifts the spawning ground northward and extends the spawning period of chub mackerel in the western North Pacific. *Mar. Ecol. Prog. Ser.*, **624**, 155–166, doi:10.3354/meps13037.
- Karp, M.A., et al., 2019: Accounting for shifting distributions and changing productivity in the development of scientific advice for fishery management. *ICES J. Mar. Sci.*, **76**(5), 1305–1315, doi:10.1093/icesjms/fsz048.
- Kawarazuka, N. and C. Béné, 2011: The potential role of small fish species in improving micronutrient deficiencies in developing countries: building evidence. *Public Health Nutr.*, **14**(11), 1927–1938, doi:10.1017/S1368980011000814.
- Kennedy, R.J. and W.W. Crozier, 2010: Evidence of changing migratory patterns of wild Atlantic salmon *Salmo salar* smolts in the River Bush, Northern Ireland, and possible associations with climate change. *J. Fish Biol.*, **76**(7), 1786–1805, doi:10.1111/j.1095-8649.2010.02617.x.
- Khan, A., A. Charles and D. Armitage, 2018a: Place-based or sector-based adaptation? A case study of municipal and fishery policy integration. *Clim. Policy*, **18**(1), 14–23, doi:10.1080/14693062.2016.1228520.
- Khan, F.N., A.M. Collins, P.K. Nayak and D. Armitage, 2018b: Women's perspectives of small-scale fisheries and environmental change in Chilika lagoon, India. *Marit. Stud.*, **17**(2), 145–154, doi:10.1007/s40152-018-0100-1.
- Kirezci, E., et al., 2020: Projections of global-scale extreme sea levels and resulting episodic coastal flooding over the 21st century. *Sci. Rep.*, **10**(1), 11629, doi:10.1038/s41598-020-67736-6.
- Kleisner, K., D. Zeller, R. Froese and D. Pauly, 2013: Using global catch data for inferences on the world's marine fisheries. *Fish Fish.*, **14**(3), 293–311, doi:10.1111/j.1467-2979.2012.00469.x.

- Kleypas, J., et al., 2021: Designing a blueprint for coral reef survival. *Biol. Conserv.*, **257**, 109107, doi:10.1016/j.biocon.2021.109107.
- Klöck, C. and P.D. Nunn, 2019: Adaptation to climate change in small island developing states: a systematic literature review of academic research. *J. Environ. Dev.*, **28**(2), 196–218, doi:10.1177/1070496519835895.
- Koenigstein, S., L.-H. Hentschel, L.C. Heel and C. Drinkorn, 2020: A game-based education approach for sustainable ocean development. *ICES J. Mar. Sci.*, **77**(5), 1629–1638, doi:10.1093/icesjms/fsaa035.
- Koschinsky, A., et al., 2018: Deep-sea mining: interdisciplinary research on potential environmental, legal, economic, and societal implications. *Integr. Environ. Assess. Manag.*, **14**(6), 672–691, doi:10.1002/ieam.4071.
- Kovach, R.P., S.C. Ellison, S. Pyare and D.A. Tallmon, 2015: Temporal patterns in adult salmon migration timing across southeast Alaska. *Glob. Change Biol.*, **21**(5), 1821–1833, doi:10.1111/gcb.12829.
- Kovach, R.P., A.J. Gharrett and D.A. Tallmon, 2012: Genetic change for earlier migration timing in a pink salmon population. *Proc. R. Soc. B.*, **279**(1743), 3870–3878, doi:10.1098/rspb.2012.1158.
- Kovach, R.P., et al., 2013: Earlier migration timing, decreasing phenotypic variation, and biocomplexity in multiple Salmonid species. *PLoS ONE*, **8**(1), e53807, doi:10.1371/journal.pone.0053807.
- Krelling, A.P., A.T. Williams and A. Turra, 2017: Differences in perception and reaction of tourist groups to beach marine debris that can influence a loss of tourism revenue in coastal areas. *Mar. Policy*, **85**, 87–99, doi:10.1016/j.marpol.2017.08.021.
- Kung, A., et al., 2021: Governing deep sea mining in the face of uncertainty. *J. Environ. Manag.*, **279**, 111593, doi:10.1016/j.jenvman.2020.111593.
- Kurekin, A.A., et al., 2019: Operational monitoring of illegal fishing in Ghana through exploitation of satellite Earth observation and AIS data. *Remote Sens.*, **11**(3), 293, doi:10.3390/rs11030293.
- Kurniawan, F., L. Adrianto, D.G. Bengen and L.B. Prasetyo, 2016: Vulnerability assessment of small islands to tourism: the case of the marine tourism park of the Gili Matra Islands, Indonesia. *Glob. Ecol. Conserv.*, **6**, 308–326, doi:10.1016/j.gecco.2016.04.001.
- Lachkar, Z., M. Lévy and K.S. Smith, 2019: Strong intensification of the Arabian Sea oxygen minimum zone in response to Arabian Gulf warming. *Geophys. Res. Lett.*, **46**(10), 5420–5429, doi:10.1029/2018GL081631.
- Laikre, L., M.K. Schwartz, R.S. Waples, N. Ryman and The GeM Working Group, 2010: Compromising genetic diversity in the wild: unmonitored large-scale release of plants and animals. *Trends Ecol. Evol.*, **25**(9), 520–529, doi:10.1016/j.tree.2010.06.013.
- Lam, V.W.Y., et al., 2020: Climate change, tropical fisheries and prospects for sustainable development. *Nat. Rev. Earth Environ.*, **1**(9), 440–454, doi:10.1038/s43017-020-0071-9.
- Langan, J.A., et al., 2021: Climate alters the migration phenology of coastal marine species. *Mar. Ecol. Prog. Ser.*, **660**, 1–18, doi:10.3354/meps13612.
- Larkin, D.J., R.J. Buck, J. Fieberg and S.M. Galatowitsch, 2019: Revisiting the benefits of active approaches for restoring damaged ecosystems. A comment on Jones HP et al., 2018 Restoration and repair of Earth's damaged ecosystems. *Proc. R. Soc. B.*, **286**(1907), 20182928, doi:10.1098/rspb.2018.2928.
- Lasso, A. and H. Dahles, 2018: Are tourism livelihoods sustainable? Tourism development and economic transformation on Komodo Island, Indonesia. *Asia Pac. J. Tour. Res.*, **23**(5), 473–485, doi:10.1080/10941665.2018.1467939.
- Le Blanc, D., C. Freire and M. Vierros, 2017: *Mapping the Linkages between Oceans and Other Sustainable Development Goals*. Working Papers, Vol. 149. UN Department of Economic and Social Affairs (DESA), New York, NY, USA. 34 pp. Available at: <https://doi.org/10.18356/3adc8369-en> (accessed 06/11/2020).
- Leal Filho, W., et al., 2018: Fostering coastal resilience to climate change vulnerability in Bangladesh, Brazil, Cameroon and Uruguay: a cross-country comparison. *Mitig. Adapt. Strateg. Glob. Change*, **23**(4), 579–602, doi:10.1007/s11027-017-9750-3.
- Leary, D., 2019: Agreeing to disagree on what we have or have not agreed on: the current state of play of the BBNJ negotiations on the status of marine genetic resources in areas beyond national jurisdiction. *Mar. Policy*, **99**, 21–29, doi:10.1016/j.marpol.2018.10.031.
- Leonardi, N., et al., 2018: Dynamic interactions between coastal storms and salt marshes: a review. *Geomorphology*, **301**, 92–107, doi:10.1016/j.geomorph.2017.11.001.
- Leslie, H.M. and K.L. McLeod, 2007: Confronting the challenges of implementing marine ecosystem-based management. *Front. Ecol. Environ.*, **5**(10), 540–548, doi:10.1890/060093.
- Levin, L.A., et al., 2009: Effects of natural and human-induced hypoxia on coastal benthos. *Biogeosciences*, **6**(10), 2063–2098, doi:10.5194/bg-6-2063-2009.
- Levin, L.A., et al., 2020: Climate change considerations are fundamental to management of deep-sea resource extraction. *Glob. Change Biol.*, **26**(9), 4664–4678, doi:10.1111/gcb.15223.
- Levin, P.S. and C. Möllmann, 2015: Marine ecosystem regime shifts: challenges and opportunities for ecosystem-based management. *Philos. Trans. Royal Soc. B Biol. Sci.*, **370**(1659), 20130275, doi:10.1098/rstb.2013.0275.
- Li, S. and S. Jay, 2020: Transboundary marine spatial planning across Europe: trends and priorities in nearly two decades of project work. *Mar. Policy*, **118**, 104012, doi:10.1016/j.marpol.2020.104012.
- Liu, G., et al., 2018: Predicting heat stress to inform reef management: NOAA coral reef watch's 4-month coral bleaching outlook. *Front. Mar. Sci.*, **5**, 57, doi:10.3389/fmars.2018.00057.
- Lodge, M.W., et al., 2007: *Recommended Best Practices for Regional Fisheries Management Organizations: Report of an Independent Panel to Develop a Model for Improved Governance by Regional Fisheries Management Organizations*. Chatham House, London, ISBN 978-1862031883. 141 pp.
- Logan, C.A., et al., 2021: Quantifying global potential for coral evolutionary response to climate change. *Nat. Clim. Change*, **11**, 537–542, doi:10.1038/s41558-021-01037-2.
- Lombardo, S.M., et al., 2019: Evidence for temperature-dependent shifts in spawning times of anadromous alewife (*Alosa pseudoharengus*) and blueback herring (*Alosa aestivalis*). *Can. J. Fish. Aquat. Sci.*, **77**(4), 741–751, doi:10.1139/cjfas-2019-0140.
- Long, R.D., A. Charles and R.L. Stephenson, 2015: Key principles of marine ecosystem-based management. *Mar. Policy*, **57**, 53–60, doi:10.1016/j.marpol.2015.01.013.
- Longhurst, A., S. Sathyendranath, T. Platt and C. Caverhill, 1995: An estimate of global primary production in the ocean from satellite radiometer data. *J. Plankton Res.*, **17**(6), 1245–1271, doi:10.1093/plankt/17.6.1245.
- Lotze, H.K., et al., 2019: Global ensemble projections reveal trophic amplification of ocean biomass declines with climate change. *Proc. Natl. Acad. Sci. U.S.A.*, **116**(26), 12907–12912, doi:10.1073/pnas.1900194116.
- Lovelock, C.E. and R. Reef, 2020: Variable impacts of climate change on blue carbon. *One Earth*, **3**(2), 195–211, doi:10.1016/j.oneear.2020.07.010.
- Lowe, B.S., et al., 2019: Adapting to change in inland fisheries: analysis from Lake Tanganyika, East Africa. *Reg. Environ. Change*, **19**(6), 1765–1776, doi:10.1007/s10113-019-01516-5.
- Lowerre-Barbieri, S.K., I. A. Catalán, A. Frugård Opdal and C. Jørgensen, 2019: Preparing for the future: integrating spatial ecology into ecosystem-based management. *ICES J. Mar. Sci.*, **76**(2), 467–476, doi:10.1093/icesjms/fsy209.
- Lubchenco, J. and K. Grorud-Colvert, 2015: Making waves: the science and politics of ocean protection. *Science*, **350**(6259), 382–383, doi:10.1126/science.aad5443.
- Luh, J., et al., 2017: Expert assessment of the resilience of drinking water and sanitation systems to climate-related hazards. *Sci. Total Environ.*, **592**, 334–344, doi:10.1016/j.scitotenv.2017.03.084.
- Mackas, D.L., R. Goldblatt and A.G. Lewis, 1998: Interdecadal variation in developmental timing of *Neocalanus plumchrus* populations at Ocean Station P in the subarctic North Pacific. *Can. J. Fish. Aquat. Sci.*, **55**(8), 1878–1893, doi:10.1139/f98-080.

- Macreadie, P.I., et al., 2019: The future of blue carbon science. *Nat. Commun.*, **10**(1), 3998, doi:10.1038/s41467-019-11693-w.
- Magnan, A.K., E.L.F. Schipper and V.K.E. Duvat, 2020: Frontiers in climate change adaptation science: advancing guidelines to design adaptation pathways. *Curr. Clim. Change Rep.*, **6**(4), 166–177, doi:10.1007/s40641-020-00166-8.
- Maharjan, A., et al., 2020: Migration and household adaptation in climate-sensitive hotspots in South Asia. *Curr. Clim. Change Rep.*, **6**(1), 1–16, doi:10.1007/s40641-020-00153-z.
- Mahon, R. and L. Fanning, 2019: Regional ocean governance: integrating and coordinating mechanisms for polycentric systems. *Mar. Policy*, **107**, 103589, doi:10.1016/j.marpol.2019.103589.
- Marzeion, B. and A. Levermann, 2014: Loss of cultural world heritage and currently inhabited places to sea-level rise. *Environ. Res. Lett.*, **9**(3), 034001, doi:10.1088/1748-9326/9/3/034001.
- Maslo, B., et al., 2019: Optimizing conservation benefits for threatened beach fauna following severe natural disturbances. *Sci. Total Environ.*, **649**, 661–671, doi:10.1016/j.scitotenv.2018.08.319.
- Mason, N., et al., 2020: Global opportunities and challenges for transboundary conservation. *Nat. Ecol. Evol.*, **4**(5), 694–701, doi:10.1038/s41559-020-1160-3.
- Masria, A., M. Iskander and A. Negm, 2015: Coastal protection measures, case study (Mediterranean zone, Egypt). *J. Coast. Conserv.*, **19**(3), 281–294, doi:10.1007/s11852-015-0389-5.
- Matich, P., K.B. Moore and J.D. Plumlee, 2020: Effects of Hurricane Harvey on the trophic status of juvenile sport fishes (*Cynoscion nebulosus*, *Sciaenops ocellatus*) in an estuarine nursery. *Estuaries Coasts*, **43**(5), 997–1012, doi:10.1007/s12237-020-00723-2.
- Maureaud, A., et al., 2021: Are we ready to track climate-driven shifts in marine species across international boundaries? A global survey of scientific bottom trawl data. *Glob. Change Biol.*, **27**(2), 220–236, doi:10.1111/gcb.15404.
- Maxwell, S.L., et al., 2020a: Area-based conservation in the twenty-first century. *Nature*, **586**(7828), 217–227, doi:10.1038/s41586-020-2773-z.
- Maxwell, S.M., K.M. Gjerde, M.G. Conners and L.B. Crowder, 2020b: Mobile protected areas for biodiversity on the high seas. *Science*, **367**(6475), 252–254, doi:10.1126/science.aaz9327.
- Mazaris, A.D., A.S. Kallimanis, S.P. Sgardelis and J.D. Pantis, 2008: Do long-term changes in sea surface temperature at the breeding areas affect the breeding dates and reproduction performance of Mediterranean loggerhead turtles? Implications for climate change. *J. Exp. Mar. Biol. Ecol.*, **367**(2), 219–226, doi:10.1016/j.jembe.2008.09.025.
- McGeady, R., C. Lordan and A.M. Power, 2021: Shift in the larval phenology of a marine ectotherm due to ocean warming with consequences for larval transport. *Limnol. Oceanogr.*, **66**(2), 543–557, doi:10.1002/lno.11622.
- McGinty, N., A.M. Power and M.P. Johnson, 2011: Variation among northeast Atlantic regions in the responses of zooplankton to climate change: not all areas follow the same path. *J. Exp. Mar. Biol. Ecol.*, **400**(1–2), 120–131, doi:10.1016/j.jembe.2011.02.013.
- McLeod, E., et al., 2019: The future of resilience-based management in coral reef ecosystems. *J. Environ. Manag.*, **233**, 291–301, doi:10.1016/j.jenvman.2018.11.034.
- McLeod, E., et al., 2018: Raising the voices of Pacific Island women to inform climate adaptation policies. *Mar. Policy*, **93**, 178–185, doi:10.1016/j.marpol.2018.03.011.
- McNamara, K.E., et al., 2020: An assessment of community-based adaptation initiatives in the Pacific Islands. *Nat. Clim. Change*, **10**(7), 628–639, doi:10.1038/s41558-020-0813-1.
- McQueen, K. and C.T. Marshall, 2017: Shifts in spawning phenology of cod linked to rising sea temperatures. *ICES J. Mar. Sci.*, **74**(6), 1561–1573, doi:10.1093/icesjms/fsx025.
- Melbourne-Thomas, J., et al., 2021: Poleward bound: adapting to climate-driven species redistribution. *Rev. Fish Biol. Fish.*, doi:10.1007/s11160-021-09641-3.
- Menéndez, P., et al., 2020: The global flood protection benefits of mangroves. *Sci. Rep.*, **10**(1), 4404, doi:10.1038/s41598-020-61136-6.
- Merryfield, W.J., et al., 2020: Current and emerging developments in subseasonal to decadal prediction. *Bull. Am. Meteorol. Soc.*, **101**(6), E869–E896, doi:10.1175/BAMS-D-19-0037.1.
- Michalena, E., et al., 2020: Promoting sustainable and inclusive oceans management in Pacific islands through women and science. *Mar. Pollut. Bull.*, **150**, 110711, doi:10.1016/j.marpolbul.2019.110711.
- Miladinova, S., A. Stips, E. Garcia-Gorriz and D. Macias Moy, 2017: Black Sea thermohaline properties: long-term trends and variations. *J. Geophys. Res. Oceans*, **122**(7), 5624–5644, doi:10.1002/2016JC012644.
- Miller, D.D., et al., 2018: Adaptation strategies to climate change in marine systems. *Glob. Change Biol.*, **24**(1), e1–e14, doi:10.1111/gcb.13829.
- Miller, K., et al., 2010: Climate change, uncertainty, and resilient fisheries: institutional responses through integrative science. *Prog. Oceanogr.*, **87**(1–4), 338–346, doi:10.1016/j.pocean.2010.09.014.
- Mo, Y., M.S. Kearney and R.E. Turner, 2020: The resilience of coastal marshes to hurricanes: the potential impact of excess nutrients. *Environ. Int.*, **138**, 105409, doi:10.1016/j.envint.2019.105409.
- Moe, B., et al., 2009: Climate change and phenological responses of two seabird species breeding in the high-Arctic. *Mar. Ecol. Prog. Ser.*, **393**, 235–246, doi:10.3354/meps08222.
- Mohamed Shaffril, H.A., et al., 2020: Systematic literature review on adaptation towards climate change impacts among indigenous people in the Asia Pacific regions. *J. Clean. Prod.*, **258**, 120595, doi:10.1016/j.jclepro.2020.120595.
- Monios, J. and G. Wilmsmeier, 2020: Deep adaptation to climate change in the maritime transport sector – a new paradigm for maritime economics? *Marit. Policy Manag.*, **47**(7), 853–872, doi:10.1080/03088839.2020.1752947.
- Monsinjon, J., et al., 2019: Effects of temperature and demography on the phenology of loggerhead sea turtles in Brazil. *Mar. Ecol. Prog. Ser.*, **623**, 209–219, doi:10.3354/meps12988.
- Moore, P.J., R.C. Thompson and S.J. Hawkins, 2011: Phenological changes in intertidal con-specific gastropods in response to climate warming. *Glob. Change Biol.*, **17**(2), 709–719, doi:10.1111/j.1365-2486.2010.02270.x.
- Morgan, M.J., P.J. Wright and R.M. Rideout, 2013: Effect of age and temperature on spawning time in two gadoid species. *Fish. Res.*, **138**, 42–51, doi:10.1016/j.fishres.2012.02.019.
- Morita, K., 2018: Earlier migration timing of salmonids: an adaptation to climate change or maladaptation to the fishery? *Can. J. Fish. Aquat. Sci.*, **76**(3), 475–479, doi:10.1139/cjfas-2018-0078.
- Morris, R.L., et al., 2020: Key principles for managing recovery of kelp forests through restoration. *BioScience*, **70**(8), 688–698, doi:10.1093/biosci/biaa058.
- Moser, S.C., J.A. Ekstrom, J. Kim and S. Heitsch, 2019: Adaptation finance archetypes: local governments' persistent challenges of funding adaptation to climate change and ways to overcome them. *Ecol. Soc.*, **24**(2), 28, doi:10.5751/ES-10980-240228.
- Muhling, B.A., et al., 2018: Regional-scale surface temperature variability allows prediction of Pacific bluefin tuna recruitment. *ICES J. Mar. Sci.*, **75**(4), 1341–1352, doi:10.1093/icesjms/fsy017.
- Naidoo, R., et al., 2019: Evaluating the impacts of protected areas on human well-being across the developing world. *Sci. Adv.*, **5**(4), eaav3006, doi:10.1126/sciadv.aav3006.
- Nalau, J., et al., 2018: The role of Indigenous and traditional knowledge in ecosystem-based adaptation: a review of the literature and case studies from the Pacific Islands. *Weather Clim. Soc.*, **10**(4), 851–865, doi:10.1175/WCAS-D-18-0032.1.
- National Academies of Sciences, 2019: *A Research Review of Interventions to Increase the Persistence and Resilience of Coral Reefs*. The National Academies Press, Washington, D.C. 258 pp.
- National Academies of Sciences Engineering and Medicine, 2019: *A Decision Framework for Interventions to Increase the Persistence and Resilience of Coral Reefs*. The National Academies Press, Washington, D.C.

- Neijnens, F.K., K. Siteur, J. van de Koppel and M. Rietkerk, 2021: Early warning signals for rate-induced critical transitions in salt marsh ecosystems. *Ecosystems*, **24**, 1825–1836, doi:10.1007/s10021-021-00610-2.
- Nicholls, R.J., 2018: Adapting to sea-level rise. In: *Resilience: The Science of Adaptation to Climate Change* [Zommers, Z. and K. Alverson(eds.)]. Elsevier, Amsterdam, Netherlands; Oxford, UK; Cambridge, USA, pp. 13–29. ISBN 978-0128118917.
- Nichols, C.R., et al., 2019: Collaborative science to enhance coastal resilience and adaptation. *Front. Mar. Sci.*, **6**, 404, doi:10.3389/fmars.2019.00404.
- Nichols, G., I. Lake and C. Heavyside, 2018: Climate change and water-related infectious diseases. *Atmosphere*, **9**(10), 385, doi:10.3390/atmos9100385.
- Nickols, K.J., et al., 2019: Setting ecological expectations for adaptive management of marine protected areas. *J. Appl. Ecol.*, **56**(10), 2376–2385, doi:10.1111/1365-2664.13463.
- Nilsson, M., et al., 2018: Mapping interactions between the sustainable development goals: lessons learned and ways forward. *Sustain. Sci.*, **13**(6), 1489–1503, doi:10.1007/s11625-018-0604-z.
- Nilsson, M., D. Griggs and M. Visbeck, 2016: Policy: map the interactions between sustainable development goals. *Nature*, **534**(7607), 320–322, doi:10.1038/534320a.
- Niner, H.J., et al., 2018: Deep-sea mining with no net loss of biodiversity—an impossible aim. *Front. Mar. Sci.*, **5**, 53, doi:10.3389/fmars.2018.00053.
- Norström, A.V., et al., 2020: Principles for knowledge co-production in sustainability research. *Nat. Sustain.*, **3**(3), 182–190, doi:10.1038/s41893-019-0448-2.
- Northrop, E., M. Konar, N. Frost and E. Hollaway, 2020: *A Sustainable and Equitable Blue Recovery to the COVID-19 Crisis*. [Secretariat of the High Level Panel for a Sustainable Ocean Economy (ed.)]. World Resources Institute, Accessed 31st December 2021, <http://www.oceanpanel.org/bluererecovery>.
- Nurhidayah, L. and A. McIlgorm, 2019: Coastal adaptation laws and the social justice of policies to address sea level rise: an Indonesian insight. *Ocean Coast. Manag.*, **171**, 11–18, doi:10.1016/j.ocemoaman.2019.01.011.
- Nursey-Bray, M., P. Fidelman and M. Owusu, 2018: Does co-management facilitate adaptive capacity in times of environmental change? Insights from fisheries in Australia. *Mar. Policy*, **96**, 72–80, doi:10.1016/j.marpol.2018.07.016.
- Obura, D.O., et al., 2021: Integrate biodiversity targets from local to global levels. *Science*, **373**(6556), 746–748, doi:10.1126/science.abb2234.
- Ogier, E., et al., 2020: Responding to climate change: participatory evaluation of adaptation options for key marine fisheries in Australia's south east. *Front. Mar. Sci.*, **7**, 97, doi:10.3389/fmars.2020.00097.
- Ojea, E., S.E. Lester and D. Salgueiro-Otero, 2020: Adaptation of fishing communities to climate-driven shifts in target species. *One Earth*, **2**(6), 544–556, doi:10.1016/j.oneear.2020.05.012.
- Ojea, E., I. Pearlman, S.D. Gaines and S.E. Lester, 2017: Fisheries regulatory regimes and resilience to climate change. *Ambio*, **46**(4), 399–412, doi:10.1007/s13280-016-0850-1.
- Oppenheimer, M., B.C. Glavovic , J. Hinkel, R. van de Wal, A.K. Magnan, A. Abd-Elgawad, R. Cai, M. Cifuentes-Jara, R.M. DeConto, T. Ghosh, J. Hay, F. Isla, B. Marzeion, B. Meyssignac, and Z. Sebesvari, et al., 2019: Sea Level Rise and Implications for Low Lying Islands, Coasts and Communities. In: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. [Pörtner, H.-O., D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. S. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama and N. Weyer (eds.)] (In press).
- Orejas, C., et al., 2020: Towards a common approach to the assessment of the environmental status of deep-sea ecosystems in areas beyond national jurisdiction. *Mar. Policy*, **121**, 104182, doi:10.1016/j.marpol.2020.104182.
- Oremus, K.L., et al., 2020: Governance challenges for tropical nations losing fish species due to climate change. *Nat. Sustain.*, **3**(4), 277–280, doi:10.1038/s41893-020-0476-y.
- Osland, M.J., et al., 2020: A tropical cyclone-induced ecological regime shift: mangrove forest conversion to mudflat in Everglades National Park (Florida, USA). *Wetlands*, **40**(5), 1445–1458, doi:10.1007/s13157-020-01291-8.
- OSPAR Commission, 2019: *2018 Status Report on the OSPAR Network of Marine Protected Areas*. Biodiversity and Ecosystem Series.OSPAR Commission, London, UK, Accessed 31st December 2021, <https://oap.ospar.org/en/ospar-assessments/committee-assessments/biodiversity-committee/status-ospar-network-marine-protected-areas/assessment-reports-mpa/2018/>. (73 pp).
- Otero, J., et al., 2014: Basin-scale phenology and effects of climate variability on global timing of initial seaward migration of Atlantic salmon (*Salmo salar*). *Glob. Change Biol.*, **20**(1), 61–75, doi:10.1111/gcb.12363.
- Ouyang, X. and S.Y. Lee, 2020: Improved estimates on global carbon stock and carbon pools in tidal wetlands. *Nat. Commun.*, **11**(1), 317, doi:10.1038/s41467-019-14120-2.
- Ovando, D., et al., 2021: Assessing the population-level conservation effects of marine protected areas. *Conserv. Biol.*, **35**(6), 1861–1870, doi:10.1111/cobi.13782.
- Owen, G., 2020: What makes climate change adaptation effective? A systematic review of the literature. *Glob. Environ. Change*, **62**, 102071, doi:10.1016/j.gloenvcha.2020.102071.
- Palacios-Abrantes, J., G. Reygondeau, C.C.C. Wabnitz and W.W.L. Cheung, 2020: The transboundary nature of the world's exploited marine species. *Sci. Rep.*, **10**(1), 17668, doi:10.1038/s41598-020-74644-2.
- Parmesan, C. and G. Yohe, 2003: A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, **421**(6918), 37–42, doi:10.1038/nature01286.
- Partelow, S., et al., 2020: Environmental governance theories: a review and application to coastal systems. *Ecol. Soc.*, **25**(4), 19, doi:10.5751/ES-12067-250419.
- Pasquini, L. and R.M. Cowling, 2015: Opportunities and challenges for mainstreaming ecosystem-based adaptation in local government: evidence from the Western Cape, South Africa. *Environ. Dev. Sustain.*, **17**(5), 1121–1140, doi:10.1007/s10668-014-9594-x.
- Paulus, C.A., M.R. Pellokila, Y.U.L. Sobang and E. Azmanajaya, 2019: The alternative livelihood development strategy in order to improve local fishermen revenue in the border region of Indonesia and Timor Leste. *Aquac. Aquar. Conserv. Legis.*, **12**(1), 269.
- Payne, M.R., et al., 2016: Uncertainties in projecting climate-change impacts in marine ecosystems. *ICES J. Mar. Sci.*, **73**(5), 1272–1282, doi:10.1093/icesjms/fsv231.
- Payne, M.R., et al., 2017: Lessons from the first generation of marine ecological forecast products. *Front. Mar. Sci.*, **4**, 289, doi:10.3389/fmars.2017.00289.
- Peck, M. A. and J.K. Pinnegar, 2018: Chapter 5: Climate change impacts, vulnerabilities and adaptations: North Atlantic and Arctic marine fisheries. In: *Impacts of Climate Change on Fisheries and Aquaculture: Synthesis of Current Knowledge, adaptation and Mitigation Options* [Barange, M., et al.(eds.)]. Food and Agriculture Organisation of the United Nations, Rome, Italy, pp. 87–112. ISBN 978-9251306079.
- Pecl, G.T., et al., 2017: Biodiversity redistribution under climate change: impacts on ecosystems and human well-being. *Science*, **355**(6332), eaai9214, doi:10.1126/science.aai9214.
- Peer, A.C. and T.J. Miller, 2014: Climate change, migration phenology, and fisheries management interact with unanticipated consequences. *N. Am. J. Fish. Manag.*, **34**(1), 94–110, doi:10.1080/02755947.2013.847877.
- Peñaherrera-Palma, C., et al., 2018: Evaluating abundance trends of iconic species using local ecological knowledge. *Biol. Conserv.*, **225**, 197–207, doi:10.1016/j.biocon.2018.07.004.
- Pennino, M.G., et al., 2021: The missing layers: integrating sociocultural values into marine spatial planning. *Front. Mar. Sci.*, **8**(848), 633198, doi:10.3389/fmars.2021.633198.
- Pentz, B., N. Klenk, S. Ogle and J.A.D. Fisher, 2018: Can regional fisheries management organizations (RFMOs) manage resources effectively during climate change? *Mar. Policy*, **92**, 13–20, doi:10.1016/j.marpol.2018.01.011.

- Perry, J., 2010: Pacific islanders pay heavy price for abandoning traditional diet. *Bull. World Health Organ.*, **88**, 484–485, doi:10.2471/BLT.10.010710.
- Persson, J., et al., 2012: Scale-dependent effects of climate on two copepod species, *Calanus glacialis* and *Pseudocalanus minutus*, in an Arctic-boreal sea. *Mar. Ecol. Prog. Ser.*, **468**, 71–83, doi:10.3354/meps09944.
- Pethybridge, H.R., et al., 2020: Contrasting futures for Australia's fisheries stocks under IPCC RCP8.5 emissions – a multi-ecosystem model approach. *Front. Mar. Sci.*, **7**(846), 577964, doi:10.3389/fmars.2020.577964.
- Petzold, J., et al., 2020: Indigenous knowledge on climate change adaptation: a global evidence map of academic literature. *Environ. Res. Lett.*, **15**, 113007, doi:10.1088/1748-9326/abb330.
- Pham, T.T., 2020: Tourism in marine protected areas: Can it be considered as an alternative livelihood for local communities? *Mar. Policy*, **115**, 103891, doi:10.1016/j.marpol.2020.103891.
- Philippart, C.J.M., et al., 2003: Climate-related changes in recruitment of the bivalve *Macoma balthica*. *Limnol. Oceanogr.*, **48**(6), 2171–2185, doi:10.4319/lo.2003.48.6.2171.
- Philips, E.J., S. Badylak, N.G. Nelson and K.E. Havens, 2020: Hurricanes, El Niño and harmful algal blooms in two sub-tropical Florida estuaries: direct and indirect impacts. *Sci. Rep.*, **10**(1), 1910, doi:10.1038/s41598-020-58771-4.
- Pinsky, M.L., 2021: Diversification spins a heatwave safety net for fisheries. *Proc. Natl. Acad. Sci. U.S.A.*, **118**(3), e2024412118, doi:10.1073/pnas.2024412118.
- Pinsky, M.L., et al., 2019: Greater vulnerability to warming of marine versus terrestrial ectotherms. *Nature*, **569**(7754), 108–111, doi:10.1038/s41586-019-1132-4.
- Pinsky, M.L., et al., 2021: Fish and fisheries in hot water: What is happening and how do we adapt? *Popul. Ecol.*, **63**(1), 17–26, doi:10.1002/1438-390X.12050.
- Pinsky, M.L. and M. Fogarty, 2012: Lagged social-ecological responses to climate and range shifts in fisheries. *Clim. Change*, **115**(3), 883–891, doi:10.1007/s10584-012-0599-x.
- Pinsky, M.L., et al., 2018: Preparing ocean governance for species on the move. *Science*, **360**(6394), 1189–1191, doi:10.1126/science.aat2360.
- Pinto, C.A., T.M. Silveira and S.B. Teixeira, 2020: Beach nourishment practice in mainland Portugal (1950–2017): overview and retrospective. *Ocean Coast. Manag.*, **192**, 105211, doi:10.1016/j.ocecoaman.2020.105211.
- Pinto, P.J., G.M. Kondolf and P.L.R. Wong, 2018: Adapting to sea level rise: emerging governance issues in the San Francisco Bay Region. *Environ. Sci. Policy*, **90**, 28–37, doi:10.1016/j.envsci.2018.09.015.
- Poloczanska, E.S., et al., 2013: Global imprint of climate change on marine life. *Nat. Clim. Change*, **3**(10), 919–925, doi:10.1038/nclimate1958.
- Queirós, A.M., J. Fernandes, L. Genevier and C.P. Lynam, 2018: Climate change alters fish community size-structure, requiring adaptive policy targets. *Fish Fish.*, **19**(4), 613–621, doi:10.1111/faf.12278.
- Queirós, A.M., et al., 2016: Solutions for ecosystem-level protection of ocean systems under climate change. *Glob. Change Biol.*, **22**(12), 3927–3936, doi:10.1111/gcb.13423.
- Quinn, T., et al., 2019: How local water and waterbody meanings shape flood risk perception and risk management preferences. *Sustain. Sci.*, **14**(3), 565–578, doi:10.1007/s11625-019-00665-0.
- Quinn, T.P. and D.J. Adams, 1996: Environmental changes affecting the migratory timing of American shad and sockeye salmon. *Ecology*, **77**(4), 1151–1162, doi:10.2307/2265584.
- Racault, M.-F., et al., 2019: Environmental reservoirs of *Vibrio cholerae*: challenges and opportunities for ocean-color remote sensing. *Remote Sens.*, **11**(23), 2763, doi:10.3390/rs11232763.
- Räisänen, J., 2017: Future climate change in the Baltic Sea region and environmental impacts. In: *Oxford Research Encyclopedia of Climate Science*. Oxford University Press, doi:10.1093/acrefore/9780190228620.013.634.
- Ramp, C., et al., 2015: Adapting to a warmer ocean—seasonal shift of baleen whale movements over three decades. *PLoS ONE*, **10**(3), e0121374, doi:10.1371/journal.pone.0121374.
- Rangel Soares, L.C., et al., 2002: Inequities in access to and use of drinking water services in Latin America and the Caribbean. *Pan Am. J. Public Health*, **11**(5/6), 386–396.
- Raymond-Yakoubian, J. and R. Daniel, 2018: An Indigenous approach to ocean planning and policy in the Bering Strait region of Alaska. *Mar. Policy*, **97**, 101–108, doi:10.1016/j.marpol.2018.08.028.
- Recha, C.W., M.N. Mukopi and J.O. Otieno, 2015: Socio-economic determinants of adoption of rainwater harvesting and conservation techniques in semi-arid Tharaka sub-county, Kenya. *Land Degrad. Dev.*, **26**(7), 765–773, doi:10.1002/ldr.2326.
- Reddin, C.J., et al., 2020: Marine clade sensitivities to climate change conform across timescales. *Nat. Clim. Change*, **10**(3), 249–253, doi:10.1038/s41558-020-0690-7.
- Reguero, B.G., et al., 2018: Coral reefs for coastal protection: a new methodological approach and engineering case study in Grenada. *J. Environ. Manag.*, **210**, 146–161, doi:10.1016/j.jenvman.2018.01.024.
- Reguero, B.G., et al., 2020: Financing coastal resilience by combining nature-based risk reduction with insurance. *Ecol. Econ.*, **169**, 106487, doi:10.1016/j.ecolecon.2019.106487.
- Reimann, L., et al., 2018: Mediterranean UNESCO World Heritage at risk from coastal flooding and erosion due to sea-level rise. *Nat. Commun.*, **9**(1), 4161, doi:10.1038/s41467-018-06645-9.
- Reum, J.C.P., et al., 2020: Ensemble projections of future climate change impacts on the eastern bering sea food web using a multispecies size spectrum model. *Front. Mar. Sci.*, **7**, 124, doi:10.3389/fmars.2020.00124.
- Rilov, G., et al., 2020: A fast-moving target: achieving marine conservation goals under shifting climate and policies. *Ecol. Appl.*, **30**(1), e02009, doi:10.1002/eap.2009.
- Rilov, G., et al., 2019: Adaptive marine conservation planning in the face of climate change: What can we learn from physiological, ecological and genetic studies? *Glob. Ecol. Conserv.*, **17**, e00566, doi:10.1016/j.gecco.2019.e00566.
- Rinkevich, B., 2021: Augmenting coral adaptation to climate change via coral gardening (the nursery phase). *J. Environ. Manag.*, **291**, 112727, doi:10.1016/j.jenvman.2021.112727.
- Robards, M.D. and T.P. Quinn, 2002: The migratory timing of adult summer-run steelhead in the Columbia River over six decades of environmental change. *Trans. Am. Fish. Soc.*, **131**(3), 523–536, doi:10.1577/1548-8659(2002)131<0523:TMTOAS>2.0.CO;2.
- Roberts, C.M., et al., 2017: Marine reserves can mitigate and promote adaptation to climate change. *Proc. Natl. Acad. Sci. U.S.A.*, **114**(24), 6167–6175, doi:10.1073/pnas.1701262114.
- Roberts, K.E., R.S. Valkan and C.N. Cook, 2018: Measuring progress in marine protection: a new set of metrics to evaluate the strength of marine protected area networks. *Biol. Conserv.*, **219**, 20–27, doi:10.1016/j.biocon.2018.01.004.
- Robinson, J.P.W., et al., 2020: Diversification insulates fisher catch and revenue in heavily exploited tropical fisheries. *Sci. Adv.*, **6**(8), eaaz587, doi:10.1126/sciadv.aaz0587.
- Rogers, L.A., et al., 2019: Shifting habitats expose fishing communities to risk under climate change. *Nat. Clim. Change*, **9**(7), 512–516, doi:10.1038/s41558-019-0503-z.
- Rosendo, S., L. Celliers and M. Mechisso, 2018: Doing more with the same: a reality-check on the ability of local government to implement Integrated Coastal Management for climate change adaptation. *Mar. Policy*, **87**, 29–39, doi:10.1016/j.marpol.2017.10.001.
- Ross, H., D.S. Adhuri, A.Y. Abdurrahim and A. Phelan, 2019: Opportunities in community-government cooperation to maintain marine ecosystem services in the Asia-Pacific and Oceania. *Ecosyst. Serv.*, **38**, 100969, doi:10.1016/j.ecoser.2019.100969.
- Rubenstein, M. A., R. Christophersen and J.I. Ransom, 2019: Trophic implications of a phenological paradigm shift: bald eagles and salmon in a changing climate. *J. Appl. Ecol.*, **56**(3), 769–778, doi:10.1111/1365-2664.13286.

- Rubio, I., U. Ganzedo, A.J. Hobday and E. Ojea, 2020: Southward re-distribution of tropical tuna fisheries activity can be explained by technological and management change. *Fish Fish.*, **21**(3), 511–521, doi:10.1111/faf.12443.
- Rudolph, T.B., et al., 2020: A transition to sustainable ocean governance. *Nat. Commun.*, **11**(1), 3600, doi:10.1038/s41467-020-17410-2.
- Rumore, D., T. Schenk and L. Susskind, 2016: Role-play simulations for climate change adaptation education and engagement. *Nat. Clim. Change*, **6**(8), 745–750, doi:10.1038/nclimate3084.
- Russi, D., et al., 2016: *Socio-Economic Benefits of the EU Marine Protected Areas*. Institute for European Environmental Policy (IEEP) for DG Environment, London, UK and Brussels, Belgium, Accessed 31st December 2021, [http://minisites.ieep.eu/assets/2131/Socio\\_Economic\\_Benefits\\_of\\_EU\\_MPAs.pdf](http://minisites.ieep.eu/assets/2131/Socio_Economic_Benefits_of_EU_MPAs.pdf). (93 pp).
- Sainz, J.F., et al., 2019: Spatial planning of marine aquaculture under climate decadal variability: a case study for mussel farms in Southern California. *Front. Mar. Sci.*, **6**, 253, doi:10.3389/fmars.2019.00253.
- Sala, E., et al., 2021: Protecting the global ocean for biodiversity, food and climate. *Nature*, **592**(7854), 397–402, doi:10.1038/s41586-021-03371-z.
- Samhouri, J.F., et al., 2019: An ecosystem-based risk assessment for California fisheries co-developed by scientists, managers, and stakeholders. *Biol. Conserv.*, **231**, 103–121, doi:10.1016/j.biocon.2018.12.027.
- Samora-Arvela, A. F., et al., 2017: Green infrastructure, climate change and spatial planning. *J. Spat. Organ. Dyn.*, **5**(3), 176–188.
- Sauve, D., G. Divoky and V.L. Friesen, 2019: Phenotypic plasticity or evolutionary change? An examination of the phenological response of an arctic seabird to climate change. *Funct. Ecol.*, **33**(11), 2180–2190, doi:10.1111/1365-2435.13406.
- Scharfe, M. and K.H. Wiltshire, 2019: Modeling of intra-annual abundance distributions: constancy and variation in the phenology of marine phytoplankton species over five decades at Helgoland Roads (North Sea). *Ecol. Model.*, **404**, 46–60, doi:10.1016/j.ecolmodel.2019.01.001.
- Schartup, A.T., et al., 2019: Climate change and overfishing increase neurotoxicant in marine predators. *Nature*, **572**(7771), 648–650, doi:10.1038/s41586-019-1468-9.
- Schlingmann, A., et al., 2021: Global patterns of adaptation to climate change by Indigenous Peoples and local communities. A systematic review. *Curr. Opin. Environ. Sustain.*, **51**, 55–64, doi:10.1016/j.cosust.2021.03.002.
- Schlüter, A., K. Van Assche, A.-K. Hornidge and N. Väidianu, 2020: Land-sea interactions and coastal development: an evolutionary governance perspective. *Mar. Policy*, **112**, 103801, doi:10.1016/j.marpol.2019.103801.
- Schlüter, M.H., et al., 2010: Phenological shifts of three interacting zooplankton groups in relation to climate change. *Glob. Change Biol.*, **16**(11), 3144–3153, doi:10.1111/j.1365-2486.2010.02246.x.
- Schroeder, I.D., et al., 2009: Winter pre-conditioning of seabird phenology in the California Current. *Mar. Ecol. Prog. Ser.*, **393**, 211–223, doi:10.3354/meps08103.
- Schuhbauer, A., et al., 2017: How subsidies affect the economic viability of small-scale fisheries. *Mar. Policy*, **82**, 114–121, doi:10.1016/j.marpol.2017.05.013.
- Scobie, M., 2019a: Climate change, human rights, and migration. In: *Emerging Threats to Human Rights: Resources, Violence, and Deprivation of Citizenship* [Smith-Cannoy, H.(ed.)]. Temple University Press, USA.
- Scobie, M., 2019b: Global marine and ocean governance, and Caribbean SIDS. In: *Global Environmental Governance and Small States: Architectures and Agency in the Caribbean*. Edward Elgar Publishing, Cheltenham, UK, pp. 90–117. ISBN 978-1786437266.
- Scott, F., et al., 2020: Hydro-morphological characterization of coral reefs for wave runup prediction. *Front. Mar. Sci.*, **7**, 361, doi:10.3389/fmars.2020.00361.
- Seddon, N., et al., 2020: Understanding the value and limits of nature-based solutions to climate change and other global challenges. *Philos. Trans. Royal Soc. B Biol. Sci.*, **375**(1794), 20190120, doi:10.1098/rstb.2019.0120.
- Shaffril, H.A.M., A. Abu Samah and J.L. D'Silva, 2017: Adapting towards climate change impacts: strategies for small-scale fishermen in Malaysia. *Mar. Policy*, **81**, 196–201, doi:10.1016/j.marpol.2017.03.032.
- Shahidul Islam, M. and M. Tanaka, 2004: Impacts of pollution on coastal and marine ecosystems including coastal and marine fisheries and approach for management: a review and synthesis. *Mar. Pollut. Bull.*, **48**(7), 624–649, doi:10.1016/j.marpolbul.2003.12.004.
- Shelton, C., 2014: *Climate Change Adaptation in Fisheries and Aquaculture: Compilation of Initial Examples*. FAO Fisheries and Aquaculture Circular No. 1088. Food and Agriculture Organization of the United Nations, Rome, Italy, Accessed 31st December 2021, <http://www.fao.org/3/a-i3569e.pdf>. (34 pp).
- Shirvani, A., S.M.J. Nazemosadat and E. Kahya, 2015: Analyses of the Persian Gulf sea surface temperature: prediction and detection of climate change signals. *Arab. J. Geosci.*, **8**(4), 2121–2130, doi:10.1007/s12517-014-1278-1.
- Siders, A.R., M. Hino and K.J. Mach, 2019: The case for strategic and managed climate retreat. *Science*, **365**(6455), 761–763, doi:10.1126/science.aax8346.
- Silver, J.M., et al., 2019: Advancing coastal risk reduction science and implementation by accounting for climate, ecosystems, and people. *Front. Mar. Sci.*, **6**, 556, doi:10.3389/fmars.2019.00556.
- Singh, G.G., et al., 2018: A rapid assessment of co-benefits and trade-offs among sustainable development goals. *Mar. Policy*, **93**, 223–231, doi:10.1016/j.marpol.2017.05.030.
- Soares, M.B., M. Daly and S. Dessai, 2018: Assessing the value of seasonal climate forecasts for decision-making. *Wiley Interdiscip. Rev. Clim. Change*, **9**(4), e523, doi:10.1002/wcc.523.
- Spillman, C.M. and G.A. Smith, 2021: A new operational seasonal thermal stress prediction tool for coral reefs around Australia. *Front. Mar. Sci.*, **8**(808), 687833, doi:10.3389/fmars.2021.687833.
- Steiner, Z., A.V. Turchyn, E. Harpaz and J. Silverman, 2018: Water chemistry reveals a significant decline in coral calcification rates in the southern Red Sea. *Nat. Commun.*, **9**(1), 3615, doi:10.1038/s41467-018-06030-6.
- Stephens, G., R. Bell and J. Lawrence, 2018: Developing signals to trigger adaptation to sea-level rise. *Environ. Res. Lett.*, **13**(10), 104004, doi:10.1088/1748-9326/aadf96.
- Stephenson, R.L., et al., 2016: Integrating fishers' knowledge research in science and management. *ICES J. Mar. Sci.*, **73**(6), 1459–1465, doi:10.1093/icesjms/fsw025.
- Stewart, E., J. Dawson and M. Johnston, 2015: Risks and opportunities associated with change in the cruise tourism sector: community perspectives from Arctic Canada. *Polar J.*, **5**(2), 403–427, doi:10.1080/2154896X.2015.1082283.
- Stewart-Sinclair, P.J., K.S. Last, B.L. Payne and T.A. Wilding, 2020a: A global assessment of the vulnerability of shellfish aquaculture to climate change and ocean acidification. *Ecol. Evol.*, **10**(7), 3518–3534, doi:10.1002/ece3.6149.
- Stewart-Sinclair, P.J., et al., 2020b: Blue restoration – building confidence and overcoming barriers. *Front. Mar. Sci.*, **7**, 748, doi:10.3389/fmars.2020.541700.
- Sultana, P., et al., 2019: Transforming local natural resource conflicts to cooperation in a changing climate: Bangladesh and Nepal lessons. *Clim. Policy*, **19**(sup1), S94–S106, doi:10.1080/14693062.2018.1527678.
- Sumaila, U.R., et al., 2010: A bottom-up re-estimation of global fisheries subsidies. *J. Bioecon.*, **12**(3), 201–225, doi:10.1007/s10818-010-9091-8.
- Sumaila, U.R. and T.C. Tai, 2020: End overfishing and increase the resilience of the ocean to climate change. *Front. Mar. Sci.*, **7**, 523, doi:10.3389/fmars.2020.00523.
- Sumaila, U.R., et al., 2021: Financing a sustainable ocean economy. *Nat. Commun.*, **12**(1), 3259, doi:10.1038/s41467-021-23168-y.
- Sumby, J., M. Haward, E.A. Fulton and G.T. Pecl, 2021: Hot fish: the response to climate change by regional fisheries bodies. *Mar. Policy*, **123**, 104284, doi:10.1016/j.marpol.2020.104284.
- Takasaki, Y., 2016: Learning from disaster: community-based marine protected areas in Fiji. *Envir. Dev. Econ.*, **21**(1), 53–77, doi:10.1017/S1355770X15000108.

- Taylor, S.F.W., et al., 2021: The complex relationship between asset wealth, adaptation, and diversification in tropical fisheries. *Ocean Coast. Manag.*, **212**, 105808, doi:10.1016/j.ocecoaman.2021.105808.
- Taylor, S.G., 2008: Climate warming causes phenological shift in Pink Salmon, *Oncorhynchus gorbuscha*, behavior at Auke Creek, Alaska. *Glob. Change Biol.*, **14**(2), 229–235, doi:10.1111/j.1365-2486.2007.01494.x.
- Terra Stori, F., C.M. Peres, A. Turra and R.L. Pressey, 2019: Traditional ecological knowledge supports ecosystem-based management in disturbed coastal marine social-ecological systems. *Front. Mar. Sci.*, **6**, 571, doi:10.3389/fmars.2019.00571.
- Thaxton, W.C., J.C. Taylor and R.G. Asch, 2020: Climate-associated trends and variability in ichthyoplankton phenology from the longest continuous larval fish time series on the east coast of the United States. *Mar. Ecol. Prog. Ser.*, **650**, 269–287, doi:10.3354/meps13404.
- The BACC II Author Team, 2015: *Second Assessment of Climate Change for the Baltic Sea Basin*. Regional Climate Studies. Springer, Cham, ISBN 978-3319160054. 501 pp.
- Theuerkauf, S.J., et al., 2019: A global spatial analysis reveals where marine aquaculture can benefit nature and people. *PLoS ONE*, **14**(10), e0222282, doi:10.1371/journal.pone.0222282.
- Thiault, L., et al., 2019: Escaping the perfect storm of simultaneous climate change impacts on agriculture and marine fisheries. *Sci. Adv.*, **5**(11), eaaw9976, doi:10.1126/sciadv.aaw9976.
- Thiele, L.P., 2020: Nature 4.0: assisted evolution, de-extinction, and ecological restoration technologies. *Glob. Environ. Polit.*, **20**(3), 9–27, doi:10.1162/glep\_a\_00559.
- Thompson, M.S.A., et al., 2020: A feeding guild indicator to assess environmental change impacts on marine ecosystem structure and functioning. *J. Appl. Ecol.*, **57**(9), 1769–1781, doi:10.1111/1365-2664.13662.
- Tilot, V., et al., 2021: Traditional dimensions of seabed resource management in the context of deep sea mining in the Pacific: learning from the socio-ecological interconnectivity between island communities and the ocean realm. *Front. Mar. Sci.*, **8**, 257, doi:10.3389/fmars.2021.637938.
- Tittensor, D.P., et al., 2019: Integrating climate adaptation and biodiversity conservation in the global ocean. *Sci. Adv.*, **5**(11), eaay9969, doi:10.1126/sciadv.aay9969.
- Tittensor, D.P., et al., 2018: A protocol for the intercomparison of marine fishery and ecosystem models: Fish-MIP v1.0. *Geosci. Model Dev.*, **11**(4), 1421–1442, doi:10.5194/gmd-11-1421-2018.
- Toimil, A., et al., 2020: Addressing the challenges of climate change risks and adaptation in coastal areas: a review. *Coast. Eng.*, **156**, 103611, doi:10.1016/j.coastaleng.2019.103611.
- Tommasi, D., et al., 2017: Managing living marine resources in a dynamic environment: the role of seasonal to decadal climate forecasts. *Prog. Oceanogr.*, **152**, 15–49, doi:10.1016/j.pocean.2016.12.011.
- Tompkins, E.L., et al., 2020: Adapting to change: people and policies. In: *Deltas in the Anthropocene* [Nicholls, R.J., W.N. Adger, C.W. Hutton and S.E. Hanson(eds.)]. Springer International Publishing, Cham, pp. 201–222. ISBN 978-3030235178.
- Tunin-Ley, A., et al., 2009: Phytoplankton biodiversity and NW Mediterranean Sea warming: changes in the dinoflagellate genus *Ceratium* in the 20th century. *Mar. Ecol. Prog. Ser.*, **375**, 85–99, doi:10.3354/meps07730.
- UN, 2019: *The Sustainable Development Goals Report 2019*. United Nations, New York, NY, USA, Accessed 31st December 2021, <https://unstats.un.org/sdgs/report/2019/The-Sustainable-Development-Goals-Report-2019.pdf>. (60 pp).
- UNEP-WCMC, 2021: Chapter 3. In: *Protected Planet Report 2020* [UNEP-WCMC and IUCN(eds.)]. UNEP-WCMC and IUCN, Cambridge, UK; Gland, Switzerland.
- UNEP-WCMC and IUCN, 2020: *Protected Planet: The World Database on Protected Areas (WDPA)*. Cambridge, UK, Accessed 31st December 2021, <https://www.protectedplanet.net>.
- Unger, S., et al., 2016: *Achieving the Sustainable Development Goal for the Oceans*. IASS Policy Brief 1/2017. Institute for Advanced Sustainability Studies (IASS), Potsdam, Germany, Accessed 31st December 2021, [https://www.iass-potsdam.de/sites/default/files/files/policy\\_brief\\_1\\_2017\\_en\\_achieving\\_the\\_sdgs\\_for\\_oceans.pdf](https://www.iass-potsdam.de/sites/default/files/files/policy_brief_1_2017_en_achieving_the_sdgs_for_oceans.pdf). (11 pp).
- Van Coppenolle, R. and S. Temmerman, 2020: Identifying global hotspots where coastal wetland conservation can contribute to nature-based mitigation of coastal flood risks. *Glob. Planet. Change*, **187**, 103125, doi:10.1016/j.gloplacha.2020.103125.
- van der Voort, T., J. Quist, C. Pahl-Wostl and M. Haasnot, 2017: Envisioning robust climate change adaptation futures for coastal regions: a comparative evaluation of cases in three continents. *Mitig. Adapt. Strateg. Glob. Change*, **22**(3), 519–546, doi:10.1007/s11027-015-9686-4.
- van Oppen, M.J.H., J.K. Oliver, H.M. Putnam and R.D. Gates, 2015: Building coral reef resilience through assisted evolution. *Proc. Natl. Acad. Sci. U.S.A.*, **112**(8), 2307–2313, doi:10.1073/pnas.1422301112.
- van Walraven, L., et al., 2017: Long-term patterns in fish phenology in the western Dutch Wadden Sea in relation to climate change. *J. Sea Res.*, **127**, 173–181, doi:10.1016/j.seares.2017.04.001.
- Van Walraven, L., et al., 2015: Long-term patterns in 50 years of scyphomedusae catches in the western Dutch Wadden Sea in relation to climate change and eutrophication. *J. Plankton Res.*, **37**(1), 151–167, doi:10.1093/plankt/fbu088.
- Vezzulli, L., et al., 2012: Long-term effects of ocean warming on the prokaryotic community: evidence from the vibrios. *ISME J.*, **6**(1), 21–30, doi:10.1038/ismej.2011.89.
- Viitasalo, M., et al., 2015: Environmental impacts—marine ecosystems. In: *Second Assessment of Climate Change for the Baltic Sea Basin* [The BACC II Author Team (ed.)]. Springer International Publishing, Cham, pp. 363–380. ISBN 978-3319160061.
- Vikolainen, V., J. Flikweert, H. Bressers and K. Lulofs, 2017: Governance context for coastal innovations in England: the case of sandscaping in North Norfolk. *Ocean Coast. Manag.*, **145**, 82–93, doi:10.1016/j.ocecoaman.2017.05.012.
- Vladimirova, K. and D. Le Blanc, 2016: Exploring links between education and sustainable development goals through the lens of UN flagship reports. *Sustain. Dev.*, **24**(4), 254–271, doi:10.1002/sd.1626.
- Volkov, S. and V.I. Pozdnyakov, 2021: Effects of environmental conditions on spring arrival, the timing of nesting, and the reproductive effort of Ross's Gull, *Rhodostethia rosea*, in the Delta of Lena River, Yakutia. *Biol Bull Russ Acad Sci*, **48**, 1332–1341, doi:10.31857/S0044513421010104.
- von Haaren, C., J. Bug and J. Barkmann, 2019: Identification and evaluation of habitat development potentials. In: *Landscape Planning with Ecosystem Services: Theories and Methods for Application in Europe* [von Haaren, C., A.A. Lovett and C. Albert(eds.)]. Springer Netherlands, Dordrecht, pp. 267–276. ISBN 978-9402416817.
- Voorberg, W. and R. Van der Veer, 2020: Co-management as a successful strategy for marine conservation. *J. Mar. Sci. Eng.*, **8**(7), 491, doi:10.3390/jmse8070491.
- Wabnitz, C.C.C., et al., 2018: Climate change impacts on marine biodiversity, fisheries and society in the Arabian Gulf. *PLoS ONE*, **13**(5), e0194537, doi:10.1371/journal.pone.0194537.
- Wamsler, C., et al., 2016: Operationalizing ecosystem-based adaptation: harnessing ecosystem services to buffer communities against climate change. *Ecol. Soc.*, **21**(1), 31, doi:10.5751/ES-08266-210131.
- Wang, J., Z. Dai, X. Mei and S. Fagherazzi, 2020: Tropical cyclones significantly alleviate mega-deltaic erosion induced by high riverine flow. *Geophys. Res. Lett.*, **47**(19), e2020GL089065, doi:10.1029/2020GL089065.
- Wanless, S., M. Frederiksen, J. Walton and M.P. Harris, 2009: Long-term changes in breeding phenology at two seabird colonies in the western North Sea. *Ibis*, **151**(2), 274–285, doi:10.1111/j.1474-919X.2008.00906.x.
- Ware, D. and Z. Banhalmi-Zakar, 2020: Strategies for governments to help close the coastal adaptation funding gap. *Ocean Coast. Manag.*, **198**, 105223, doi:10.1016/j.ocecoaman.2020.105223.

- Weatherdon, L.V., et al., 2016: Observed and projected impacts of climate change on marine fisheries, aquaculture, coastal tourism, and human health: an update. *Front. Mar. Sci.*, **3**, 48, doi:10.3389/fmars.2016.00048.
- Weaver, P.P.E., D.S.M. Billett and C.L. Van Dover, 2018: Environmental risks of deep-sea mining. In: *Handbook on Marine Environment Protection: Science, Impacts and Sustainable Management* [Salomon, M. and T. Markus(eds.)]. Springer International Publishing, Cham, pp. 215–245. ISBN 978-3319601564.
- White, A.T., H.P. Vogt and T. Arin, 2000: Philippine coral reefs under threat: the economic losses caused by reef destruction. *Mar. Pollut. Bull.*, **40**(7), 598–605, doi:10.1016/S0025-326X(00)0022-9.
- White, T.D., et al., 2020: Tracking the response of industrial fishing fleets to large marine protected areas in the Pacific Ocean. *Conserv. Biol.*, **34**(6), 1571–1578, doi:10.1111/cobi.13584.
- Wilkinson, B.P., Y.G. Satgé, J.S. Lamb and P.G.R. Jodice, 2019: Tropical cyclones alter short-term activity patterns of a coastal seabird. *Mov. Ecol.*, **7**(1), 30, doi:10.1186/s40462-019-0178-0.
- Wilkinson, E., L. Schipper, C. Simonet and Z. Kubik, 2016: *Climate Change, Migration and the 2030 Agenda for Sustainable Development*. Overseas Development Institute, Accessed 31st December 2021, <https://odi.org/en/publications/climate-change-migration-and-the-2030-agenda-for-sustainable-development/>. (15 pp).
- Williams, D.S., et al., 2020: A method for enhancing capacity of local governance for climate change adaptation. *Earth's Future*, **8**(7), e2020EF001506, doi:10.1029/2020EF001506.
- Williams, J.E., et al., 2015: Climate change adaptation and restoration of western trout streams: opportunities and strategies. *Fisheries*, **40**(7), 304–317, doi:10.1080/03632415.2015.1049692.
- Wilson, K.L., D.P. Tittensor, B. Worm and H.K. Lotze, 2020: Incorporating climate change adaptation into marine protected area planning. *Glob. Change Biol.*, **26**(6), 3251–3267, doi:10.1111/gcb.15094.
- Wiltshire, K.H., et al., 2010: Helgoland Roads, North Sea: 45 years of change. *Estuaries Coasts*, **33**(2), 295–310, doi:10.1007/s12237-009-9228-y.
- Wiltshire, K.H. and B.F.J. Manly, 2004: The warming trend at Helgoland Roads, North Sea: phytoplankton response. *Helgol. Mar. Res.*, **58**(4), 269–273, doi:10.1007/s10152-004-0196-0.
- Winter, G., et al., 2020: Steps to develop early warning systems and future scenarios of storm wave-driven flooding along coral reef-lined coasts. *Front. Mar. Sci.*, **7**, 199, doi:10.3389/fmars.2020.00199.
- Wood, A.L., J.R.A. Butler, M. Sheaves and J. Wani, 2013: Sport fisheries: opportunities and challenges for diversifying coastal livelihoods in the Pacific. *Mar. Policy*, **42**, 305–314, doi:10.1016/j.marpol.2013.03.005.
- Woodruff, S.C., M. Mullin and M. Roy, 2020: Is coastal adaptation a public good? The financing implications of good characteristics in coastal adaptation. *J. Environ. Plan. Manag.*, **63**(12), 2082–2101, doi:10.1080/09640568.2019.1703656.
- Wright, G., 2015: Marine governance in an industrialised ocean: a case study of the emerging marine renewable energy industry. *Mar. Policy*, **52**, 77–84, doi:10.1016/j.marpol.2014.10.021.
- Wright, G., et al., 2019: Marine spatial planning in areas beyond national jurisdiction. *Mar. Policy*, **132**, 103384, doi:10.1016/j.marpol.2018.12.003.
- Xie, W., et al., 2017: Application of terrestrial laser scanner on tidal flat morphology at a typhoon event timescale. *Geomorphology*, **292**, 47–58, doi:10.1016/j.geomorph.2017.04.034.
- Young, T., et al., 2019: Adaptation strategies of coastal fishing communities as species shift poleward. *ICES J. Mar. Sci.*, **76**(1), 93–103, doi:10.1093/icesjms/fsy140.
- Zickgraf, C., 2021: Climate change, slow onset events and human mobility: reviewing the evidence. *Curr. Opin. Environ. Sustain.*, **50**, 21–30, doi:10.1016/j.cosust.2020.11.007.