

**Cross-Chapter Paper 4: Mediterranean Region
Supplementary Material**

Cross-Chapter Paper Leads: Elham Ali (Egypt), Wolfgang Cramer (France)

Cross-Chapter Paper Authors: Jofre Carnicer (Spain), Elena Georgopoulou (Greece), Nathalie Hilmi (Monaco), Gonéri Le Cozannet (France), Piero Lionello (Italy)

Cross-Chapter Paper Contributing Authors: Ahmed Abdelrehim (Egypt), Mine Cinar (USA), Islam Abou El-Magd (Egypt), Shekoofeh Farahmand (Iran), François Gemenne (Belgium), Lena Reimann (Germany), Alain Safa (France), Sergio Vicente-Serrano (Spain), Francesca Spagnuolo (Italy), Duygu Sevgi Sevilgen (Monaco), Samuel Somot (France), Rémi Thiéblemont (France), Cristina Tirado (USA), Yves Trambly (France)

Cross-Chapter Paper Review Editors: Karim Hilmi (Morocco), Marta Rivera-Ferre (Spain)

Cross-Chapter Paper Scientist: Duygu Sevgi Sevilgen (Monaco)

Date of Draft: 1 October 2021

Notes: TSU Compiled Version

Table of Contents

SMCCP4.1 Detection and Attribution of Climate Change Impacts in the Mediterranean Basin.....	2
SMCCP4.2 Projected Climate Risks	3
SMCCP4.3 Sustainable Development in the Mediterranean Basin	12
SMCCP4.4 Mediterranean Sea-level Projections	12
References	13

1 **SMCCP4.1 Detection and Attribution of Climate Change Impacts in the Mediterranean Basin**
 2
 3

4 **Table SMCCP4.1:** References supporting Figure CCP4.5 (Attribution of observed impacts of climate change in the
 5 Mediterranean region).

Impact	Supporting references
Thermal discomfort	Heatwaves are increasing due to climate change in the Mediterranean, and amplified in cities due to urbanization practices, ultimately increasing mortality and morbidity rates as well as energy consumption (<i>high agreement, robust evidence in the north, medium evidence in the south</i>) (WGI Chapters 10 and 12; WGII Chapters 9 and 13; Kuglitsch et al., 2010; Salvati et al., 2017; Zinzi and Carnielo, 2017; Pyrgou and Santamouris, 2018; Salameh et al., 2019; Maggiotto et al., 2021)
Pluvial and river flooding	There is a mixed signal of increasing and decreasing flood trends in the Mediterranean (Gaume et al., 2016; Blöschl et al., 2017; Gudmundsson et al., 2017; Kundzewicz et al., 2017; Siam and Eltahir, 2017), and there is <i>low confidence</i> in attribution to climate change due to the major impacts of human interventions such as land use change, groundwater exploitation, urbanization and non-optimal flood risk management (WGI Chapters 11 and 12 for extreme precipitation changes; Llasat et al., 2013; Mediero et al., 2014; Ziv et al., 2014; Baahmed et al., 2015; Polemio and Lonigro, 2015; Gaume et al., 2016; Llasat et al., 2016; Paprotny et al., 2018; Ribes et al., 2019; Trambly et al., 2019; Vicente-Serrano et al., 2019; Argaman et al., 2020) ()
Water availability and quality	There is <i>high confidence</i> in detection and attribution of climate change impacts on water availability in the North-Western Mediterranean, and <i>medium confidence</i> in other subregions. There is emerging evidence on the impacts to water quality. (WGI Chapters 8 and 11; WGII Chapters 9 and 13; Hoerling et al., 2012; Ruffault et al., 2013; Vicente-Serrano et al., 2014; Aguilera et al., 2015; Quintana-Seguí et al., 2016; Van Vliet et al., 2016; Gosling et al., 2017; Spinoni et al., 2017; Caloiero et al., 2018; Mathbout et al., 2018; Okkan and Kirdemir, 2018; Braca et al., 2019; Grillakis, 2019)
Wildfires	There is <i>medium confidence</i> that in Mediterranean Europe, wildfires are decreasing owing to good practices (Brotons et al., 2013; Turco et al., 2014; Turco et al., 2016), despite the increasing hazard caused by increased drought frequency and severity (WGI Chapter 11; Pausas and Fernández-Muñoz, 2012; Karali et al., 2014; Fernandes et al., 2016; Turco et al., 2017b; Ruffault et al., 2018). There is low confidence in any trends on wildfire for the south Mediterranean due to the lack of attribution studies, limited monitoring of direct human interventions and also limited fuel availability in the south east Mediterranean (Meddour-Sahar, 2015; Turco et al., 2017a; Curt et al., 2020).
Coastal flooding and erosion	Chronic flooding in Venice (Box 13.1) and generally Mediterranean extreme water levels during storms increase consistently with sea-level rise (Marcos et al., 2009). Flood and erosion hazards and risks have not been attributed to climate change yet, as they highly depend on coastal management strategies (WGI Chapter 13Frihy et al., 2010). Pocket beaches in the Mediterranean might be early responders to sea-level rise (Brunel and Sabatier, 2009). Some permanent flooding is observed in subsiding areas such as the region near Thessaloniki (Greece) (Raucoules et al., 2008).
Marine ecosystems	New evidence published since AR5 (AR5 WGII Chapters 5 and 18) confirms that a shift in Mediterranean marine ecosystems, characterized by biodiversity decline and invasive species, has occurred since the 1980s, with <i>high confidence</i> of a major climate change imprint superimposed with the impacts of human activities (Chapter 3; Fortibuoni et al., 2015; Stergiou et al., 2016; Corrales et al., 2017; Givan et al., 2018; Azzurro et al., 2019; Kim et al., 2019; Albano et al., 2021).
Inland ecosystems	New evidence since AR5 WGII Chapter 18 confirms that terrestrial and freshwater ecosystems are impacted by climate change in the Mediterranean, resulting in loss of habitats and biodiversity and range shifts (e.g. birds), including through cascading impacts such as drought and wildfires (<i>high confidence</i> in northern Mediterranean, <i>medium confidence</i> in Southern Mediterranean) (Chapters 9 and 13 and CCP1; Stefanescu et al., 2011; Peñuelas et al., 2018; Bartsch et al., 2020).

Fisheries	New evidence published since AR5 WGII Chapter 18 (Sabatés et al., 2006; Lejeusne et al., 2010; Sabatés et al., 2012) further confirms that climate warming has had a negative impact on marine exploited fish stocks (<i>high confidence</i>), superimposed on the impacts of human activities such as overfishing. The economic value of proliferating species is generally less than that of declining species (<i>high confidence</i>) (Coll et al., 2014; Tsikliras and Stergiou, 2014; Stergiou et al., 2016; Azzurro et al., 2019).
Agriculture and viticulture	Changes in seasonality are attributed to climate change and the drying trends affect agriculture negatively (<i>low confidence</i> in the south-east Mediterranean and <i>medium confidence</i> in other sub-regions) (El-Maayar and Lange, 2013; Garcia-Mozo et al., 2015; Moore and Lobell, 2015; Oteros et al., 2015; Seif-Ennasr et al., 2016; Di Lena et al., 2018).

1
2
3
4
5
6
7
8
9

SMCCP4.2 Projected Climate Risks

The following tables provide the quantitative basis and literature references for the “burning ember” illustrations in Figure CCP4.8.

Table SMCCP4.2a: Supporting material for water availability and quality

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
Undetectable to Moderate	0.8 – 1.0	High	<ul style="list-style-type: none"> • Increase of the fractional area under soil drought in the northern Mediterranean by 4-14% (Grillakis, 2019) • Reduction (median) in mean annual runoff in Spain by 5%, compared to 1980-2010 (Gosling et al., 2017) • Reduction of groundwater recharge in Italy by 7-12% from 1996-2015 (Braca et al., 2019) • Reduction of reservoir inflows in Turkey by 15% from 1980-2005 (Okkan and Kirdemir, 2018) • Change of gross hydropower potential by -5% up to +20% in the northern Mediterranean and by 5-60% in the southern Mediterranean, compared to 1971-2000 (Van Vliet et al., 2016) • Reduction of cooling water discharge capacity by 5-60%, from 1971-2000 (Van Vliet et al., 2016) • The Standard Precipitation index (SPI) displays decreasing trends in the Mediterranean Basin (trends less clear in southern France). The typical characteristics of droughts (12-24 months SPI) mean that they have relevance for water management (Caloiero et al., 2018). • The SPI (driven by precipitation) and SPEI (driven by precipitation and temperature) display remarkable drying trends in southern Europe, especially in summer and autumn during the last three decades, both in terms of frequency and intensity (Spinoni et al., 2017). • Increased severity of droughts in Spain attributed to higher evaporative demand due to rising temperatures (Vicente-Serrano et al., 2014) • Increase in drought intensity and duration as well as seasonality changes are detected in subregions of Mediterranean France (Ruffault et al., 2013) • The Mediterranean drying after 1970 during the winter season is partly attributed to anthropogenic climate change (Hoerling et al., 2012) • Identification of drought clusters in north-western Mediterranean in the 1940s since the 2000s, and in eastern Mediterranean since the 1980s (Quintana-Seguí et al., 2016) • A drying trend is reported in the Eastern Mediterranean (Mathbout et al., 2018)

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
			<ul style="list-style-type: none"> There is <i>medium to high confidence</i> that anthropogenic climate change increased drought severity in the Mediterranean (AR6 WGI Chapters 8 and 11)
Moderate to High	1.4 – 2.0	<i>High</i>	<ul style="list-style-type: none"> <i>High confidence</i> in increase of hydrological, agricultural and ecological droughts in the Mediterranean between 1.5 and 2°C of GWL (WGI Chapter 11). Aridity is projected to expand in the South Mediterranean (CCP3) Reduction of streamflow (90th percentile) by 12-16% in northern Mediterranean, compared to 1971-2000 (Marx et al., 2018) Reduction of average annual discharge in Spain by 14-37% from 1971-2000 (Marcos-Garcia et al., 2017) and by 13-14% from 2003-2013 in Italy (Pumo et al., 2016) Reduction (median) of mean annual runoff in Spain by 20% compared to 1980-2010 (Gosling et al., 2017) and of surface runoff in Turkey by 7-20% from 1970-2000 (Bucak et al., 2017) Reduction of low flows in France by 12% from 1980-2009 (Andrew and Sauquet, 2017) and by 15% or more over southern Europe (Roudier et al., 2016) Reduction of inflow in Spanish dams by 15-17% from to 1992-2011 (Zabalza-Martínez et al., 2018) Reduction of groundwater recharge in Italy by 10-18% from baseline (Masia et al., 2018; Braca et al., 2019) Reduction of reservoir inflows in Turkey by 21% from 1980-2005 (Okkan and Kirdemir, 2018) and of reservoir maximum water level by 14-31% from 1970-2000 (Bucak et al., 2017) Reduction of mean annual streamflow volume in Israel by 45-47% from 1996-2005 (Peleg et al., 2015) Reduction of water availability in southwestern Mediterranean by up to 40% from to 1976-2005 (Tramblay et al., 2018) Reduction of groundwater volume by 26-42% in a semi-arid catchment in Italy compared to baseline (Guyennon et al., 2017) and of groundwater availability in Greece by 12-18% from historic levels (Koutroulis et al., 2016) Reduction of hydropower production in Spain by 13-33% from baseline (Lobanova et al., 2016; Solaun and Cerdá, 2017), and by 10-20% on average over the entire Mediterranean (Turner et al., 2017) Reduction of gross hydropower potential by 60% or more from 1981-2000 (Van Vliet et al., 2016; Zhang et al., 2018) Reduction of available freshwater for cooling of thermal power plants by 6-18% from baseline in Spain and other northern Mediterranean locations (Payet-Burin et al., 2018; Tobin et al., 2018) In France, increase of area under risk of severe drought by 14-66% (EEA, 2017; Lozano et al., 2017; Peñuelas et al., 2017; Varela et al., 2019) Increase of the highest probability of four consecutive drought years by 20-82% compared to 1967-2016 (Lehner et al., 2017) Reduction of annual runoff by 9-37% (Schleussner et al., 2016; Sellami et al., 2016)
High to Very High	2.7-3.2	<i>High</i>	<p>Warming at 2.7-3.2°C:</p> <ul style="list-style-type: none"> Reduction of hydropower production in Spain by 55% from baseline (Lobanova et al., 2016; Solaun and Cerdá, 2017) Reduction (median) of mean annual runoff in Spain by 20% compared to 1980-2010 (Gosling et al., 2017) Reduction (median) in groundwater water availability in Greece by 27% (Koutroulis et al., 2016)

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
			<ul style="list-style-type: none"> Reduction of streamflow (90th percentile) by 35% in northern Mediterranean compared to 1971-2000 (Marx et al., 2018) In Turkey, reduction (median) of surface runoff by 52% and reservoir maximum water level by 75-78% from 1970-2000 (Bucak et al., 2017) <p>Warming higher than 3.5°C:</p> <ul style="list-style-type: none"> <i>Very likely</i> increase of hydrological, agricultural and ecological droughts in the Mediterranean at 4°C of GWL (WGI Chapter 11). In Italy, reduction of: a) groundwater volume by 57% in a semi-arid catchment compared to baseline (Guyennon et al., 2017), b) groundwater recharge by 42% from baseline (Braca et al., 2019), c) of average annual discharge by 32% from 2003-2013 (Pumo et al., 2016). Reduction of water availability in southwestern Mediterranean by up to 55% from to 1976-2005 (Tramblay et al., 2018) Reduction of annual runoff in Portugal by 40-70% from 1961-1990 (Mourato et al., 2015) Reduction of annual discharge in Tunisia by 41-58% from 1970-2000 (Dakhlaoui et al., 2020) Reduction of surface runoff in Morocco by 55-63% from 1989-2009 (Marchane et al., 2017) Increase of the highest probability of four consecutive drought years by 220% compared to 1967-2016 (Lehner et al., 2017) Reduction of gross hydropower potential by 20-60% in northern Mediterranean and by 40-60% or more in southern Mediterranean from 1981-2000 (Van Vliet et al., 2016) Reduction of hydropower production in Spain by 30-55% from baseline (Lobanova et al., 2016; Solaun and Cerdá, 2017) Reduction of cooling water discharge capacity by more than 60% from 1971-2000 (Van Vliet et al., 2016)

1
2
3

Table SMCCP4.2b: Supporting material for wildfires

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
Undetectable to Moderate	0.8 - 1.5	<i>Medium</i>	<ul style="list-style-type: none"> Change of burnt area in northern Mediterranean by -5% to +54% (Wu et al., 2015; Turco et al., 2018) In Italy, change of fire probability by -4.2% to +11.6%, of fire potential index by -5.5% to +11.8%, and of high flame length probability by -50% to +25% (EEA, 2017; Lozano et al., 2017; Peñuelas et al., 2017; Varela et al., 2019). Wildfire hazard increasing due to increased drought frequency and severity (Pausas and Fernández-Muñoz, 2012; Karali et al., 2014; Fernandes et al., 2016; Turco et al., 2017b; Ruffault et al., 2018), but human interventions remain key driver of wildfire risk (Meddour-Sahar, 2015; Turco et al., 2017a; Curt et al., 2020), decrease of wildfire risk in the northern Mediterranean (Brotons et al., 2013; Turco et al., 2014; Turco et al., 2016) Times of emergence of increased wildfire risk by the mid-21st century in the north-western Mediterranean (Lozano et al., 2017)
Moderate to High	1.6 - 2.5	<i>Medium</i>	<ul style="list-style-type: none"> Increase of areas prone to fire risk by 53% in Corsica (Garbolino et al., 2016). Area with 7 or more days with extreme fire weather increasing by 82-217% in southern France (EEA, 2017; Lozano et al., 2017; Peñuelas et al., 2017; Varela et al., 2019). Increase of fire risk in Turkey by 20% on average (Satir et al., 2016)

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
			<ul style="list-style-type: none"> Increase of the number of days in summer with Fire Weather Index (FWI) ≥ 15 by 20-50% in central-northern Italy and by 2-20% in southern Italy (Faggian, 2018). Increase of burnt area in northern Mediterranean by 62-87% (Turco et al., 2018)
High to Very High	3.0 - 3.7	Low	<ul style="list-style-type: none"> Increase of burnt area in northern Mediterranean by 37-187% (Wu et al., 2015; Turco et al., 2018)

1
2
3

Table SMCCP4.2c: Supporting material for inland ecosystems

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
Undetectable to Moderate	0.7-1.2	Medium	<ul style="list-style-type: none"> New evidence since AR5 WGII Chapter 18 confirms that terrestrial and freshwater ecosystems are impacted by climate change in the Mediterranean, resulting in loss of habitats and biodiversity and range shifts (e.g. birds), including through cascading impacts such as drought and wildfires (<i>high confidence</i> in northern Mediterranean, <i>medium confidence</i> in Southern Mediterranean) (Chapters 4, 9 and 13 and CCP1; Stefanescu et al., 2011; Peñuelas et al., 2018; Bartsch et al., 2020).
Moderate to High	1.5-2	High	<ul style="list-style-type: none"> Terrestrial ecosystems negatively impacted by drought and wildfires (Guiot and Cramer, 2016; Matías et al., 2019; Sánchez-Salguero et al., 2020), and freshwater species and ecosystems (including fishes, insects, molluscs) negatively affected by reduced river flow and alteration of the water quality Chapters 2 and 4, CCP1; See Tables SMCCP4.2a and SMCCP4.2b; Jarić et al., 2019; Lefebvre et al., 2019) Mediterranean island endemic species especially at risk due to low possibilities to migrate (CCP1).
High to Very High	2-3	Medium	<ul style="list-style-type: none"> Terrestrial species such as land plants, insects, birds, reptiles, and mammals are projected to be negatively affected, with a steep increase projected risks between 2 and 3°C of warming (CCP1; Manes et al., 2021) Arid conditions gaining space in the Mediterranean region, with e.g. 13-30% of Mediterranean Natura 2000 area and 15 to 23% of Natura 2000 sites projected to become arid above 3°C GWL (Barredo et al., 2016). Substantial loss of habitat in freshwaters, e.g., 25% (RCP4.5) and 40% (RCP8.5) for brown trout in 2100 (Santiago et al., 2016)

4
5
6

Table SMCCP4.2d: Supporting material for marine ecosystems

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
Undetectable to Moderate	0.5-1	High	<ul style="list-style-type: none"> New evidence published since AR5 (AR5 WGII Chapters 5 and 18) confirms that a shift in Mediterranean ecosystems is detected since the 1980s with <i>high confidence</i> of a climate change imprint and continued tropicalization (<i>high confidence</i>) and mass mortality events (<i>very high confidence</i>) (Chapter 3) (Fortibuoni et al., 2015; Stergiou et al., 2016; Corrales et al., 2017; Givan et al., 2018; Azzurro et al., 2019; Kim et al., 2019; Albano et al., 2021) New evidence published since (Sabatés et al., 2006; Lejeusne et al., 2010; Sabatés et al., 2012) and AR5 WGII Chapter 18 confirms that climate warming has a negative impact on marine exploited fish stocks (<i>high confidence</i>), superimposed on direct human impacts. The value of proliferating species is generally less than those of

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
			declining species (<i>high confidence</i>) (Coll et al., 2014; Tsikliras and Stergiou, 2014; Stergiou et al., 2016; Azzurro et al., 2019).
Moderate to High	1-1.5	High	<ul style="list-style-type: none"> Observed shift in marine ecosystems projected to continue and intensify, with e.g. loss of seagrass habitat (WGI Chapter 3), decrease of primary production and biomass in the West Mediterranean (Macias et al., 2015; Moullec et al., 2019), decreasing biodiversity of meso-zooplankton (Benedetti et al., 2018), continued tropicalization (<i>high confidence</i>) (Lloret et al., 2015) (Corrales et al., 2018) Mass mortality events due to marine heatwaves (<i>very high confidence</i>) (Chapter 3) and increased risks of extinctions as climate warms projected to continue (CCPI; Ben Rais Lasram et al., 2010; Garrabou et al., 2021; Manes et al., 2021)
High to Very High	1.5-2.5	High	<ul style="list-style-type: none"> Increased ecosystem shifts, mass mortality events and risks of extinction for endemic species between 1.5 and 2.5°C (Chapters 3 9 and 13; Garrabou et al., 2021; Manes et al., 2021) Loss of habitats and nesting areas for e.g. turtles due to beach losses induced by projected sea-level rise (WGI Chapter 3). 25% of endemic marine species in the Mediterranean projected to be at high risk of extinction due to climate change (range of projections covered: from 1.5°C to 3°C) (CCPI; Manes et al., 2021) Abundance of small to medium pelagic fish is projected to decline up to 33% by 2100 (Albouy et al., 2013; Burrows et al., 2014; Stergiou et al., 2016; Raybaud et al., 2017; Albano et al., 2021).

1
2
3

Table SMCCP4.2e: Supporting material for food production and security

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
Undetectable to Moderate	1.0 – 1.4	High	<ul style="list-style-type: none"> Decrease of wheat yield by 5% in Egypt, by 0-25% in Italy, by 18.8% in Morocco (Chourghal et al., 2016; Dettori et al., 2017; Iocola et al., 2017; Brouziyne et al., 2018; Kheir et al., 2019). Decrease of sunflower crop water productivity by 15.7% in Morocco (Chourghal et al., 2016; Dettori et al., 2017; Iocola et al., 2017; Brouziyne et al., 2018; Kheir et al., 2019). Decrease of olive yield by 17% in Portugal, by up to 8% in Sardinia-Sicily in Italy, and by 15-18% in Spain (Fraga et al., 2020).
Moderate to High	1.5 – 2.1	High	<ul style="list-style-type: none"> Decrease of wheat yield by 9% in Egypt, by 4.5-25% in Italy (Chourghal et al., 2016; Dettori et al., 2017; Iocola et al., 2017; Brouziyne et al., 2018; Kheir et al., 2019), by 5-55% in Algeria (Bouregaa, 2019; Cammarano et al., 2019). Decrease of wheat crop water productivity in Morocco by 21.7% (Chourghal et al., 2016; Dettori et al., 2017; Iocola et al., 2017; Brouziyne et al., 2018; Kheir et al., 2019). Decrease of maize yield by 20-29% in Italy (Chourghal et al., 2016; Dettori et al., 2017; Iocola et al., 2017; Brouziyne et al., 2018; Kheir et al., 2019), by 3-10% in Greece (Georgopoulou et al., 2017; Iocola et al., 2017) Decrease of barley yield by 18-25% in Algeria (Bouregaa, 2019; Cammarano et al., 2019), by up to 12% in Greece (Georgopoulou et al., 2017; Iocola et al., 2017), and by 27% on average in the entire Mediterranean (Bouregaa, 2019; Cammarano et al., 2019). Decrease of rice yield by 6.4% in France and by 19.6% in Italy (Bregaglio et al., 2017). Decrease of sunflower crop water productivity by 44.7% in Morocco (Chourghal et al., 2016; Dettori et al., 2017; Iocola et al.,

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
			<p>2017; Brouziyne et al., 2018; Kheir et al., 2019). Decrease of sunflower yield by 65% in Greece (Georgopoulou et al., 2017; Iocola et al., 2017).</p> <ul style="list-style-type: none"> • Decrease of olive yield by 20% in Portugal (Fraga et al., 2020), by 3-8% in Italy (Fraga et al., 2020), by 19-21% in Spain (Fraga et al., 2020), and by 10-60% in Algeria (Bouregaa, 2019; Cammarano et al., 2019). Decrease of net primary production of olive groves by 0.02-11% in Italy (Brilli et al., 2019; Fraga et al., 2020). • Decrease of potato yield by up to -45% in northern Mediterranean (Zhao et al., 2015; Georgopoulou et al., 2017). • Decrease of various vegetable yields by 4-34% in Greece (Georgopoulou et al., 2017; Iocola et al., 2017). • Decrease of tomato yield by -19-81% over the entire Mediterranean (Saadi et al., 2015). • Apples at maturity in northeastern Spain could be of inferior quality, and chilling requirements may be unsatisfied (Funes et al., 2016; Rodríguez et al., 2019). • By 2060, more than 20% of exploited fishes and invertebrates currently found in eastern Mediterranean could become locally extinct (Jones and Cheung, 2015; Cheung et al., 2016; MedECC, 2020). High losses of clawed lobster production by the end of the century have also been projected under RCP4.5 (Boavida-Portugal et al., 2018). For much of the Mediterranean, fisheries' revenue may decrease by 15-30% by 2050 relative to 2000 under RCP8.5 (Lam et al., 2016).
High to Very High	2.3–4.0	<i>Medium</i>	<ul style="list-style-type: none"> • Decrease of wheat yield by 5-59% in Algeria (Chourghal et al., 2016; Dettori et al., 2017; Iocola et al., 2017; Brouziyne et al., 2018; Bouregaa, 2019; Cammarano et al., 2019; Kheir et al., 2019), by 13-17% in Egypt, by more than 14% in Italy (Chourghal et al., 2016; Dettori et al., 2017; Iocola et al., 2017; Brouziyne et al., 2018; Kheir et al., 2019). • Decrease of barley yield by 22-29% in Algeria (Bouregaa, 2019; Cammarano et al., 2019). • Decrease of olive yield by 15-64% in Algeria (Bouregaa, 2019; Cammarano et al., 2019). • Change of tomato yield by -94% to +12% in Tunisia, depending on soil type and sowing date, or even non-viable crops under some combinations of soil type-sowing date (Bird et al., 2016), • In northeastern Spain, 28-72% of the years after 2070 could have winters not fulfilling chilling requirements for apple trees (Funes et al., 2016; Rodríguez et al., 2019), while chilling requirements will be severely compromised for other fruit trees in Spain (Funes et al., 2016; Rodríguez et al., 2019). • Early maturation may result in unbalanced wines through higher sugar and lower acids in the grape must after 2050 under RCP8.5 (Fraga et al., 2016; Koufos et al., 2018). Reduction in table quality vines and wine grape production in southern Europe due to a future increase in the cumulative thermal stress and dryness during the growing season (Cardell et al., 2019). • Net irrigation requirements for date palms in Tunisia under RCP8.5 are expected to increase by 34% in 2050 compared to 2018 to sustain date production (Haj-Amor et al., 2020).

1 **Table SMCCP4.2f:** Supporting material for health and wellbeing.

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
Undetectable to Moderate	0.7-1.3	<i>Medium</i>	<ul style="list-style-type: none"> Population exposed to high or very high heat stress risk compared to 1986-2005 is expected to increase by 53 million people in northern Mediterranean and by 82 million people in southwestern Mediterranean (Gasparrini et al., 2017; Rohat et al., 2019). 1.3-1.6 times higher heat-related excess mortality in Italy and Spain compared to 1987-2010 (Gasparrini et al., 2017; Rohat et al., 2019). Increase of specific cooling demand by 50-143% in office buildings at northern Mediterranean cities (Cellura et al., 2018).
Moderate to High	1.5-2.0	<i>High</i>	<ul style="list-style-type: none"> Population exposed to high or very high heat stress risk compared to 1986-2005 expected to increase by 93 million people in northern Mediterranean (Gasparrini et al., 2017; Rohat et al., 2019) In northern Mediterranean, about 13-20,000 additional attributable deaths per warm season compared to the reference (i.e. apparent temperatures at the historical levels observed during the period 1971-2001) (Kendrovski et al., 2017). Heat-related excess mortality in Italy and Spain could increase by 2.9-3.7 times compared to 1987-2010 (Gasparrini et al., 2017; Rohat et al., 2019). Decadal mortality risk ratio compared to the historical mortality risk could equal 3-7 in southern Mediterranean (Ahmadalipour and Moradkhani, 2018). More than 20,000 climate-related deaths over the entire Mediterranean due to sub-optimal fruit and vegetable consumption (Springmann et al., 2016). The predicted probability of districts with West Nile Virus infections over the entire Mediterranean could range from 0.02 to more than 0.5, compared to an average of 0.0125 in 2014 (Semenza et al., 2016). Increase of the epidemic potential of dengue fever in southern Europe (Liu-Helmersson et al., 2019)
High to Very High	2.1-3.1	<i>Medium</i>	<p>Under warming levels of 2.1 – 3.1°C:</p> <ul style="list-style-type: none"> Increase of specific cooling demand by 50-278% in office buildings at northern Mediterranean cities (Cellura et al., 2018) Heat-related excess mortality in Italy and Spain could increase by 3.6-3.7 times compared to 1987-2010 (Gasparrini et al., 2017; Rohat et al., 2019). Slight increase of the epidemic potential of dengue fever in southern Europe (Liu-Helmersson et al., 2019) Supporting evidence from Chapter 13 (Figure 13.30 and SM13.10). <p>Under warming levels exceeding 3.7°C:</p> <ul style="list-style-type: none"> Decadal mortality risk ratio compared to the historical mortality risk could equal 8-20 in southern Mediterranean (Ahmadalipour and Moradkhani, 2018). In northern Mediterranean, about 50,000 additional attributable deaths per warm season compared to the reference (i.e. apparent temperatures at the historical levels observed during the period 1971-2001) (Kendrovski et al., 2017). Heat-related excess mortality in Italy and Spain could increase by 6.3-6.8 times compared to 1987-2010 (Gasparrini et al., 2017; Rohat et al., 2019). Increase of the epidemic potential of dengue fever in southern Europe (Liu-Helmersson et al., 2019) Increase of specific cooling demand by 134-375% in office buildings at northern Mediterranean cities (Cellura et al., 2018). Supporting evidence from Chapter 13 (Figure 13.30 and SM13.10).

2

3

1 **Table SMCCP4.2g:** Supporting material for coastal risks

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
Undetectable to Moderate	0.7-1.5	<i>Medium</i>	<ul style="list-style-type: none"> Chronic flooding in Venice (Box 13.1) Mediterranean extreme water levels during storms increase consistently with sea-level rise (Marcos et al., 2009). Flood and erosion hazards and risks not attributed yet, highly depend on coastal management strategies (WGI Chapter 13; Frihy et al., 2010). Pocket beaches in the Mediterranean might be early responders to sea-level rise (Brunel and Sabatier, 2009). Some permanent flooding is observed in subsiding areas such as around Thessaloniki (Greece) (Raucoules et al., 2008). Times of emergence of climate induced erosion after 2050 in Mediterranean beaches (Le Cozannet et al., 2016)
Moderate to High	1.5-2.0	<i>High</i>	<ul style="list-style-type: none"> Multicriteria coastal risk analysis at the scale of the Mediterranean shows potential for high risks all around the basin, with more marked potential for high risk in the South-Eastern Mediterranean (Satta et al., 2017). Chronic flooding taking place at high-tide significant concern in Venice (Box 13.1). The MOSE barrier will prevent such flooding, but the closure of the barrier is projected to reach 2 months per year with 50cm of sea-level rise, and adversely impacts ship traffic and lagoon water exchange (Box 13.1). Chronic flooding also projected to affect other low-lying areas in the Mediterranean such as the Ebro Delta (Sayol and Marcos, 2018), but a full picture of the problem is still missing at the scale of the Mediterranean. Extreme water levels will change in response to two contrasting impacts of climate change in the Mediterranean: reduced storminess and increased or accelerating mean sea-level rise due to climate change, with the latter dominating over the farther by the mid-21st century (WGI Chapters 9 and 12; Lionello et al., 2017), possibly becoming the main driver of extreme changes e.g. in the North Western Mediterranean by 2050 (Le Cozannet et al., 2015; Sayol and Marcos, 2018).
High to Very High	2.0-2.8	<i>High</i>	<ul style="list-style-type: none"> Hazards (e.g., extreme water levels amplification factors and allowances) and risks (e.g., economic average annual losses) projected to increase more quickly in the Mediterranean than in other regions of the world by 2050 (AR6 SROCC; Hallegatte et al., 2013), when GWL exceed 1.6 to 2.4°C (best estimate; AR6 WGI SPM). Shoreline retreat caused by permanent flooding projected to become widespread in the Mediterranean for RCP8.5 by the mid century, or for RCP4.5 during the second half of the century, and further aggravates for higher levels of climate forcing (Snoussi et al., 2008; Le Cozannet et al., 2016; Antonioli et al., 2017; Anzidei et al., 2017; Ciro Aucelli et al., 2017; Enríquez et al., 2017; Jiménez et al., 2017; Antonioli et al., 2020). Storm surge superimposed with a rise of 0.5 m in mean sea-level could result in the loss of up to 90% of pocket beaches in the Aegean archipelago, at least temporarily (Monioudi et al., 2017). Sediment inputs from major rivers limited by upstream dams, reducing potential compensation of erosion by sediments (Besset et al., 2017). Overtopping and other coastal risks induced by the combination of sea-level rise, storms and waves significant for Mediterranean ports and related activities above 50 cm of sea-level rise with respect to 1986-2005, especially in the South-Western Mediterranean (Sánchez-Arcilla et al., 2016; Sierra et al., 2016; Izaguirre et al., 2021). Wave agitations in ports increases due to sea-level rise despite projected decreasing offshore waves heights, reducing

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
			<p>operability times in ports, with shallower ports being projected to be more severely affected (Sierra et al., 2017)</p> <ul style="list-style-type: none"> • Groundwater salinization projected to increase with sea-level rise depending on the geological context and the processes through which aquifers are exploited and recharged. Sea-level rise not the only process causing salinization, but is projected to add another constraint to groundwater management and to sectors such as agriculture in a number of Mediterranean coastal floodplains and deltas, e.g. in some governorates of the Nile Delta for 1m of sea-level rise by 2100 (Mabrouk et al., 2018; Mastrocicco and Colombani, 2021; Pisinaras et al., 2021).

1
2
3
4

Table SMCCP4.2h: Supporting material for committed impacts of sea-level rise to cultural heritage, infrastructures and communities

Risk with no/low adaptation	Range of temperature transition	Confidence level for transition	Explanation (text & references)
Undetectable to Moderate	0.5-1.2	Low	<ul style="list-style-type: none"> • Mediterranean sea-levels controlled by sea-level changes in the Atlantic through Gibraltar Strait at multi-decadal timescales (Calafat et al., 2012), Mediterranean sea-level projections are close to global trends (Thiéblemont et al., 2019) (see SMCCP4.4). • Pre-industrial temperatures resulted in stable sea-level (AR5 WGI Chapter 13), whereas now recent greenhouse gas emissions have committed 0.7 to 1.1 m of sea-level rise by 2300 (AR6 WGI Chapter 9) (Nauels et al., 2019). • Compared to other regions, the Mediterranean Region is characterized by a very high number of cultural heritage sites exposed to sea-level rises, some of which have been preserved since more than 2,000 years (Marzeion and Levermann, 2014). • To date, 37 Mediterranean UNESCO World Heritage sites are at risk from flooding, and 42 at risk of erosion (Reimann et al., 2018). This includes Venice, where high-tide flooding has increased consistently with relative sea-level changes (Box 13.1).
Moderate to High	1.2-1.5	Medium	<ul style="list-style-type: none"> • Sea level projected to rise between 0.3 to 3.1 m by 2300 for SSP1-2.6 (<i>low confidence</i>) (WGI, Chapter 9). • Sea level committed to rise by 2 to 3 m after 2,000 years, and by 6 to 7 m after 10,000 years for 1.5°C of GWL, but these long term projections incorporate processes in which there is <i>low confidence</i> (WGI Chapter 9). • The number of UNESCO World Heritage sites exposed to flooding (erosion) increases from 37 to 40 (from 42 to 46) for a sea-level rise of 1.2 m (“high-end scenario” based on (Kopp et al., 2014) in (Reimann et al., 2018)). Flood frequencies, depths and erosion rates increase significantly in each of the cultural heritage sites already affected today (Reimann et al., 2018).
High to Very High	1.5-2.0	High	<ul style="list-style-type: none"> • Sea-levels projected to rise between 0.3 to 3.1 m by 2300 for SSP1-2.6 (<i>low confidence</i>) (WGI, Chapter 9). • Sea-levels committed to rise by 2 to 6 m after 2000 years, and by 8 to 13 m after 10,000 years for 2°C of GWL, but these long term projections incorporate processes in which there is <i>low confidence</i> (WGI, Chapter 9). • In at least 13 countries of the Mediterranean region, stabilization of global warming at about 2°C leads to drowning areas where at least 10% of the population currently live (Clark et al., 2016). The South-Eastern Mediterranean low-lying areas projected to be drowned are particularly densely populated (Clark et al., 2016).

5

SMCCP4.3 Sustainable Development in the Mediterranean Basin

Table SMCCP4.3: Indicators for the achievement of the Sustainable Development Goals (SDGs) as illustrated in Figure CCP4.9

SDG1 No poverty: (sdg1_320pov; Poverty headcount ratio at \$3.20/day (% population))
SDG2 Zero hunger: (sdg2_crlyld; Cereal yield (t/ha))
SDG3 Good health and well-being: (sdg3_uhc; Universal Health Coverage Tracer Index (0-100))
SDG4 Quality education: (sdg4_second; Lower secondary education completion rate (%))
SDG5 Gender equality: (sdg5_lfpr; Ratio of female to male labour force participation rate)
SDG6 Clean water and sanitation: (sdg6_water; Population using at least basic drinking water services (%))
SDG7 Affordable and clean energy: (sdg7_co2twh; CO2 emissions from fuel combustion / electricity output (MtCO2/TWh))
SDG8 Decent work and economic growth: (sdg9_intuse; Population using the internet (%))
SDG9 Industry, innovation and infrastructure: (sdg9_rdex; Research and development expenditure (% GDP))
SDG10 Reduced inequalities: (sdg10_adjgini; Gini Coefficient adjusted for top income (1-100))
SDG11 Sustainable cities and communities: (sdg11_pm25; Annual mean concentration of particulate matter of less than 2.5 microns of diameter (PM2.5) ($\mu\text{g}/\text{m}^3$)),
SDG12 Responsible consumption and production:
SDG13 Climate action: (sdg13_co2pc; Energy-related CO2 emissions per capita (tCO2/capita))
SDG14 Life below water: (sdg14_cpma; Mean area that is protected in marine sites important to biodiversity (%))
SDG15 Life on land: (sdg15_redlist; Red List Index of species survival (0-1))
SDG16 Peace, justice and strong institutions: (sdg16_homicides; Homicides (per 100,000 population))
SDG17 Partnerships for the goals: (sdg17_govex; Government Health and Education spending (% GDP))

The list of Northern Mediterranean countries (N) includes Bosnia and Herzegovina, Croatia, Cyprus, France, Greece, Italy, Malta, Monaco, Portugal and Spain. Southern Mediterranean countries (S) analysed are Algeria, Egypt, Jordan, Lebanon, Morocco, Syrian Arab Republic, Tunisia and Turkey.

Data extracted from (Sachs et al., 2019). All analyses reported significantly different mean SDG indicator values between N and S regions with $p < 0.0001$ (Tukey-Kramer test).

SMCCP4.4 Mediterranean Sea-level Projections

Mediterranean sea-level projections presented in the FAQ are downscaled from those presented in AR6 WGI Chapter 9 and Technical Summary, following the method presented in Thiéblemont et al. (2019) (Table SMCCP4.4). Until 2100, the Mediterranean stereodynamic sea-levels are assumed to follow stereodynamic sea-level changes in the Atlantic near Gibraltar. The procedure ignores intra-basin stereodynamic sea-level changes, but it avoids biases due to the resolution of models and the coarse representation of water exchanges through the Gibraltar straight (Landerer et al., 2014; Meyssignac et al., 2017; Parras-Berrocal et al., 2020). Furthermore, observational evidence indicates that multi-decadal sea-level changes in the Mediterranean basin follow the Gibraltar forcing (Section 4.1; Calafat et al., 2012). Beyond 2150, Mediterranean stereodynamic sea-levels are assumed to follow the mean global thermal expansion.

Table SMCCP4.4: Mean sea-level projections in the Mediterranean relative to 1995-2014. There is *medium confidence* in the median values [likely range] shown for the five SSP scenarios. There is *low confidence* in the median values [17-83rd percentiles] shown for SSP5-8.5-L, a scenario based on a Structured-Expert Judgement or assuming marine ice-cliffs instabilities (See AR6 WGI Chapter 9 for details). Mediterranean projections include the mean Glacial Isostatic Adjustment in the Mediterranean, but not local vertical ground motions due to natural or anthropogenic processes such as tectonics or groundwater extractions.

	2050 (m)	2100 (m)	2150 (m)
SSP1-1.9	0.22 [0.15-0.30]	0.43 [0.31-0.61]	0.52 [0.32 - 0.81]
SSP1-2.6	0.22 [0.16-0.30]	0.46 [0.34-0.66]	0.62 [0.39 - 0.95]
SSP2-4.5	0.23 [0.17-0.31]	0.57 [0.44-0.79]	0.83 [0.57 - 1.23]
SSP3-7.0	0.24 [0.18-0.31]	0.69 [0.56-0.94]	1.08 [0.80 - 1.57]
SSP5-8.5	0.26 [0.20-0.33]	0.79 [0.64-1.06]	1.22 [0.91 - 1.78]
SSP5-8.5-L	0.24 [0.18-0.33]	0.87 [0.65-1.38]	1.89 [0.96-5.01]

References

- Aguilera, R., R. Marce and S. Sabater, 2015: Detection and attribution of global change effects on river nutrient dynamics in a large Mediterranean basin. *Biogeosciences*, **12**(13), 4085-4098, doi:10.5194/bg-12-4085-2015.
- Ahmadalipour, A. and H. Moradkhani, 2018: Escalating heat-stress mortality risk due to global warming in the Middle East and North Africa (MENA). *Environment International*, **117**, 215-225, doi:10.1016/j.envint.2018.05.014.
- Albano, P. G. et al., 2021: Native biodiversity collapse in the eastern Mediterranean. *Proceedings of the Royal Society B: Biological Sciences*, **288**(1942), 20202469, doi:10.1098/rspb.2020.2469.
- Albouy, C. et al., 2013: Projected climate change and the changing biogeography of coastal Mediterranean fishes. *Journal of Biogeography*, **40**(3), 534-547, doi:10.1111/jbi.12013.
- Andrew, J. T. and E. Sauquet, 2017: Climate change impacts and water management adaptation in two Mediterranean-climate watersheds: Learning from the Durance and Sacramento rivers. *Water*, **9**(2), 126, doi:10.3390/w9020126.
- Antonioli, F. et al., 2017: Sea-level rise and potential drowning of the Italian coastal plains: Flooding risk scenarios for 2100. *Quaternary Science Reviews*, **158**, 29-43, doi:10.1016/j.quascirev.2016.12.021.
- Antonioli, F. et al., 2020: Relative sea-level rise and potential submersion risk for 2100 on 16 coastal plains of the Mediterranean Sea. *Water*, **12**(8), 2173.
- Anzidei, M. et al., 2017: Flooding scenarios due to land subsidence and sea-level rise: a case study for Lipari Island (Italy). *Terra Nova*, **29**(1), 44-51, doi:10.1111/ter.12246.
- Argaman, E., R. Barth, Y. Moshe and M. Ben-Hur, 2020: Long-term effects of climatic and hydrological variation on natural vegetation production and characteristics in a semiarid watershed: The northern Negev, Israel. *Science of The Total Environment*, **747**, 141146, doi:10.1016/j.scitotenv.2020.141146.
- Azzurro, E. et al., 2019: Climate change, biological invasions, and the shifting distribution of Mediterranean fishes: A large-scale survey based on local ecological knowledge. *Global Change Biology*, **25**(8), 2779-2792, doi:10.1111/gcb.14670.
- Baahmed, D., L. Oudin and M. Errih, 2015: Current runoff variations in the Macta catchment (Algeria): is climate the sole factor? *Hydrological Sciences Journal*, **60**(7-8), 1331-1339, doi:10.1080/02626667.2014.975708.
- Barredo, J. I., G. Caudullo and A. Dosio, 2016: Mediterranean habitat loss under future climate conditions: Assessing impacts on the Natura 2000 protected area network. *Applied Geography*, **75**, 83-92, doi:10.1016/j.apgeog.2016.08.003.
- Bartsch, S. et al., 2020: Impact of precipitation, air temperature and abiotic emissions on gross primary production in Mediterranean ecosystems in Europe. *European Journal of Forest Research*, **139**(1), 111-126, doi:10.1007/s10342-019-01246-7.
- Ben Rais Lasram, F. et al., 2010: The Mediterranean Sea as a 'cul-de-sac' for endemic fishes facing climate change. *Global Change Biology*, **16**(12), 3233-3245, doi:10.1111/j.1365-2486.2010.02224.x.
- Benedetti, F., F. Guilhaumon, F. Adloff and S.-D. Ayata, 2018: Investigating uncertainties in zooplankton composition shifts under climate change scenarios in the Mediterranean Sea. *Ecography*, **41**(2), 345-360, doi:10.1111/ecog.02434.
- Besset, M., E. J. Anthony and F. Sabatier, 2017: River delta shoreline reworking and erosion in the Mediterranean and Black Seas: the potential roles of fluvial sediment starvation and other factors. *Elementa: Science of the Anthropocene*, **5**, doi:10.1525/elementa.139.
- Bird, D. N. et al., 2016: Modelling climate change impacts on and adaptation strategies for agriculture in Sardinia and Tunisia using AquaCrop and value-at-risk. *Science of The Total Environment*, **543**, 1019-1027, doi:10.1016/j.scitotenv.2015.07.035.
- Blöschl, G. et al., 2017: Changing climate shifts timing of European floods. *Science*, **357**(6351), 588, doi:10.1126/science.aan2506.
- Boavida-Portugal, J. et al., 2018: Climate change impacts on the distribution of coastal lobsters. *Marine Biology*, **165**(12), 186, doi:10.1007/s00227-018-3441-9.
- Bouregaa, T., 2019: Impact of climate change on yield and water requirement of rainfed crops in the Setif region. *Management of Environmental Quality: An International Journal*, **30**(4), 851-863, doi:10.1108/MEQ-06-2018-0110.
- Braca, G. et al., 2019: Evaluation of national and regional groundwater resources under climate change scenarios using a GIS-based water budget procedure. *Rendiconti Lincei. Scienze Fisiche e Naturali*, **30**(1), 109-123, doi:10.1007/s12210-018-00757-6.
- Bregaglio, S. et al., 2017: Identifying trends and associated uncertainties in potential rice production under climate change in Mediterranean areas. *Agricultural and Forest Meteorology*, **237-238**, 219-232, doi:10.1016/j.agrformet.2017.02.015.
- Brilli, L. et al., 2019: Carbon sequestration capacity and productivity responses of Mediterranean olive groves under future climates and management options. *Mitigation and Adaptation Strategies for Global Change*, **24**(3), 467-491, doi:10.1007/s11027-018-9824-x.
- Brotans, L. et al., 2013: How fire history, fire suppression practices and climate change affect wildfire regimes in Mediterranean landscapes. *PLoS ONE*, **8**(5), e62392, doi:10.1371/journal.pone.0062392.

- 1 Brouziyne, Y. et al., 2018: Modeling sustainable adaptation strategies toward a climate-smart agriculture in a
2 Mediterranean watershed under projected climate change scenarios. *Agricultural Systems*, **162**, 154-163,
3 doi:10.1016/j.agsy.2018.01.024.
- 4 Brunel, C. and F. Sabatier, 2009: Potential influence of sea-level rise in controlling shoreline position on the French
5 Mediterranean Coast. *Geomorphology*, **107**(1), 47-57, doi:10.1016/j.geomorph.2007.05.024.
- 6 Bucak, T. et al., 2017: Future water availability in the largest freshwater Mediterranean lake is at great risk as evidenced
7 from simulations with the SWAT model. *Science of The Total Environment*, **581-582**, 413-425,
8 doi:10.1016/j.scitotenv.2016.12.149.
- 9 Burrows, M. T. et al., 2014: Geographical limits to species-range shifts are suggested by climate velocity. *Nature*,
10 **507**(7493), 492-495, doi:10.1038/nature12976.
- 11 Calafat, F. M., D. P. Chambers and M. N. Tsimplis, 2012: Mechanisms of decadal sea level variability in the eastern
12 North Atlantic and the Mediterranean Sea. *Journal of Geophysical Research: Oceans*, **117**(C9),
13 doi:10.1029/2012JC008285.
- 14 Caloiero, T., S. Veltri, P. Caloiero and F. Frustaci, 2018: Drought analysis in Europe and in the Mediterranean Basin
15 using the Standardized Precipitation Index. *Water*, **10**(8), doi:10.3390/w10081043.
- 16 Cammarano, D. et al., 2019: The impact of climate change on barley yield in the Mediterranean basin. *European
17 Journal of Agronomy*, **106**, 1-11, doi:10.1016/j.eja.2019.03.002.
- 18 Cardell, M. F., A. Amengual and R. Romero, 2019: Future effects of climate change on the suitability of wine grape
19 production across Europe. *Regional Environmental Change*, **19**(8), 2299-2310, doi:10.1007/s10143-019-01502-x.
- 20 Cellura, M., F. Guarino, S. Longo and G. Tumminia, 2018: Climate change and the building sector: Modelling and
21 energy implications to an office building in southern Europe. *Energy for Sustainable Development*, **45**, 46-65,
22 doi:10.1016/j.esd.2018.05.001.
- 23 Cheung, W. W. L. et al., 2016: Structural uncertainty in projecting global fisheries catches under climate change.
24 *Ecological Modelling*, **325**, 57-66, doi:10.1016/j.ecolmodel.2015.12.018.
- 25 Chourghal, N., J. P. Lhomme, F. Huard and A. Aidaoui, 2016: Climate change in Algeria and its impact on durum
26 wheat. *Regional Environmental Change*, **16**(6), 1623-1634, doi:10.1007/s10113-015-0889-8.
- 27 Ciro Aucelli, P. P. et al., 2017: Coastal inundation risk assessment due to subsidence and sea level rise in a
28 Mediterranean alluvial plain (Voturno coastal plain – southern Italy). *Estuarine, Coastal and Shelf Science*, **198**,
29 597-609, doi:10.1016/j.ecss.2016.06.017.
- 30 Clark, P. U. et al., 2016: Consequences of twenty-first-century policy for multi-millennial climate and sea-level change.
31 *Nature Climate Change*, **6**(4), 360-369, doi:10.1038/nclimate2923.
- 32 Coll, M. et al., 2014: Assessing fishing and marine biodiversity changes using fishers' perceptions: The Spanish
33 Mediterranean and Gulf of Cadiz case study. *PLoS ONE*, **9**(1), e85670, doi:10.1371/journal.pone.0085670.
- 34 Corrales, X. et al., 2018: Future scenarios of marine resources and ecosystem conditions in the Eastern Mediterranean
35 under the impacts of fishing, alien species and sea warming. *Scientific Reports*, **8**(1), 14284, doi:10.1038/s41598-
36 018-32666-x.
- 37 Corrales, X. et al., 2017: Hindcasting the dynamics of an Eastern Mediterranean marine ecosystem under the impacts of
38 multiple stressors. *Marine Ecology Progress Series*, **580**, 17-36, doi:10.3354/meps12271.
- 39 Curt, T., A. Aini and S. Dupire, 2020: Fire activity in Mediterranean forests (the Algerian case). *Fire*, **3**(4), 58.
- 40 Dakhlaoui, H., J. Seibert and K. Hakala, 2020: Sensitivity of discharge projections to potential evapotranspiration
41 estimation in Northern Tunisia. *Regional Environmental Change*, **20**(2), 34, doi:10.1007/s10113-020-01615-8.
- 42 Dettori, M., C. Cesaraccio and P. Duce, 2017: Simulation of climate change impacts on production and phenology of
43 durum wheat in Mediterranean environments using CERES-Wheat model. *Field Crops Research*, **206**, 43-53,
44 doi:10.1016/j.fcr.2017.02.013.
- 45 Di Lena, B., O. Silvestroni, V. Lanari and A. Palliotti, 2018: Climate change effects on cv. Montepulciano in some
46 wine-growing areas of the Abruzzi region (Italy). *Theoretical and Applied Climatology*, doi:10.1007/s00704-018-
47 2545-y.
- 48 EEA, 2017: *Climate change, impacts and vulnerability in Europe 2016: An indicator-based report*. European
49 Environment Agency. Publications Office of the European Union, Luxembourg, 424 pp. Available at:
50 <https://www.eea.europa.eu/publications/climate-change-impacts-and-vulnerability-2016> (accessed 15/10/2020).
- 51 El-Maayar, M. and M. A. Lange, 2013: A methodology to infer crop yield response to climate variability and change
52 using long-term observations. *Atmosphere*, **4**, 365-382, doi:10.3390/atmos4040365.
- 53 Enríquez, A. R. et al., 2017: Changes in beach shoreline due to sea level rise and waves under climate change scenarios:
54 application to the Balearic Islands (western Mediterranean). *Natural Hazards and Earth System Sciences*, **17**(7),
55 1075-1089, doi:10.5194/nhess-17-1075-2017.
- 56 Faggian, P., 2018: Estimating fire danger over Italy in the next decades. *Euro-Mediterranean Journal for
57 Environmental Integration*, **3**(1), 15, doi:10.1007/s41207-018-0053-1.
- 58 Fernandes, P. M., A. M. G. Barros, A. Pinto and J. A. Santos, 2016: Characteristics and controls of extremely large
59 wildfires in the western Mediterranean Basin. *Journal of Geophysical Research: Biogeosciences*, **121**(8), 2141-
60 2157, doi:10.1002/2016JG003389.
- 61 Fortibuoni, T. et al., 2015: Climate impact on Italian fisheries (Mediterranean Sea). *Regional Environmental Change*,
62 **15**(5), 931-937, doi:10.1007/s10113-015-0781-6.

- 1 Fraga, H., I. García de Cortázar Aauri, A. C. Malheiro and J. A. Santos, 2016: Modelling climate change impacts on
2 viticultural yield, phenology and stress conditions in Europe. *Global Change Biology*, **22**(11), 3774-3788,
3 doi:10.1111/gcb.13382.
- 4 Fraga, H., J. G. Pinto, F. Viola and J. A. Santos, 2020: Climate change projections for olive yields in the Mediterranean
5 Basin. *International Journal of Climatology*, **40**(2), 769-781, doi:10.1002/joc.6237.
- 6 Frihy, O. E. S., E. A. Deabes, S. M. Shereet and F. A. Abdalla, 2010: Alexandria-Nile Delta coast, Egypt: update and
7 future projection of relative sea-level rise. *Environmental Earth Sciences*, **61**(2), 253-273, doi:10.1007/s12665-
8 009-0340-x.
- 9 Funes, I. et al., 2016: Future climate change impacts on apple flowering date in a Mediterranean subbasin. *Agricultural*
10 *Water Management*, **164**, 19-27, doi:10.1016/j.agwat.2015.06.013.
- 11 Garbolino, E., V. Sanseverino-Godfrin and G. Hinojos-Mendoza, 2016: Describing and predicting of the vegetation
12 development of Corsica due to expected climate change and its impact on forest fire risk evolution. *Safety Science*,
13 **88**, 180-186, doi:10.1016/j.ssci.2016.02.006.
- 14 Garcia-Mozo, H., J. Oteros and C. Galan, 2015: Phenological changes in olive (*Olea europaea* L.) reproductive cycle in
15 southern Spain due to climate change. *Annals of Agricultural and Environmental Medicine*, **22**(3), 421-428,
16 doi:10.5604/12321966.1167706.
- 17 Garrabou, J. et al., 2021: Sliding toward the collapse of Mediterranean coastal marine rocky ecosystems. In: *Ecosystem*
18 *Collapse and Climate Change* [Canadell, J. G. and R. B. Jackson (eds.)]. Springer International Publishing, Cham,
19 pp. 291-324. ISBN 978-3-030-71330-0.
- 20 Gasparrini, A. et al., 2017: Projections of temperature-related excess mortality under climate change scenarios. *The*
21 *Lancet Planetary Health*, **1**(9), e360-e367, doi:10.1016/S2542-5196(17)30156-0.
- 22 Gaume, E. et al., 2016: Mediterranean extreme floods and flash floods. In: *The Mediterranean Region under Climate*
23 *Change: A Scientific Update* [Thiébaud, S., J.-P. Moatti, V. Ducrocq, E. Gaume, F. Dulac, E. Hamonou, Y.-J.
24 Shin, J. Guiot, W. Cramer, G. Boulet, J.-F. Guégan, R. Barouki, I. Annesi-Maesano, P. Marty, E. Torquebiau, J. F.
25 Soussana, Y. Aumeeruddy-Thomas, J.-L. Chotte and D. Lacroix (eds.)]. IRD Éditions, Marseille, France, pp. 133-
26 144. ISBN 9782709922203.
- 27 Georgopoulou, E. et al., 2017: Climate change impacts and adaptation options for the Greek agriculture in 2021–2050:
28 A monetary assessment. *Climate Risk Management*, **16**, 164-182, doi:10.1016/j.crm.2017.02.002.
- 29 Givan, O., D. Edelist, O. Sonin and J. Belmaker, 2018: Thermal affinity as the dominant factor changing Mediterranean
30 fish abundances. *Global Change Biology*, **24**(1), E80-E89, doi:10.1111/gcb.13835.
- 31 Gosling, S. N. et al., 2017: A comparison of changes in river runoff from multiple global and catchment-scale
32 hydrological models under global warming scenarios of 1°C, 2°C and 3°C. *Climatic Change*, **141**(3), 577-595,
33 doi:10.1007/s10584-016-1773-3.
- 34 Grillakis, M. G., 2019: Increase in severe and extreme soil moisture droughts for Europe under climate change. *Science*
35 *of The Total Environment*, **660**, 1245-1255, doi:10.1016/j.scitotenv.2019.01.001.
- 36 Gudmundsson, L., S. I. Seneviratne and X. Zhang, 2017: Anthropogenic climate change detected in European
37 renewable freshwater resources. *Nature Climate Change*, **7**, 813, doi:10.1038/nclimate3416.
- 38 Guiot, J. and W. Cramer, 2016: Climate change: The 2015 Paris Agreement thresholds and Mediterranean basin
39 ecosystems. *Science*, **354**(6311), 465-468, doi:10.1126/science.aah5015.
- 40 Guyennon, N., F. Salerno, I. Portoghese and E. Romano, 2017: Climate change adaptation in a Mediterranean semi-arid
41 catchment: Testing managed aquifer recharge and increased surface reservoir capacity. *Water*, **9**(9), 689,
42 doi:10.3390/w9090689.
- 43 Haj-Amor, Z., T. Kumar Acharjee, L. Dhaouadi and S. Bouri, 2020: Impacts of climate change on irrigation water
44 requirement of date palms under future salinity trend in coastal aquifer of Tunisian oasis. *Agricultural Water*
45 *Management*, **228**, 105843, doi:10.1016/j.agwat.2019.105843.
- 46 Hallegatte, S., C. Green, R. J. Nicholls and J. Corfee-Morlot, 2013: Future flood losses in major coastal cities. *Nature*
47 *Climate Change*, **3**(9), 802-806, doi:10.1038/nclimate1979.
- 48 Hoerling, M. et al., 2012: On the increased frequency of Mediterranean drought. *Journal of Climate*, **25**(6), 2146-2161,
49 doi:10.1175/JCLI-D-11-00296.1.
- 50 Iocola, I. et al., 2017: Can conservation tillage mitigate climate change impacts in Mediterranean cereal systems? A soil
51 organic carbon assessment using long term experiments. *European Journal of Agronomy*, **90**, 96-107,
52 doi:10.1016/j.eja.2017.07.011.
- 53 Izaguirre, C. et al., 2021: Climate change risk to global port operations. *Nature Climate Change*, **11**(1), 14-20,
54 doi:10.1038/s41558-020-00937-z.
- 55 Jarić, I. et al., 2019: Susceptibility of European freshwater fish to climate change: Species profiling based on life-
56 history and environmental characteristics. *Global Change Biology*, **25**(2), 448-458,
57 doi:<https://doi.org/10.1111/gcb.14518>.
- 58 Jiménez, J. A. et al., 2017: Impacts of sea-level rise-induced erosion on the Catalan coast. *Regional Environmental*
59 *Change*, **17**(2), 593-603, doi:10.1007/s10113-016-1052-x.
- 60 Jones, M. C. and W. W. L. Cheung, 2015: Multi-model ensemble projections of climate change effects on global
61 marine biodiversity. *ICES Journal of Marine Science*, **72**(3), 741-752, doi:10.1093/icesjms/fsu172.
- 62 Karali, A. et al., 2014: Sensitivity and evaluation of current fire risk and future projections due to climate change: the
63 case study of Greece. *Nat. Hazards Earth Syst. Sci.*, **14**(1), 143-153, doi:10.5194/nhess-14-143-2014.

- 1 Kendrovski, V. et al., 2017: Quantifying projected heat mortality impacts under 21st-century warming conditions for
2 selected European countries. *International Journal of Environmental Research and Public Health*, **14**(7), 729,
3 doi:10.3390/ijerph14070729.
- 4 Kheir, A. M. S. et al., 2019: Impacts of rising temperature, carbon dioxide concentration and sea level on wheat
5 production in North Nile delta. *Science of The Total Environment*, **651**, 3161-3173,
6 doi:10.1016/j.scitotenv.2018.10.209.
- 7 Kim, G.-U., K.-H. Seo and D. Chen, 2019: Climate change over the Mediterranean and current destruction of marine
8 ecosystem. *Scientific Reports*, **9**(1), 18813, doi:10.1038/s41598-019-55303-7.
- 9 Kopp, R. E. et al., 2014: Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge
10 sites. *Earth's Future*, **2**(8), 383-406, doi:<https://doi.org/10.1002/2014EF000239>.
- 11 Koufos, G. C., T. Mavromatis, S. Koundouras and G. V. Jones, 2018: Response of viticulture-related climatic indices
12 and zoning to historical and future climate conditions in Greece. *International Journal of Climatology*, **38**(4),
13 2097-2111, doi:10.1002/joc.5320.
- 14 Koutroulis, A. G. et al., 2016: Cross sectoral impacts on water availability at +2°C and +3°C for east Mediterranean
15 island states: The case of Crete. *Journal of Hydrology*, **532**, 16-28, doi:10.1016/j.jhydrol.2015.11.015.
- 16 Kuglitsch, F. G. et al., 2010: Heat wave changes in the eastern Mediterranean since 1960. *Geophysical Research
17 Letters*, **37**(4), doi:10.1029/2009gl041841.
- 18 Kundzewicz, Z. W., I. Pińskwar and G. R. Brakenridge, 2017: Changes in river flood hazard in Europe: a review.
19 *Hydrology Research*, **49**(2), 294-302, doi:10.2166/nh.2017.016.
- 20 Lam, V. W. Y., W. W. L. Cheung, G. Reygondeau and U. R. Sumaila, 2016: Projected change in global fisheries
21 revenues under climate change. *Scientific Reports*, **6**(1), 32607, doi:10.1038/srep32607.
- 22 Landerer, F. W., P. J. Gleckler and T. Lee, 2014: Evaluation of CMIP5 dynamic sea surface height multi-model
23 simulations against satellite observations. *Climate Dynamics*, **43**(5), 1271-1283, doi:10.1007/s00382-013-1939-x.
- 24 Le Cozannet, G. et al., 2016: Uncertainties in sandy shorelines evolution under the Bruun rule assumption. *Frontiers in
25 Marine Science*, **3**(49), doi:10.3389/fmars.2016.00049.
- 26 Le Cozannet, G. et al., 2015: Evaluating uncertainties of future marine flooding occurrence as sea-level rises.
27 *Environmental Modelling & Software*, **73**, 44-56, doi:10.1016/j.envsoft.2015.07.021.
- 28 Lefebvre, G. et al., 2019: Predicting the vulnerability of seasonally-flooded wetlands to climate change across the
29 Mediterranean Basin. *Science of The Total Environment*, **692**, 546-555, doi:10.1016/j.scitotenv.2019.07.263.
- 30 Lehner, F. et al., 2017: Projected drought risk in 1.5°C and 2°C warmer climates. *Geophysical Research Letters*, **44**(14),
31 7419-7428, doi:10.1002/2017gl074117.
- 32 Lejeune, C. et al., 2010: Climate change effects on a miniature ocean: the highly diverse, highly impacted
33 Mediterranean Sea. *Trends in Ecology & Evolution*, **25**(4), 250-260, doi:10.1016/j.tree.2009.10.009.
- 34 Lionello, P., D. Conte, L. Marzo and L. Scarascia, 2017: The contrasting effect of increasing mean sea level and
35 decreasing storminess on the maximum water level during storms along the coast of the Mediterranean Sea in the
36 mid 21st century. *Global and Planetary Change*, **151**, 80-91, doi:10.1016/j.gloplacha.2016.06.012.
- 37 Liu-Helmersson, J., J. Rocklöv, M. Sewe and Å. Brännström, 2019: Climate change may enable *Aedes aegypti*
38 infestation in major European cities by 2100. *Environmental Research*, **172**, 693-699,
39 doi:10.1016/j.envres.2019.02.026.
- 40 Llasat, M. C. et al., 2013: Towards a database on societal impact of Mediterranean floods within the framework of the
41 HYMEX project. *Natural Hazards and Earth System Sciences*, **13**(5), 1337-1350, doi:10.5194/nhess-13-1337-
42 2013.
- 43 Llasat, M. C. et al., 2016: Trends in flash flood events versus convective precipitation in the Mediterranean region: The
44 case of Catalonia. *Journal of Hydrology*, **541**, 24-37, doi:10.1016/j.jhydrol.2016.05.040.
- 45 Lloret, J. et al., 2015: How a multidisciplinary approach involving ethnoecology, biology and fisheries can help explain
46 the spatio-temporal changes in marine fish abundance resulting from climate change. *Global Ecology and
47 Biogeography*, **24**(4), 448-461, doi:10.1111/geb.12276.
- 48 Lobanova, A. et al., 2016: Impacts of changing climate on the hydrology and hydropower production of the Tagus
49 River basin. *Hydrological Processes*, **30**(26), 5039-5052, doi:10.1002/hyp.10966.
- 50 Lozano, O. M. et al., 2017: Assessing climate change impacts on wildfire exposure in Mediterranean areas. *Risk
51 Analysis*, **37**(10), 1898-1916, doi:10.1111/risa.12739.
- 52 Mabrouk, M., A. Jonoski, G. H. P. Oude Essink and S. Uhlenbrook, 2018: Impacts of sea level rise and groundwater
53 extraction scenarios on fresh groundwater resources in the Nile Delta Governorates, Egypt. *Water*, **10**(11), 1690.
- 54 Macias, D. M., E. Garcia-Gorriiz and A. Stips, 2015: Productivity changes in the Mediterranean Sea for the twenty-first
55 century in response to changes in the regional atmospheric forcing. *Frontiers in Marine Science*, **2**(79),
56 doi:10.3389/fmars.2015.00079.
- 57 Maggiotto, G. et al., 2021: Heat waves and adaptation strategies in a mediterranean urban context. *Environmental
58 Research*, **197**, 111066, doi:10.1016/j.envres.2021.111066.
- 59 Manes, S. et al., 2021: Endemism increases species' climate change risk in areas of global biodiversity importance.
60 *Biological Conservation*, **257**, 109070, doi:<https://doi.org/10.1016/j.biocon.2021.109070>.
- 61 Marchane, A. et al., 2017: Climate change impacts on surface water resources in the Rheraya catchment (High Atlas,
62 Morocco). *Hydrological Sciences Journal*, **62**(6), 979-995, doi:10.1080/02626667.2017.1283042.

- 1 Marcos-Garcia, P., A. Lopez-Nicolas and M. Pulido-Velazquez, 2017: Combined use of relative drought indices to
2 analyze climate change impact on meteorological and hydrological droughts in a Mediterranean basin. *Journal of*
3 *Hydrology*, **554**, 292-305, doi:10.1016/j.jhydrol.2017.09.028.
- 4 Marcos, M., M. N. Tsimplis and A. G. P. Shaw, 2009: Sea level extremes in southern Europe. *Journal of Geophysical*
5 *Research: Oceans*, **114**(C1), doi:10.1029/2008JC004912.
- 6 Marx, A. et al., 2018: Climate change alters low flows in Europe under global warming of 1.5, 2, and 3 °C. *Hydrology*
7 *and Earth System Sciences*, **22**(2), 1017-1032, doi:10.5194/hess-22-1017-2018.
- 8 Marzeion, B. and A. Levermann, 2014: Loss of cultural world heritage and currently inhabited places to sea-level rise.
9 *Environmental Research Letters*, **9**(3), 034001, doi:10.1088/1748-9326/9/3/034001.
- 10 Masia, S. et al., 2018: Assessment of irrigated agriculture vulnerability under climate change in Southern Italy. *Water*,
11 **10**(2), 209, doi:10.3390/w10020209.
- 12 Mastrocicco, M. and N. Colombani, 2021: The issue of groundwater salinization in coastal areas of the Mediterranean
13 region: A review. *Water*, **13**(1), 90.
- 14 Mathbout, S. et al., 2018: Observed changes in daily precipitation extremes at annual timescale over the Eastern
15 Mediterranean during 1961-2012. *Pure and Applied Geophysics*, **175**(11), 3875-3890, doi:10.1007/s00024-017-
16 1695-7.
- 17 Matías, L., M. Abdelaziz, O. Godoy and L. Gómez-Aparicio, 2019: Disentangling the climatic and biotic factors driving
18 changes in the dynamics of *Quercus suber* populations across the species' latitudinal range. *Diversity and*
19 *Distributions*, **25**(4), 524-535, doi:10.1111/ddi.12873.
- 20 Meddour-Sahar, O., 2015: Wildfires in Algeria: problems and challenges. *iForest - Biogeosciences and Forestry*, **8**(6),
21 818-826, doi:10.3832/ifer1279-007.
- 22 MedECC, 2020: Summary for Policymakers. In: *Climate and Environmental Change in the Mediterranean Basin –*
23 *Current Situation and Risks for the Future. First Mediterranean Assessment Report* [Cramer, W., J. Guiot and K.
24 Marini (eds.)]. Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 11-40.
- 25 Mediero, L., D. Santillán, L. Garrote and A. Granados, 2014: Detection and attribution of trends in magnitude,
26 frequency and timing of floods in Spain. *Journal of Hydrology*, **517**, 1072-1088,
27 doi:10.1016/j.jhydrol.2014.06.040.
- 28 Meyssignac, B. et al., 2017: Evaluating model simulations of Twentieth-Century sea-level rise. Part II: Regional sea-
29 level changes. *Journal of Climate*, **30**(21), 8565-8593, doi:10.1175/JCLI-D-17-0112.1.
- 30 Monioudi, I. N. et al., 2017: Assessment of island beach erosion due to sea level rise: the case of the Aegean
31 archipelago (Eastern Mediterranean). *Nat. Hazards Earth Syst. Sci.*, **17**(3), 449-466, doi:10.5194/nhess-17-449-
32 2017.
- 33 Moore, F. C. and D. B. Lobell, 2015: The fingerprint of climate trends on European crop yields. *Proceedings of the*
34 *National Academy of Sciences of the United States of America*, **112**(9), 2670-2675, doi:10.1073/pnas.1409606112.
- 35 Moullec, F. et al., 2019: An end-to-end model reveals losers and winners in a warming Mediterranean Sea. *Frontiers in*
36 *Marine Science*, **6**(345), doi:10.3389/fmars.2019.00345.
- 37 Mourato, S., M. Moreira and J. Corte-Real, 2015: Water resources impact assessment under climate change scenarios in
38 Mediterranean watersheds. *Water Resources Management*, **29**(7), 2377-2391, doi:10.1007/s11269-015-0947-5.
- 39 Nauels, A. et al., 2019: Attributing long-term sea-level rise to Paris Agreement emission pledges. *Proceedings of the*
40 *National Academy of Sciences*, **116**(47), 23487-23492, doi:10.1073/pnas.1907461116.
- 41 Okkan, U. and U. Kirdemir, 2018: Investigation of the behavior of an agricultural-operated dam reservoir under RCP
42 scenarios of AR5-IPCC. *Water Resources Management*, **32**(8), 2847-2866, doi:10.1007/s11269-018-1962-0.
- 43 Oteros, J. et al., 2015: Variations in cereal crop phenology in Spain over the last twenty-six years (1986-2012). *Climatic*
44 *Change*, **130**(4), 545-558, doi:10.1007/s10584-015-1363-9.
- 45 Paprotny, D., A. Sebastian, O. Morales-Nápoles and S. N. Jonkman, 2018: Trends in flood losses in Europe over the
46 past 150 years. *Nature Communications*, **9**(1), 1985, doi:10.1038/s41467-018-04253-1.
- 47 Parras-Berrocal, I. M. et al., 2020: The climate change signal in the Mediterranean Sea in a regionally coupled
48 atmosphere–ocean model. *Ocean Sci.*, **16**(3), 743-765, doi:10.5194/os-16-743-2020.
- 49 Pausas, J. G. and S. Fernández-Muñoz, 2012: Fire regime changes in the Western Mediterranean Basin: from fuel-
50 limited to drought-driven fire regime. *Climatic Change*, **110**(1), 215-226, doi:10.1007/s10584-011-0060-6.
- 51 Payet-Burin, R., F. Bertoni, C. Davidsen and P. Bauer-Gottwein, 2018: Optimization of regional water - power systems
52 under cooling constraints and climate change. *Energy*, **155**, 484-494, doi:10.1016/j.energy.2018.05.043.
- 53 Peleg, N., E. Shamir, K. P. Georgakakos and E. Morin, 2015: A framework for assessing hydrological regime
54 sensitivity to climate change in a convective rainfall environment: a case study of two medium-sized eastern
55 Mediterranean catchments, Israel. *Hydrology and Earth System Sciences*, **19**(1), 567-581, doi:10.5194/hess-19-
56 567-2015.
- 57 Peñuelas, J. et al., 2017: Impacts of global change on Mediterranean forests and their services. *Forests*, **8**(12), 463,
58 doi:10.3390/f8120463.
- 59 Peñuelas, J. et al., 2018: Assessment of the impacts of climate change on Mediterranean terrestrial ecosystems based on
60 data from field experiments and long-term monitored field gradients in Catalonia. *Environmental and*
61 *Experimental Botany*, **152**, 49-59, doi:10.1016/j.envexpbot.2017.05.012.
- 62 Pinaras, V., C. Paraskevas and A. Panagopoulos, 2021: Investigating the effects of agricultural water management in a
63 Mediterranean coastal aquifer under current and projected climate conditions. *Water*, **13**(1), 108.

- 1 Polemio, M. and T. Lonigro, 2015: Trends in climate, short-duration rainfall, and damaging hydrogeological events
2 (Apulia, Southern Italy). *Natural Hazards*, **75**(1), 515-540, doi:10.1007/s11069-014-1333-y.
- 3 Pumo, D., D. Caracciolo, F. Viola and L. V. Noto, 2016: Climate change effects on the hydrological regime of small
4 non-perennial river basins. *Science of The Total Environment*, **542**, 76-92, doi:10.1016/j.scitotenv.2015.10.109.
- 5 Pyrgou, A. and M. Santamouris, 2018: Increasing probability of heat-related mortality in a Mediterranean city due to
6 urban warming. *International Journal of Environmental Research and Public Health*, **15**(8), 1571.
- 7 Quintana-Seguí, P. et al., 2016: Drought: observed trends, future projections. In: *The Mediterranean Region under*
8 *Climate Change: A Scientific Update* [Thiébaud, S., J.-P. Moatti, V. Ducrocq, E. Gaume, F. Dulac, E. Hamonou,
9 Y.-J. Shin, J. Guiot, W. Cramer, G. Boulet, J.-F. Guégan, R. Barouki, I. Annesi-Maesano, P. Marty, E.
10 Torquebiau, J. F. Soussana, Y. Aumeeruddy-Thomas, J.-L. Chotte and D. Lacroix (eds.)]. IRD Éditions,
11 Marseille, France, pp. 123-131. ISBN 9782709922203.
- 12 Raucoules, D. et al., 2008: Ground deformation detection of the greater area of Thessaloniki (Northern Greece) using
13 radar interferometry techniques. *Natural Hazards and Earth System Sciences*, **8**(4), 779-788, doi:10.5194/nhess-8-
14 779-2008.
- 15 Raybaud, V., M. Bacha, R. Amara and G. Beaugrand, 2017: Forecasting climate-driven changes in the geographical
16 range of the European anchovy (*Engraulis encrasicolus*). *ICES Journal of Marine Science*, **74**(5), 1288-1299,
17 doi:10.1093/icesjms/fsx003.
- 18 Reimann, L. et al., 2018: Mediterranean UNESCO World Heritage at risk from coastal flooding and erosion due to sea-
19 level rise. *Nature Communications*, **9**(1), 4161, doi:10.1038/s41467-018-06645-9.
- 20 Ribes, A. et al., 2019: Observed increase in extreme daily rainfall in the French Mediterranean. *Climate Dynamics*,
21 **52**(1), 1095-1114, doi:10.1007/s00382-018-4179-2.
- 22 Rodríguez, A. et al., 2019: Chilling accumulation in fruit trees in Spain under climate change. *Natural Hazards and*
23 *Earth System Sciences*, **19**(5), 1087-1103, doi:10.5194/nhess-19-1087-2019.
- 24 Rohat, G. et al., 2019: Influence of changes in socioeconomic and climatic conditions on future heat-related health
25 challenges in Europe. *Global and Planetary Change*, **172**, 45-59, doi:10.1016/j.gloplacha.2018.09.013.
- 26 Roudier, P. et al., 2016: Projections of future floods and hydrological droughts in Europe under a +2°C global warming.
27 *Climatic Change*, **135**(2), 341-355, doi:10.1007/s10584-015-1570-4.
- 28 Ruffault, J. et al., 2018: Extreme wildfire events are linked to global-change-type droughts in the northern
29 Mediterranean. *Nat. Hazards Earth Syst. Sci.*, **18**(3), 847-856, doi:10.5194/nhess-18-847-2018.
- 30 Ruffault, J., N. K. Martin-StPaul, S. Rambal and F. Mouillot, 2013: Differential regional responses in drought length,
31 intensity and timing to recent climate changes in a Mediterranean forested ecosystem. *Climatic Change*, **117**(1),
32 103-117, doi:10.1007/s10584-012-0559-5.
- 33 Saadi, S. et al., 2015: Climate change and Mediterranean agriculture: Impacts on winter wheat and tomato crop
34 evapotranspiration, irrigation requirements and yield. *Agricultural Water Management*, **147**, 103-115,
35 doi:10.1016/j.agwat.2014.05.008.
- 36 Sabatés, A., P. Martín, J. Lloret and V. Raya, 2006: Sea warming and fish distribution: the case of the small pelagic
37 fish, *Sardinella aurita*, in the western Mediterranean. *Global Change Biology*, **12**(11), 2209-2219,
38 doi:10.1111/j.1365-2486.2006.01246.x.
- 39 Sabatés, A., P. Martín and V. Raya, 2012: Changes in life-history traits in relation to climate change: bluefish
40 (*Pomatomus saltatrix*) in the northwestern Mediterranean. *ICES Journal of Marine Science*, **69**(6), 1000-1009,
41 doi:10.1093/icesjms/fss053.
- 42 Sachs, J. et al., 2019: *Sustainable Development Report 2019*. Bertelsmann Stiftung and Sustainable Development
43 Solutions Network (SDSN), New York, NY, USA. Available at: [https://sdgindex.org/reports/sustainable-](https://sdgindex.org/reports/sustainable-development-report-2019/)
44 [development-report-2019/](https://sdgindex.org/reports/sustainable-development-report-2019/) (accessed 16/10/2020).
- 45 Salameh, A. A. M. et al., 2019: Spatio-temporal analysis for extreme temperature indices over the Levant region.
46 *International Journal of Climatology*, **39**(15), 5556-5582, doi:10.1002/joc.6171.
- 47 Salvati, A., H. Coch Roura and C. Cecere, 2017: Assessing the urban heat island and its energy impact on residential
48 buildings in Mediterranean climate: Barcelona case study. *Energy and Buildings*, **146**, 38-54,
49 doi:10.1016/j.enbuild.2017.04.025.
- 50 Sánchez-Arcilla, A. et al., 2016: A review of potential physical impacts on harbours in the Mediterranean Sea under
51 climate change. *Regional Environmental Change*, **16**(8), 2471-2484, doi:10.1007/s10113-016-0972-9.
- 52 Sánchez-Salguero, R. et al., 2020: Shifts in growth responses to climate and exceeded drought-vulnerability thresholds
53 characterize dieback in two Mediterranean deciduous oaks. *Forests*, **11**(7), 714.
- 54 Santiago, J. M. et al., 2016: Brown trout thermal niche and climate change: expected changes in the distribution of cold-
55 water fish in central Spain. *Ecology*, **9**(3), 514-528, doi:<https://doi.org/10.1002/eco.1653>.
- 56 Satir, O., S. Berberoglu and A. Cilek, 2016: Modelling long term forest fire risk using fire weather index under climate
57 change in Turkey. *Applied Ecology and Environmental Research* **14**(4), 537-551,
58 doi:10.15666/aeer/1404_537551.
- 59 Satta, A., M. Puddu, S. Venturini and C. Giupponi, 2017: Assessment of coastal risks to climate change related impacts
60 at the regional scale: The case of the Mediterranean region. *International Journal of Disaster Risk Reduction*, **24**,
61 284-296, doi:10.1016/j.ijdrr.2017.06.018.

- 1 Sayol, J. M. and M. Marcos, 2018: Assessing flood risk under sea level rise and extreme sea levels scenarios:
2 Application to the Ebro delta (Spain). *Journal of Geophysical Research: Oceans*, **123**(2), 794-811,
3 doi:10.1002/2017jc013355.
- 4 Schlessner, C. F. et al., 2016: Differential climate impacts for policy-relevant limits to global warming: the case of
5 1.5 °C and 2°C. *Earth System Dynamics*, **7**(2), 327-351, doi:10.5194/esd-7-327-2016.
- 6 Seif-Ennasr, M. et al., 2016: Climate change and adaptive water management measures in Chtouka Aït Baha region
7 (Morocco). *Science of The Total Environment*, **573**, 862-875, doi:10.1016/j.scitotenv.2016.08.170.
- 8 Sellami, H., S. Benabdallah, I. La Jeunesse and M. Vanclooster, 2016: Quantifying hydrological responses of small
9 Mediterranean catchments under climate change projections. *Science of The Total Environment*, **543**, 924-936,
10 doi:10.1016/j.scitotenv.2015.07.006.
- 11 Semenza, J. C. et al., 2016: Climate change projections of West Nile virus infections in Europe: implications for blood
12 safety practices. *Environmental Health*, **15**(1), S28, doi:10.1186/s12940-016-0105-4.
- 13 Siam, M. S. and E. A. B. Eltahir, 2017: Climate change enhances interannual variability of the Nile river flow. *Nature*
14 *Climate Change*, **7**(5), 350-354, doi:10.1038/nclimate3273.
- 15 Sierra, J. P. et al., 2016: Vulnerability of Catalan (NW Mediterranean) ports to wave overtopping due to different
16 scenarios of sea level rise. *Regional Environmental Change*, **16**(5), 1457-1468, doi:10.1007/s10113-015-0879-x.
- 17 Sierra, J. P. et al., 2017: Modelling the impact of climate change on harbour operability: The Barcelona port case study.
18 *Ocean Engineering*, **141**, 64-78, doi:10.1016/j.oceaneng.2017.06.002.
- 19 Snoussi, M., T. Ouchani and S. Niazi, 2008: Vulnerability assessment of the impact of sea-level rise and flooding on the
20 Moroccan coast: The case of the Mediterranean eastern zone. *Estuarine, Coastal and Shelf Science*, **77**(2), 206-
21 213, doi:<https://doi.org/10.1016/j.ecss.2007.09.024>.
- 22 Solaun, K. and E. Cerdá, 2017: The impact of climate change on the generation of hydroelectric power - A case study in
23 Southern Spain. *Energies*, **10**(9), 1343, doi:10.3390/en10091343.
- 24 Spinoni, J., G. Naumann and J. V. Vogt, 2017: Pan-European seasonal trends and recent changes of drought frequency
25 and severity. *Global and Planetary Change*, **148**, 113-130, doi:10.1016/j.gloplacha.2016.11.013.
- 26 Springmann, M. et al., 2016: Global and regional health effects of future food production under climate change: a
27 modelling study. *The Lancet*, **387**(10031), 1937-1946, doi:10.1016/S0140-6736(15)01156-3.
- 28 Stefanescu, C., J. Carnicer and J. Peñuelas, 2011: Determinants of species richness in generalist and specialist
29 Mediterranean butterflies: the negative synergistic forces of climate and habitat change. *Ecography*, **34**(3), 353-
30 363, doi:10.1111/j.1600-0587.2010.06264.x.
- 31 Stergiou, K. I. et al., 2016: Trends in productivity and biomass yields in the Mediterranean Sea Large Marine
32 Ecosystem during climate change. *Environmental Development*, **17**, 57-74, doi:10.1016/j.envdev.2015.09.001.
- 33 Thiéblemont, R. et al., 2019: Likely and high-end impacts of regional sea-level rise on the shoreline change of
34 European sandy coasts under a high greenhouse gas emissions scenario. *Water*, **11**(12), 2607.
- 35 Tobin, I. et al., 2018: Vulnerabilities and resilience of European power generation to 1.5 degrees C, 2 degrees C and 3
36 degrees C warming. *Environmental Research Letters*, **13**(4), doi:10.1088/1748-9326/aab211.
- 37 Trambly, Y., L. Jarlan, L. Hanich and S. Somot, 2018: Future scenarios of surface water resources availability in North
38 African dams. *Water Resources Management*, **32**(4), 1291-1306, doi:10.1007/s11269-017-1870-8.
- 39 Trambly, Y. et al., 2019: Detection and attribution of flood trends in Mediterranean basins. *Hydrology and Earth*
40 *System Sciences*, **23**(11), 4419-4431, doi:10.5194/hess-23-4419-2019.
- 41 Tsikliras, A. C. and K. I. Stergiou, 2014: Mean temperature of the catch increases quickly in the Mediterranean Sea.
42 *Mar Ecol Prog Ser*, **515**, 281-284, doi:10.3354/meps11005.
- 43 Turco, M. et al., 2016: Decreasing fires in Mediterranean Europe. *PLoS ONE*, **11**(3), e0150663,
44 doi:10.1371/journal.pone.0150663.
- 45 Turco, M., N. Levin, N. Tessler and H. Saaroni, 2017a: Recent changes and relations among drought, vegetation and
46 wildfires in the Eastern Mediterranean: The case of Israel. *Global and Planetary Change*, **151**, 28-35,
47 doi:10.1016/j.gloplacha.2016.09.002.
- 48 Turco, M., M.-C. Llasat, J. von Hardenberg and A. Provenzale, 2014: Climate change impacts on wildfires in a
49 Mediterranean environment. *Climatic Change*, **125**(3), 369-380, doi:10.1007/s10584-014-1183-3.
- 50 Turco, M. et al., 2018: Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-
51 stationary climate-fire models. *Nature Communications*, **9**(1), 3821, doi:10.1038/s41467-018-06358-z.
- 52 Turco, M. et al., 2017b: On the key role of droughts in the dynamics of summer fires in Mediterranean Europe.
53 *Scientific Reports*, **7**(1), 81, doi:10.1038/s41598-017-00116-9.
- 54 Turner, S. W. D., J. Y. Ng and S. Galelli, 2017: Examining global electricity supply vulnerability to climate change
55 using a high-fidelity hydropower dam model. *Science of The Total Environment*, **590-591**, 663-675,
56 doi:10.1016/j.scitotenv.2017.03.022.
- 57 Van Vliet, M. T. H. et al., 2016: Multi-model assessment of global hydropower and cooling water discharge potential
58 under climate change. *Global Environmental Change-Human and Policy Dimensions*, **40**, 156-170,
59 doi:10.1016/j.gloenvcha.2016.07.007.
- 60 Varela, V. et al., 2019: Projection of forest fire danger due to climate change in the French Mediterranean region.
61 *Sustainability*, **11**(16), 4284, doi:10.3390/su11164284.
- 62 Vicente-Serrano, S. M. et al., 2014: Evidence of increasing drought severity caused by temperature rise in southern
63 Europe. *Environmental Research Letters*, **9**(4), 044001, doi:10.1088/1748-9326/9/4/044001.

- 1 Vicente-Serrano, S. M. et al., 2019: Climate, irrigation, and land cover change explain streamflow trends in countries
2 bordering the Northeast Atlantic. *Geophysical Research Letters*, **46**(19), 10821-10833,
3 doi:10.1029/2019GL084084.
- 4 Wu, M. et al., 2015: Sensitivity of burned area in Europe to climate change, atmospheric CO₂ levels, and demography:
5 A comparison of two fire-vegetation models. *Journal of Geophysical Research: Biogeosciences*, **120**(11), 2256-
6 2272, doi:10.1002/2015JG003036.
- 7 Zabalza-Martínez, J. et al., 2018: The influence of climate and land-cover scenarios on dam management strategies in a
8 high water pressure catchment in Northeast Spain. *Water*, **10**(11), 1668, doi:10.3390/w10111668.
- 9 Zhang, X. et al., 2018: Impacts of climate change, policy and Water-Energy-Food nexus on hydropower development.
10 *Renewable Energy*, **116**, 827-834, doi:10.1016/j.renene.2017.10.030.
- 11 Zhao, G. et al., 2015: The implication of irrigation in climate change impact assessment: a European-wide study. *Global
12 Change Biology*, **21**(11), 4031-4048, doi:10.1111/gcb.13008.
- 13 Zinzi, M. and E. Carnielo, 2017: Impact of urban temperatures on energy performance and thermal comfort in
14 residential buildings. The case of Rome, Italy. *Energy and Buildings*, **157**, 20-29,
15 doi:10.1016/j.enbuild.2017.05.021.
- 16 Ziv, B. et al., 2014: Trends in rainfall regime over Israel, 1975–2010, and their relationship to large-scale variability.
17 *Regional Environmental Change*, **14**(5), 1751-1764, doi:10.1007/s10113-013-0414-x.
18

ACCEPTED VERSION
SUBJECT TO FINAL EDITS